# PRELIMINARY SURFICIAL GEOLOGY OF THE MIAMI COUNTY PORTION OF THE OSAWATOMIE QUADRANGLE, KANSAS

# Geology by Rolfe D. Mandel and K. David Newell

# 2014

Computer compilation and cartography by John W. Dunham, Charity M. Lander, Dustin A. Fross, and Hillary C. Crabb

**Open-file Report 2014-11** 

Funded in part by the **USGS National Cooperative Geologic Mapping Program** 



Elevation contours are presented for general reference. Used in the U.S. Geological Survey's current US Topo 1:24,000-scale topographic map series, they were generated from hydrographically-improved 1/3 arc-second National Elevation Dataset (NED) data, and smoothed during processing for use at 1:24,000 scale. In some places the contours may be more generalized than the base data used for compilation of geologic outcrop patterns. Outcrop patterns on the map will typically reflect topographic variation more accurately than the associated contour lines. Repeated fluctuation of an outcrop line across a contour line should be interpreted as an indication that the mapped rock unit is maintaining a relatively constant elevation along a generalized contour.


Geology was mapped in the field using a USGS 7.5-minute 1:24,000-scale topographic map. Roads and highways are shown on the base map as represented by data from the Kansas Department of Transportation (KDOT), U.S. Census Bureau, and other sources. U.S. Department of Agriculture - Farm Services Agency (USDA-FSA) National Agriculture Imagery Program (NAIP) imagery also was used to check road

Shaded relief is based on 1-meter hydroflattened bare-earth DEMs from the State of Kansas LiDAR Database. The DEM images, in ERDAS IMAGINE format, were mosaicked into a single output DEM, downsampled to 5-meter resolution, and reprojected to decimal degrees. The output DEM was then converted to a hillshade, a multidirectional shaded-relief image using angles of illumination from 0°, 225°, 270°, and 315° azimuths, each 45° above the horizon, with a 4x vertical exaggeration.

This geologic map was funded in part by the USGS National Cooperative Geologic Mapping Program, award This map was produced using the ArcGIS system developed by Esri (Environmental Systems Research Institute,

This map is a preliminary product and has had less scientific and cartographic review than the Kansas Geological Survey's M-series geologic maps. KGS does not guarantee this map to be free from errors or inaccuracies and disclaims any responsibility or liability for interpretations made from the map or decisions based thereon. SUGGESTED REFERENCE TO THE MAP

Mandel, R. D., and Newell, K. D., 2014, Preliminary surficial geology of the Miami County portion of the Osawatomie quadrangle, Kansas: Kansas Geological Survey, Open-file Report 2014-11, scale 1:24,000,



DECLINATION, 2014

R 22 E R 23 E

**EXPLANATION Boundaries and Locations** Geologic Unit Boundaries — Observed contact ——— County boundary Township/range line Hydrology and Topography ——— Section line Perennial stream Intermittent stream Transportation Water body 169 U.S. highway Elevation contour State highway (100-meter interval) ———— Local road Elevation contour

### **GEOLOGIC UNITS** CENOZOIC

**Quaternary System** 

Undifferentiated Alluvium — Undifferentiated alluvium that typically is less than 4,000 years old comprises valley fills beneath the modern floodplain of the Marais des Cygnes River and its tributaries. This alluvium consists of clay, silt, sand, and gravel, with the coarsest sediments comprising the lower 3-6 ft (0.9-1.8 m) of the valley fills. The fine-grained floodplain facies are mostly dark gray, gray, dark grayish brown, and grayish brown silty clay, and silty clay loam. These deposits can reach thicknesses greater than 50 ft (15 m) in the Marais des

Alluvial Terraces — Alluvium that typically is 4,000 to 11,000 years old comprises valley fills beneath a low terrace (T-1) of the Marais des Cygnes River and its tributaries. This alluvium consists of clay, silt, sand, and gravel, with the coarsest sediments comprising the lower 3-6 ft (0.9-1.8 m) of the terrace fills. The fine-grained alluvium is mostly brown, dark yellowish brown, and yellowish brown silty clay loam. These deposits can reach thicknesses greater than 70 ft (21 m) in the Marais des Cygnes River valley.

Quaternary and Neogene Alluvium — Clayey alluvium often interbedded with cherty gravel is common beneath high terraces and on hilltops in Miami County. The numerical age of these alluvial deposits is unknown. However, based on their position in the landscape, the deposits on the hilltops probably aggraded during the Neogene (Aber, 1998), and it is likely that the high-terrace fills date at least to the middle Pleistocene. The lower 3-6 ft (0.9-1.8 m) of the terrace fills and hilltop deposits consist of cherty gravel. The overlying fine-grained alluvium mostly consists of dark gray, gray, dark grayish brown, and pale brown silty clay. The distribution of fine-grained alluvium interbedded with cherty alluvial gravels is indicated by the Kenoma soil series, as shown on the Soil Survey of Linn and Miami Counties, Kansas (Penner, 1981). Redoximorphic features, including strong brown and reddish brown mottles and iron and manganese oxide concretions, are common in the upper 3-10 ft (0.9-3 m) of the fine-grained alluvium. Alluvial deposits comprising the fills of high terraces are typically 20-30 ft (6-9 m) thick, but the alluvial deposits on hilltops generally are less than 15 ft (4.5 m) thick. A veneer of loess that is less than 3 ft (0.9 m) thick often caps the high-terrace and hilltop alluvial deposits.

Colluvial Apron and Alluvial Fan Deposits — Deposits of clay, silt, sand, and gravel occur on footslopes and toeslopes in valley landscapes. These deposits mostly accumulated during the early and middle Holocene, although some may date back to the terminal Pleistocene, and they comprise both colluvial aprons and alluvial fans. The colluvial aprons consist of massive deposits of poorly sorted sediments that include many angular, bedrock-derived pebbles and cobbles. The alluvial fans formed where small, intermittent streams enter the Marais des Cygnes River valley and the valleys of its major tributaries. The fan deposits are stratified and consist of wellsorted alluvium dominated by brown, dark yellowish brown, and yellowish brown silty clay, silty clay loam, and silt loam. Thin lenses of gravel are common, and buried soils often occur at the top of upward-fining sequences.

# Carboniferous System –Pennsylvanian Subsystem

## (descriptions from Miller, 1966)

Plattsburg Limestone/Vilas Shale — The lower part of the Lansing group is composed of the Plattsburg Limestone and overlying Vilas Shale (~35 ft; 11 m thick). The **Plattsburg Limestone** (~16 ft; 5 m) is composed of two limestone members and intervening shale member. In ascending order, these units are Merriam limestone, Hickory Creek shale, and Spring Hill limestone. The Merriam Limestone Member, averaging 3.5 ft (1 m), but dense limestone. The lower part (3.6 ft; 1.1 m) of the Merriam limestone is a massive, bluish-gray to light-gray, cross-bedded, oolitic limestone, locally with black, fossiliferous chert. It is commonly a resistant unit, forming scarps and outcrops. Coated grains and Osagia-like forms are usually present and in some places these constitute a large part of the limestone. Myalinid pelecypods and productid brachiopods can be common. The upper part of the Merriam limestone is a light- to medium-gray, fine-grained, massive, locally cherty limestone, but usually not fossiliferous. "Worm tubes" are present in some localities on the upper surface. The Hickory Creek Shale Member (locally absent to 2 ft;0.6 m) is yellowish-gray to orangish-gray, clayey, nodular, and calcareous. A thin, nodular limestone bed locally occurs near the middle of the unit. The shale is fossiliferous, commonly with crinoid remains and bryozoans. The **Spring Hill Limestone Member** (4-17 ft; 1.2-5 m) is light olive-gray to yellowish-gray, fine- to coarse-grained, thin- to medium-bedded, and sandy. It contains chert locally and may be oolitic in places at its top or conglomeratic with limonite pebbles, shale and ironstone fragments, and small abraded calcite crystals. Shale partings and thin, wavy carbonaceous streaks may be present within this member. Triticites sp. and Osagia-like forms and echinoid spines are locally abundant. A zone of large Composita sp. is commonly found near the base of the unit. The Vilas Shale (5 ft; 1.5 m, thickening westward to 30 ft; 9 m) is grayish-olive to light-gray, sandy, and blocky. In the western part of the county it contains a considerable amount of moderate yellowish-brown siltstone and orangish-gray cross-bedded sandstone, which may be up to 4 ft (1.2 m) thick. Plant remains are common in the sandy layers, but otherwise it is relatively unfossiliferous. Kansas City Group

Wyandotte Limestone/Bonner Springs Shale — The uppermost part of the Kansas City Group is composed of

the Wyandotte Limestone and the overlying Bonner Springs Shale. The Wyandotte Limestone is composed of five alternating shale and limestone members. From the base upward, these five units are the Frisbie limestone, Quindaro shale, Argentine limestone, Island Creek shale, and Farley limestone. The Wyandotte is a cuestaforming formation, ranging in thickness from about 10 to 80 ft (3 to 24 m). The shale members are present only locally in Miami County. The absence of the shale members and the similarity in the lithology of the two upper limestone members makes identification of the units difficult. The Frisbie Limestone Member (2.5-3.6 ft; 0.8-1.1 m) is a light olive-gray to light brownish-gray, fine-grained, massive limestone. Locally, thin shales (<0.5 ft; 0.15 m thick) can be present in the Frisbie. Crinoid stems and small productid brachiopods are the most common fossils and locally Osagia-like forms (coated grains) may be present. The Quindaro Shale Member (locally absent to 3.3 ft, 1 m) is a dusky-yellow to dark yellowish-orange, sandy shale. In places the lower part of this unit is composed of very dark gray shale. The Quindaro is very fossiliferous with *Heterocoelia* sp., *Dielasma* sp., and Phricodothyris sp. being the most common forms. Crinoid fragments and bryozoans are also abundant. The **Argentine Limestone Member** (~35 ft; 11 m) is probably the most persistent member of the Wyandotte. It is lithologically similar to the Farley limestone which is normally the uppermost limestone member of the Wyandotte. Locally the Farley limestone can be missing and the Argentine limestone is the uppermost member. The Argentine is a light olive-gray to grayish-orange, medium-grained, thin-bedded, locally cherty limestone, which weathers into thin fragments. The Argentine has a varied fauna with the brachiopods Composita sp., Echinaria sp., Antiquatonia sp., and Phricodothyris sp. being most common. Enteletes sp. is abundant west and south of Paola. The fusulinid *Triticites* sp. is common locally. The **Island Creek Shale Member** (~1.5 ft; 0.5 m) is not well exposed. It is a grayish-orange clayey shale containing abundant gastropods. The Farley Limestone **Member** (locally absent to ~15 ft; 4.6 m), where present, is the uppermost member of the Wyandotte Limestone. It is a light olive-gray to pinkish-gray coarse-grained, wavy, thin- to thick-bedded limestone. Locally the lower part of this unit can be oolitic. The Farley limestone contains much the same fauna as the Argentine limestone. The Bonner Springs Shale (0.9-31.5 ft; 0.3-9.6 m) is the uppermost formation of the Kansas City Group. The lower part is composed of pale olive- to light-gray sandy shale which in places is a thin-bedded micaceous siltstone. The upper part is an olive-gray to yellowish-brown clayey shale in the upper part. Varying lithologies can be present near the middle of the unit – unfossiliferous grayish-red clayey shale or medium-gray clayey shale containing carbonaceous streaks, plant rootlets, and leaf impressions, or medium-grained, medium-bedded, calcareous sandstone is found in this interval. In areas where the Bonner Springs is primarily clayey shale, scattered limestone nodules are present, and a thin, argillaceous limestone bed (~1 ft; 0.3 m) is present 2.0 ft (0.6 m) below the top of the formation. The limestone bed is very fossiliferous and contains abundant pelecypods, algae, brachiopods, gastropods, and bryozoan fragments.

**Iola Limestone/Lane Shale** — The **Iola Limestone** (10-12 ft; 3-3.7 m) is composed of two limestone members and an intervening shale member – the lower limestone is the Paola limestone, the middle shale is the Muncie Creek shale, and the upper limestone is the Raytown limestone. The **Paola Limestone Member** (absent to ~2.5 ft; 0.8 m) is composed of a massive bed of dark-gray to brownish-gray, fine-grained, dense limestone. The contact with the underlying shale is fairly smooth, but the upper surface of the unit is very irregular. Locally, iron-stained "worm tubes" extend downward 1 to 3 inches (2.5-7.6 cm) from the upper surface of the Paola. Crinoid stems and small productid brachiopods are the most abundant fossils. Coated grains and bryozoans are common. The Muncie Creek Shale Member (0.5 ft; 0.15 m) is a bluish-gray to dusky-yellow sandy shale that locally has a dark-gray, carbonaceous, fissile facies that contains small (0.4-1.2 inches: 1-2 cm) phosphatic nodules. The **Raytown Limestone Member** (5-24 ft; 1.5-7 m) is a light olive-gray to light-gray medium- to coarse-grained limestone with numerous silty shale partings and abundant vugs lined with crystalline calcite. The Raytown is medium bedded in the northern part of Miami County but becomes progressively thin bedded toward the southern part. Abundant large productid brachiopods occur in the Raytown, with Echinaria sp. and Linoproductus sp. being the most common. The Lane Shale (25 ft; 8m, increasing in thickness westward to 80 ft; 24 m) is lithologically variable. In the western half of the county is an olive-gray to dusky-yellow silty to sandy shale and thin-bedded siltstone. In the eastern half it is an olive-gray to light-gray clayey shale. Thin carbonaceous streaks 1/16- to 1/4-inch (1.6- to 6.4 mm) thick are found locally in the Lane. In east-central Miami County, a thin lightgray, clayey shale containing laminae of reddish-brown calcareous silt occurs just below the overlying Wyandotte Limestone. The Lane is unfossiliferous except for sparse plant remains, and locally, small brachiopods and crinoids are present in the upper few feet of the formation.

**Drum Limestone/Chanute Shale** — The **Drum Limestone** (1.7-8 ft; 0.5-2.4 m) consists of a single massive bed of yellowish-gray to reddish-brown fine- to medium-grained cross-bedded limestone that is uniform in lithology and weathers into thin slabs. The most characteristic feature of this unit is abundant, small, white crinoid segments scattered throughout the limestone. Other fossils are Neospirifer sp., Marginifera sp., and Composita sp., and locally the small sponge Heliospongia sp. The Chanute Shale (8- 38 ft; 2.4-12 m) varies greatly in thickness and lithology over short lateral distances. It is yellowish-brown to greenish-gray sandy to clayey shale, which locally contains sandstone in the lower and middle parts and commonly in the upper part. The Chanute Shale is a relatively unfossiliferous unit except for plant impressions found locally in the sandy layers. The Chanute Shale contains the Thayer coal (0- 0.5 ft; 0.15 m), 11 to 15 feet (3.4-4.6 m) above the base of the formation. The Thayer occurs at differing stratigraphic positions in respect to the upper formational boundary, but is absent where the Chanute is thinner than 11 ft (3.4 m).

**Dennis Limestone**/Cherryvale Shale — The **Dennis Limestone** (~32 ft; 10 m) is resistant, forming prominent scarps. It is composed of three members, which, in ascending order, are the Canville limestone, Stark shale, and Winterset limestone. The Canville Limestone Member is absent in Miami County, but may be represented locally by a thin, impure, nodular limestone. The **Stark Shale Member** (0.9-6 ft; 0.3-1.8 m) is represented by a black, fissile to very thin-bedded, carbonaceous shale in the lower part. The upper part is medium-olive to greenish-gray sandy shale and is about twice the thickness of the underlying black shale. The Stark is fossiliferous in the upper, sandy part, with chonetid brachiopods being the most common fossil. The lower part is unfossiliferous except for rare inarticulate brachiopods. The Winterset Limestone Member (~29 ft; 9 m), the uppermost member of the Dennis Limestone, is light gray to olive gray, medium grained, medium bedded, dense in the lower part and more fine grained in the middle part. The lower and middle parts of the member characteristically contain shaly partings and abundant chert. The upper part is usually separated from the lower beds by a thin bed of calcareous gray shale (~1.5 ft; 0.5 m). The upper part of the Winterset is dense, finegrained, massive limestone. It contains an abundance of dark-gray chert. Locally, 1 to 3 ft (0.3-0.9 m) of oolitic limestone can be present near the top. The upper part of the Winterset is very fossiliferous and is characterized by numerous large productid brachiopods and locally by abundant gastropods. Triticites sp. is also very abundant in the upper part and is usually found in chert. The Cherryvale Shale (~60 ft; 18 m) comprises a shale-dominated unit between the top of the Dennis Limestone and the base of the Drum Limestone. It includes five members, in ascending order: the Fontana shale, the Block limestone, the Wea shale, the Westerville limestone, and the Quivira shale. The **Fontana Shale Member** (~15 ft; 4.6 m) is greenish-gray to olive-gray sandy shale. Locally a thin, nodular limestone occurs in the lower part. It is relatively unfossiliferous except for sparse chonetid brachiopods in the lower part. The **Block Limestone Member** (~4 ft; 1.2 m), the lowermost limestone bed in the Cherryvale Shale, is a bluish-gray to olive-gray medium-grained, thin-bedded fossiliferous limestone with numerous thin, fossiliferous shale partings. Triticites sp. is the most characteristic fossil, but Marginifera sp. and Syringopora sp. also are abundant. The Wea Shale Member (~18 ft; 5.5 m) is characteristically olive-gray sandy shale that contains sandstone beds locally near the base. A thin bed of maroon shale may occur near the top. Locally, pyritiferous nodules and limonite concretions occur in the Wea, but plant impressions and carbonaceous partings can be are found at its top. In places a very thin coal occurs near the top of the member. The Westerville **Limestone Member** (absent to ~2 ft; 0.6 m) is poorly exposed, but where present, is composed of conglomeratic limestone containing abundant limonite nodules and quartz sand. It can lithologically transition to yellowish-gray marly limestone. The Quivira Shale (~ 4 ft; 1.2 m) has a lower part (0.5-1 ft; 0.15-0.3 m) that is composed of black, carbonaceous, fissile shale or maroon clayey shale. The upper part consists of olive-gray sandy shale. The Quivira is unfossiliferous, except for sparse inarticulate brachiopods and conodonts in the black fissile shale.

Swope Limestone/Galesburg Shale — The Swope Limestone (34 ft; 10 m), the lower part of this map unit, is composed of two limestone members and an intervening shale member. In ascending order, these members are: Middle Creek limestone, Hushpuckney shale, and Bethany Falls limestone. The Middle Creek Limestone **Member** (1.8-2 ft) is a medium-gray to bluish-gray, fine-grained, dense limestone that is commonly expressed in outcrop as a single massive unit. A thin shale parting can be present approximately in the middle of this unit. The **Hushpuckney Shale Member** (4-7.5 ft; 1.2-2.3 m) is composed of black fissile shale overlain by grayish to yellow sandy shale. Locally, a thin bed of light olive-gray shale can overlie the black shale. The upper part of the Hushpuckney is fossiliferous, with chonetid and productid brachiopods. The Bethany Falls Limestone Member (13-28 ft; 4-8.5 m) is a light-gray to light brownish-gray, medium-grained, medium- to thick-bedded sandy limestone containing numerous thin shale partings. In most areas the Member can be divided into two distinct parts. The lower part (~12 -16 ft; 3.7-4.9 m) contains fusulinids and the upper part (0-14 ft; 4.3 m) is cherty and locally cross-bedded. Pelletal limestone or oolites, or both, may be present at the very top of the Bethany Falls. Where the pelletal or oolitic limestone is present, the upper part locally contains vertical, tubular cavities (3 to 4 feet (0.9-1.2 m) in length and 1 to 2 inches (2.5-10.2 cm) in diameter) containing iron-stained crystalline calcite. The Bethany Falls is a fossiliferous unit in which Meekella sp., Derbyia sp., Antiquatonia sp., and chonetid brachiopods are most common. Triticites sp. are found locally in the lower part of the unit. Algae are locally abundant. The Galesburg Shale (4-12 ft; 1.2-3.7 m), which overlies the Swope Limestone, consists of greenishgray to dusky-yellow, sandy to calcareous, blocky shale. It is characteristically sparsely fossiliferous and poorly

Aber, J. S., 1998, Chert Gravel and Neogene Drainage in East-Central Kansas: Current Research in Earth Sciences, Bulletin 240, part 3, Kansas Geological Survey, pp. 29-41. Miller, D. E., 1966, Geology and ground-water resources of Miami County, Kansas: Kansas Geological Survey, Bulletin 181, 66 p.; http://www.kgs.ku.edu/ General/Geology/Miami/ index.html.

Penner, H. L., 1981, Soil Survey of Linn and Miami Counties, Kansas: U.S. Department of Agriculture, Soil

Conservation Service, U.S. Government Printing Office, Washington, D.C., 102 p.

M-4	M-7/8 M-82	M-11	M-75	
1 M-5	2	3 M-13A	4 • M- M-18	17A 5
M-81	<b>M</b> -6A	<b>M</b> -12	RWD#2	
	M-38	M-30		
6	• M-36 7	8 M-32 ●	9 M-24	10
M-45		M-57	M-74	
11 M-48	M-51 M-52 12 M-54B M-53A	13 • <sup>M-60</sup>	M-62A M-79 14 M-70	15

# MIAMI COUNTY QUADRANGLES AND DRILLHOLE LOCATIONS

1 Wellsville

2 Antioch 3 Spring Hill 4 Bucyrus 5 West Line

(20-meter interval)

MO-KS 6 Rantoul 7 Paola West 8 Paola East

9 Louisburg 10 Freeman 11 Lane 12 Osawatomie 13 Fontana

14 New Lancaster 15 Drexel

Latitude Longitude N38.632° W94.671° N38.721° W95.018° M-5 N38.68° W95.004° M-6A N38.637° W94.987° N38.737° W94.92° N38.736° W94.796° M-12 N38.651° W94.804° M-13A N38.688° W94.758° M-17A N38.688° W94.637° N38.666° W94.714° N38.545° W94.638° N38.606° N38.535° W94.809° M-36 N38.574° W94.936° N38.599° W94.991° M-45 N38.498° W95.055° M-48 W95.051° N38.42° M-51 N38.456° W94.973° N38.445° M-53A N38.396° W94.947° M-54B N38.408° W94.991° W94.78° M-60 N38.412° W94.831° M-62A W94.713° N38.448° M-70 N38.419° W94.701° N38.478° N38.737° W94.703° M-76 N38.543° W94.759° N38.571° W94.656° M-78 N38.565° W94.993°

W94.736°

W95.048°

W95.002°

M-79


KANSAS

QUADRANGLE LOCATION

N38.425°

N38.671°

N38.721°



KANSAS DEPARTMENT OF TRANSPORTATION (KDOT) M-32 CORE

In 2002, the Kansas Department of Transportation (KDOT) began a project with the Kansas Geological Survey (KGS) to investigate the construction aggregate properties of limestones of the Pennsylvanian Kansas City Group in Miami County. That project involved drillcore donations with over 2,700 linear feet of core now housed at the KGS. That older project documented engineering properties of drillcore samples, but improved stratigraphic correlations were needed. The KGS is subcontracting with former KGS postdoctoral researcher Dr. Bradley Cramer of the University of Iowa to refine stratigraphic interpretations of Pennsylvanian bedrock units in Miami County. The work performed under the subcontract includes description and analysis of a series of 33 drill cores from throughout Miami County (see Table 1 and labels on index map). Descriptions of the cores include detailed cmscale core logging and the production of high-resolution stratigraphic columns for each core. Analysis of the core suite will eventually include two primary data sets: carbon isotope chemostratigraphy and micropaleontological biostratigraphy using conodonts. These two data sets will be combined as bio-chemostratigraphy and integrated into a comprehensive synthesis of the chronostratigraphic correlation of the units contained within the cores. This synthesis will directly inform the surficial mapping and geospatial understanding of both the surface and shallow subsurface geology of Miami County.