

PRELIMINARY SURFICIAL GEOLOGY OF THE SUMNER COUNTY PORTION OF THE ZYBA QUADRANGLE, KANSAS

by Anthony L. Layzell and John W. Dunham
2020

Open-File Report 2020-6

Funded in part by the
USGS National Cooperative
Geologic Mapping Program

The following descriptions consist of a compilation of several sources, including field notes and measured sections, data from shallow (<10 m [33 ft]) cores collected during the course of mapping, U.S. Department of Agriculture Natural Resources Conservation Service databases, and lithologic and other data in Swineford (1955) and Walters (1961).

GEOLOGIC UNITS

CENOZOIC

Quaternary System Pleistocene-Holocene

Undifferentiated floodplain alluvium—Floodplain deposits are associated with most of the principal streams in Sumner County, including the Arkansas, Nineschah, and Chikaskia rivers. Most streams in the county have undergone deep incision during the latest Holocene and have recently re-established floodplains within the incised channels in some areas. Hence, floodplain deposits are likely less than 1,000 years old. Floodplain alluvium is composed of silt, sand, and gravel and is relatively thin, probably not exceeding 20 ft (6 m) in thickness.

Alluvial terrace deposits—Stream terrace deposits are present in all major stream valleys in the county. Radiocarbon and luminescence ages indicate that terrace deposits are late Pleistocene to Holocene in age (30,000–1,500 yr B.P.). Terrace deposits are composed of clay, silt, sand, and gravel and likely do not exceed 70 ft (21 m) in thickness. Terrace alluvium in the principal stream valleys tends to be coarser grained, being chiefly derived from upland alluvial deposits (Qal₂) in the county as well as sediments comprising the High Plains succession in adjacent western counties (e.g., Kingman and Harper). Terrace deposits in smaller stream valleys are primarily derived from local shale units of the Permian Sumner Group and therefore tend to be finer grained.

Upland alluvium—Upland alluvial deposits are extensive in Sumner County and are found well above up to 65 ft (20 m) and often higher in the county. Upland alluvium is composed chiefly of silt, sand, and gravel. These deposits are reddish brown in color and coarser sediments are typically arkosic. Upland alluvium may be up to 90 ft (27 m) thick.

Dune sand—Isolated eolian sand deposits occur primarily on floodplain (Qal₁) and terrace (Qt) surfaces in Sumner County. These deposits are associated with major streams (e.g., the Arkansas and Chikaskia rivers) as they provide the source for the windblown sediments. Luminescence dating of the Hutchinson dunes, which comprise alluvial deposits of the Arkansas River in Reno County, indicate that eolian sand deposition has most recently occurred episodically over the last 2,100 years (Haffen et al., 2012). Dune sand thickness is highly variable but is generally less than 30 ft (9 m) thick.

PALEOZOIC

Permian System Leonardian Series

Wellington Formation—The Wellington Formation ranges from 40 to 65 ft (12 to 198 m) in thickness and consists principally of gray to greenish-gray shale with minor amounts of limestone, dolomite, silstone, gypsum, and anhydrite, representing marine, brackish, and freshwater environments (Swineford, 1955; Walters, 1961). The Wellington Formation contains the following formally recognized members, in ascending order: the Hollenberg Limestone, the Carlton Limestone, the Hutchinson Salt, and the Milan Limestone, as well as two unnamed members (Sawin et al., 2008). The Wellington Formation is bounded by the Nolans Limestone below and the Nineschah Shale above. Throughout the county, the formation either crops out or is covered unconformably by Quaternary deposits.

The Hollenberg Limestone is a persistent bed of argillaceous, dolomitic limestone and comprises the lower 40 ft (12 m) of the formation. The Carlton Limestone primarily consists of gray-green shales containing numerous discontinuous thin (<0.5 ft [15 cm]) beds of limestone and can be up to 60 ft (18 m) thick. The middle part of the Wellington Formation is composed of the Hutchinson Salt Member. No outcrops of this member are present in the county because of its high solubility. The Hutchinson Salt Member is thickest in the northwest part of the county but likely does not exceed 150 ft (45 m). A distinct color change from gray-green to red and purplish-red shale occurs in the upper part of the Wellington Formation. The Milan Limestone marks the top of the Wellington Formation and may be as much as 8 ft (2.5 m) thick. It consists of one to three thin beds of dolomitic limestone containing bright-green copper carbonate. In many places in Sumner County, the Milan cannot be recognized; hence the contact between the Wellington Formation and the overlying Nineschah Shale is indefinite.

CITED REFERENCES


- Haffen, A. F., Johnson, W. C., Hanson, P. R., Woodburn, T. L., Ludvigson, G. A., and Young, A. R., 2012, Activation history of the Hutchinson dunes in east-central Kansas, USA during the past 2200 years: *Aeolian Research* 5, p. 9–20.
- Sawin, R. S., Fransen, E. K., West, R. R., Ludvigson, G. A., and Watney, W. L., 2008, Clarification and changes in Permian stratigraphic nomenclature in Kansas, in *Current Research in Earth Sciences: Kansas Geological Survey, Bulletin 254*, part 2. <http://www.kgs.ku.edu/Current/2008/Sawin/index.html>.
- Swineford, A., 1955, Petrography of Upper Permian rocks in south-central Kansas: *Kansas Geological Survey, Bulletin 111*, 179 p.
- Walters, K. L., 1961, Geology and ground-water resources of Sumner County, Kansas: *Kansas Geological Survey, Bulletin 151*, 198 p.

ADDITIONAL SOURCE


- Bayne, C. K., 1962, Geology and ground-water resources of Cowley County, Kansas: *Kansas Geological Survey, Bulletin 158*, 219 p.

EXPLANATION

Boundaries and Locations	Hydrology
County boundary	Perennial stream
Township/range line	Intermittent stream
Section line	Water body
Transportation	Water body – manmade shoreline
U.S. highway	Topography
Local road	Elevation contour (50-foot interval)
Railroad	Elevation contour (10-foot interval)
Geologic Unit Boundaries	
Observed contact	



Lineaments were mapped using aerial photography and bare-earth LIDAR Digital Elevation Models (DEMs), based on one or more of the following criteria: (1) Visible offset in marker beds; (2) relatively low- to high-relief linear ridges that are not obvious geomorphic features due to erosion, slumping, subsidence, or dune formation; (3) linear ridges that cross drainage divides; (4) ridges that display curvilinear or backstepping (en echelon) configurations; (5) rectilinear or parallel drainage patterns; and (6) linear drainage patterns that align across drainage divides.



Elevation contours are presented for general reference. Used in the U.S. Geological Survey's current US Topo 1:24,000-scale topographic map series, they were generated from hydrographically improved 1/3 arc-second National Elevation Dataset (NED) data and smoothed during processing for use at 1:24,000 scale. In some places, the contours may be more generalized than the base data used for compilation of geologic outcrop patterns. Outcrop patterns on the map will typically reflect topographic variation more accurately than the associated contour lines. Repeated fluctuation of an outcrop line across a contour line should be interpreted as an indication that the mapped rock unit is maintaining a relatively constant elevation along a generalized contour.

1-meter LIDAR hillshades (2010 imagery) and 1-foot Kansas NG911 2014 digital imagery were used as references in the digital mapping. USDA Natural Resources Conservation Service (NRCS) SSURGO data and other geologic maps and bulletins were used to supplement the mapping. Field mapping was undertaken from October 2019 to February 2020. Roads and highways are shown on the base map as represented by data from the U.S. Census Bureau, U.S. Department of Agriculture – Farm Services Agency (USDA-FSA) National Agriculture Imagery Program (NAIP) imagery also was used to check road locations.

Shaded relief is based on 1-meter hydroflattened bare-earth DEMs from the State of Kansas LIDAR Database. The DEM images, in ERDAS IMAGINE format, were mosaicked into a single output DEM, downsampled to 2-meter resolution, and projected to decimal degrees. The output DEM was then converted to a hillshade, a multidirectional shaded-relief image using angles of illumination from 0°, 225°, 270°, and 315° azimuths, each 45° above the horizon, with a 4x vertical exaggeration.


This geologic map was funded in part by the USGS National Cooperative Geologic Mapping Program, award number G19AC00231 (FY2019).

This map was produced using the ArcGIS system developed by Esri (Environmental Systems Research Institute, Inc.).

This map is a preliminary product and has had less scientific and cartographic review than the Kansas Geological Survey's M-series geologic maps. KGS does not guarantee this map to be free from errors or inaccuracies and disclaims any responsibility or liability for interpretations made from the map or decisions based thereon.

SUGGESTED REFERENCE TO THE MAP

Layzell, A. L., and Dunham, J. W., 2020, Preliminary surficial geology of the Sumner County portion of the Zyba quadrangle, Kansas: *Kansas Geological Survey, Open-File Report 2020-6*, scale 1:24,000, unpublished.




UNIVERSAL TRANSVERSE MERCATOR PROJECTION, ZONE 14
NORTH AMERICAN DATUM OF 1983

37°22'

TRUE NORTH
MAGNETIC NORTH

APPROXIMATE MEAN
DECLINATION, 2020



QUADRANGLE LOCATION

Clearwater	Bayneville	Darby
Millerton	Zyba	Belle Plaine
Mayfield	Wellington	Dillon

ADJOINING 7.5' QUADRANGLES