

Figure 6--Occurrence of fossil mollusks in Nebraskan deposits at 11 localities in Kansas. Asterisk in location info indicates collection from test hole samples.

Molluscan species	Faunal localities										
	SE sec. 6, T. 2 S., R. 20 E., Doniphan	SW sec. 12, T. 29 S., R. 8 W., Kingman	*SW sec. 31, T. 30 S., R. 26 W., Meade	*SW sec. 34, T. 30 S., R. 28 W., Meade	*SW sec. 33, T. 30 S., R. 29 W., Meade	*NW sec. 6, T. 31 S., R. 30 W., Meade	NW sec. 19, T. 32 S., R. 28 W., Meade	SW sec. 22, T. 33 S., R. 29 W., Meade	*NE sec. 33, T. 34 S., R. 29 W., Meade	SW sec. 35, T. 34 S., R. 30 W., Meade	SW sec. 36, T. 34 S., R. 31 W., Seward
Nebraskan or older to Recent											
<i>Cionella lubrica</i> (Muller)							●	●			
<i>Ferrissia parallela</i> (Haldeman)					●		●	●		●	
<i>Ferrissia rivularis</i> (Say)		●					●	●		●	
<i>Gastrocopta holzingeri</i> (Sterki)							●	●			●
<i>Gastrocopta tappaniana</i> (C. B. Adams)		●		●		●	●	●	●	●	●
<i>Hawaiia minuscula</i> (Binney)			●	●	●		●	●	●	●	●
<i>Heliodiscus singleyanus</i> (Pilsbry)		●				●	●	●		●	
<i>Helisoma antrosa</i> (Conrad)	●		●				●	●		●	●
<i>Lymnaea humilis modicella</i> (Say)		●					●	●	●		●
<i>Physa anatina</i> Leo	●	●			●		●	●		●	●
<i>Pisidium</i> sp.		●					●	●		●	
<i>Pupoides aibilabris</i> (C. B. Adams)							●	●	●	●	
<i>Retinella electrina</i> (Gould)		●						●		●	
<i>Sphaerium</i> sp.	●	●			●		●	●			
<i>Strobilops cf. labyrinthica</i>							●	●			●
<i>Succinea grosvenori</i> Lea							●	●			●
<i>Vallonia gracilicosta</i> Reinhardt				●			●	●			
<i>Vertigo milium</i> Gould				●			●	●		●	●
<i>Zonitoides arboreus</i> (Say)		●						●			

Molluscan species	Faunal localities											
	SE sec. 6, T. 2 S., R. 20 E., Doniphan	SW sec. 12, T. 29 S., R. 8 W., Kingman	*SW sec. 31, T. 30 S., R. 26 W., Meade	*SW sec. 34, T. 30 S., R. 28 W., Meade	*SW sec. 33, T. 30 S., R. 29 W., Meade	*NW sec. 6, T. 31 S., R. 30 W., Meade	NW sec. 19, T. 32 S., R. 28 W., Meade	SW sec. 22, T. 33 S., R. 29 W., Meade	*NE sec. 33, T. 34 S., R. 29 W., Meade	SW sec. 35, T. 34 S., R. 30 W., Meade	SW sec. 36, T. 34 S., R. 31 W., Seward	
Nebraskan to Yarmouthian												
<i>Carychium peregrinum</i> Baker		●				●	●		●	●	●	
<i>Deroceras aenigma</i> Leonard			●				●	●		●	●	●
<i>Strobilops sparsicosta</i> Baker							●	●		●	●	
Restricted to Nebraskan												
<i>Amnicola crybates</i> Leonard	●	●	●	●	●	●	●	●	●	●	●	●
<i>Gastrocopta paracristata</i> Franzen and Leonard			●	●	●	●	●	●		●	●	●
<i>Gastrocopta rexroadensis</i> Franzen and Leonard								●	●		●	●
<i>Gyraulus enaulus</i> Leonard		●	●	●			●		●			
<i>Lymnaea diminuta</i> Leonard	●	●		●		●		●		●	●	●
<i>Lymnaea macella</i> Leonard		●				●		●				●
<i>Lymnaea perexilis</i> Leonard	●	●					●	●		●	●	
<i>Lymnaea turritella</i> Leonard		●				●	●	●		●	●	
<i>Menetus kansasensis</i> Baker	●	●		●				●		●	●	
<i>Polygyra mooreana</i> (Binney)									●		●	
<i>Promenetus blancoensis</i> Leonard	●	●				●		●		●	●	
<i>Vertigo hibbardi</i> Baker		●				●		●	●	●	●	

Pleistocene Geology of Kansas, by John C. Frye and A. Byron Leonard, Originally published in 1952 as Kansas Geological Survey Bulletin 99.

<http://www.kgs.ku.edu/Publications/Bulletins/99/index.html>

Aug. 22, 2005