Figure 2—Correlations and stratigraphic nomenclature for Southwestern Kansas.

	Hibbard, 1949	Frye and Leonard, 1952	Hibbard, 1956 Kansas General	Hibbard, 1956 Southwestern Kansas	Zeller, 1968 Central and Western Kansas	Skinner-Hibbard, 1972	Bayne and Others, 1971 Substage	Bayne and Others, 1971 Rock Unit	This Report, 1974
Wisconsinan	Vanhem formation	Sanborn formation: Bignell member; sand and gravel; Brady soil; Peoria member; sand and gravel	Sanborn group: Bignell formation; terrace deposits; Brady soil; Peoria formation; terrace deposits	Sanborn group: Vanhem formation; Jones Ranch local fauna	Bignell Formation; Fluvial deposits; Brady Soil; Peoria Formation; Fluvial deposits			Bignell Formation; Brady Soil; Peoria Formation; Gilman Canyon Formation	Vanhem formation: Roberts local fauna 11,000 yrs. B. P.; Jones Ranch l.f. 27,500 yrs. B. P.; Horse Quarry l. f. 35,000 yrs. B. P.; Jinglebob local fauna
Sangamonian	Kingsdown formation	Sangamon soil; Loveland member	Sangamon soil; Loveland formation	Jinglebob local fauna; Cragin Quarry local fauna; Kingsdown formation	Sangamon Soil; Loveland Formation			Sangamon Soil; Loveland Formation	Kingsdown formation: Cragin Quarry local fauna; Mount Scott local fauna; Butler Spring local fauna; Adams local fauna
Illinoisan		Crete member	Crete formation	Butler Spring local fauna	Crete Formation			Fluvial deposits	
Yarmouthian	Crooked Creek formation: Pearlette ash member	Meade formation: Yarmouth soil; Sappa member; Pearlette ash	Meade group: Unnamed formation: Yarmouth soil; Sappa member; Pearlette ash	Meade group: Crooked Creek formation: Soil Caliche; Borchers l. f.; Atwater member; Pearlette ash; Cudahy l. f.	Yarmouth Soil; Sappa Formation; Pearlette ash		Upper Kansan	Yarmouth Soil; Loess; Fluvial and Lacustrine deposits	Unnamed: *Kanoplis local fauna; Pearlette ash (restricted); Cudahy local fauna
Kansan	Crooked Creek formation: Stump Arroyo member	Meade formation: Grand Island member; Kansas Till; Atchison member	Meade group: Unnamed formation: Grand Island member; Kansas Till; Atchison member	Meade group: Crooked Creek formation: Stump Arroyo member; Seger local fauna	Grand Island Formation	Volcanic ash (type "S", 1.2 m.y.)	Medial Kansan	Cedar Bluffs Till; Fluvial deposits	
							Lower Kansan	Nickerson Till; Atchison Formation	
Aftonian	Meade formation: Missler member	Blanco formation: Afton soil; Fullerton member; Holdrege member	Meade group: Unnamed formation: Afton soil; Fullerton member; Holdrege member	Meade group: Ballard formation: Sanders local fauna; Caliche; Deer Park l. f.; Missler member; Dixon l. f.	Afton Soil; Fullerton Formation		Upper Nebraskan	Afton Soil; Iowa Point Till	Unnamed: *Wathena local fauna; Nash local fauna; Borchers local fauna; Borchers type "B" ash; Seger local fauna; *White Rock l. f.; *Dixon local fauna; Sanders local fauna
Nebraskan	Meade Gravels member	Nebraskan Till; David City fm.	Nebraskan Till; David City member	Angell member	Holdrege Formation	Sand Draw local fauna	Lower Nebraskan	Fluvial and Lacustrine deposits	Ballard fm.: Deer Park local fauna; Unnamed local fauna; Spring Creek l. f.; Benders local fauna