


Technical Series 7

Surficial Geology and Stratigraphy of Russell County, Kansas

Alan F. Arbogast and William C. Johnson



Kansas Geological Survey

1996

Cover: Shaded digital elevation model for Russell County, Kansas. Higher elevations are depicted in shades of yellow, while lower elevations appear in shades of green. The Smoky Hill drainage can be seen flowing from west to east across the lower part of Russell County, and the Saline River drainage, also flowing from west to east, can be seen in the north. The cover illustration was created by merging four 1:250,000 scale digital elevation models, projecting them to a Lambert Conformal Conic projection, cutting out the area of interest, and embedding the Russell County boundary. Illustration production courtesy of the Kansas Applied Remote Sensing Program, The University of Kansas, Lawrence, Kansas 66045.

Surficial geology and stratigraphy of Russell County, Kansas

Alan F. Arbogast¹ and William C. Johnson²

¹Department of Geography, Michigan State University, East Lansing, MI 48824

²Department of Geography, The University of Kansas, Lawrence, KS 66045

A text to accompany the *Geologic Map of Russell County*,
compiled by W. C. Johnson and A. F. Arbogast
(Kansas Geological Survey, Map Series M-37, 1996)

The Kansas Geological Survey compiled this publication according to specific standards, using what is thought to be the most reliable information available. The Kansas Geological Survey does not guarantee freedom from errors or inaccuracies and disclaims any legal responsibility or liability for interpretations made from the publication, or decisions based thereon.

Editor: Liz Brosius
Designer: Jennifer Sims



Text printed on recycled paper with soy-based ink

Contents

Abstract	1	Tertiary System—Miocene Series	15
Introduction	1	Ogallala Formation	15
Purpose and scope of the investigation	1	Three members	17
Location and nature of the study area	2	Quaternary System—Pleistocene Series	18
Previous geologic investigations	2	Pre-Illinoian deposits	19
Methods of investigation	4	Illinoian deposits	20
Stratigraphy of outcrops	5	Crete Formation	21
Cretaceous System—Lower Cretaceous Series	5	Loveland Loess	23
Dakota Formation	5	Sangamon Soil	24
Cretaceous System—Upper Cretaceous Series	7	Wisconsinan deposits	25
Graneros Shale	7	Gilman Canyon Formation	25
Greenhorn Limestone	7	Peoria Loess	28
Carlile Shale	11	Quaternary System—Holocene Series	30
Fairport Chalk Member	13	Brady Soil	30
Blue Hill Shale Member	13	Bignell Loess	31
Codell Sandstone Member	14	Fluvial deposits	31
Niobrara Chalk	14	Recent geologic history	37
Fort Hays Limestone Member	14	Acknowledgments	40
Smoky Hill Chalk Member	15	References	40

Tables

1. Geology and soil series associations	5	3. Radiocarbon ages from the Beisel-Steinle site	28
2. Fission-track dated volcanic ashes	19		

Figures

1. Index map of Kansas	2	23. Late Quaternary stratigraphic succession in Russell County	24
2. Drainage pattern for Russell County	3	24. Pleistocene sand and gravel	25
3. Classification of rocks in Russell County	6-7	25. Beisel-Steinle site	26
4. Exposure of the upper Dakota Formation	8	26. Upper section of the Beisel-Steinle site	27
5. Exposure of the upper Dakota Formation	9	27. Chemical, textural, and magnetic characteristics of the Beisel-Steinle site	29
6. Exposure of the Graneros Shale	10	28. $\delta^{13}\text{C}$ values from the Beisel-Steinle site	30
7. Graneros Shale—Greenhorn Limestone contact	10	29. Holocene alluvium in the Saline River valley	32
8. Hartland Shale and Jetmore Chalk Members	11	30. Channel way, floodplain, and low terraces of the Saline River	32
9. Exposure of the Greenhorn Limestone	12	31. Map of Wolf Creek basin	33
10. A small fence-post quarry	12	32. Generalized cross section of Wolf Creek basin	34
11. Sinkholes along Interstate 70	13	33. View of the T-4 terrace in Wolf Creek basin	35
12. Geologic profile of the Crawford sink	14	34. View of the T-3 terrace in Wolf Creek basin	35
13. Roadcut in the lower Fairport Chalk	15	35. View of the Naylor site in Wolf Creek basin	36
14. Blue Hill Shale landscape	16	36. View of the Paschal site in Wolf Creek basin	37
15. Slumping of the Blue Hill Shale	16	37. View of the Schoen site in Wolf Creek basin	38
16. Outcrop of the Blue Hill Shale	17	38. Aerial photograph of the T-2 terrace in Wolf Creek basin	39
17. Outcrop of the Codell Sandstone	18	39. Cross section of the T-2, T-1 complex, and floodplain in Wolf Creek basin	39
18. Gravel quarry exposing the Niobrara Chalk	20		
19. Smoky Hill Chalk Member	21		
20. Exposure of the Ogallala Formation	22		
21. Cross beds in the Ogallala Formation	22		
22. Quaternary time scale	23		

Abstract

Russell County is located in north-central Kansas, within the Smoky Hills physiographic province, which exhibits a wide variety of erosional and depositional landscapes. Outcrops range in age from Upper Cretaceous to Holocene, including the Dakota Formation, Graneros Shale, Greenhorn Limestone, Carlile Shale, Niobrara Chalk, Ogallala Formation, and several unconsolidated Quaternary eolian and fluvial deposits. The Dakota Formation, consisting of deltaic sandstones and mudstones, underlies the entire county but crops out primarily in the eastern one-half of the study area. The Graneros Shale is a thin formation of dominantly fissile shale, poorly exposed, that crops out conformably between the Dakota and the overlying Greenhorn Limestone. The Greenhorn Limestone consists of transgressive beds of shale and chalk and crops out countywide. Intensively dissected in some areas, the Greenhorn Limestone forms steep canyons where resistant beds of chalk and chalky limestone, such as the Fencepost Limestone, are present. The Carlile Shale consists of two distinct members in Russell County, the Fairport Chalk Member and the Blue Hill Shale Member. Exposures of the Fairport Chalk Member are common throughout the county but are thickest in the northwestern corner, whereas the Blue Hill Shale Member is exposed only in the northwestern corner of Russell County. Due to the paucity of exposures, the Codell Sandstone Member of the Carlile Shale is not differentiated from the Blue Hill Shale Member. Unconformably overlying the Carlile is the Niobrara Chalk, which consists of two distinct members, the Fort Hays Limestone Member and the Smoky Hill Chalk Member. Both members of the Niobrara Chalk are confined to the extreme northwestern corner of Russell County. Outcrops of the Miocene-age Ogallala Formation, a sedimentary sequence of Rocky Mountain-derived fluvial sediments, have been tentatively identified and are largely confined to the northwestern corner of the county.

Quaternary deposits, ranging in age from pre-Illinoian to Holocene, mantle much of Russell County, but are generally poorly exposed. Pre-Illinoian associations are tentative, consisting of the Grand Island and Sappa Formations, and are based on the topographic position of upland gravel deposits. Illinoian stratigraphy consists of the Crete Formation, Loveland Loess, and Sangamon Soil. Sands and gravels of the Crete Formation are expressed as a high terrace along the major streams in the county, but their distribution is uncertain. The Loveland Loess is widely distributed in north-central Kansas and has been recognized at one site in Russell County. The Sangamon Soil, a major pedostratigraphic feature which caps the Illinoian Loveland Loess, has been identified in Russell County. Overlying the Sangamon Soil is the middle-Wisconsinan Gilman Canyon Formation and the late-Wisconsinan Peoria Loess. The Gilman Canyon Formation is a loess that accumulated at a sufficiently slow rate for the development of a soil. Accelerated loess fall during the late Pleistocene produced the Peoria Loess, which mantles the upland.

Holocene stratigraphic elements include the Brady Soil, Bignell Loess, modern surface soil, and fluvial deposits. The Brady Soil caps the Peoria Loess and is discontinuous in Russell County but is recognizable only where it is buried by Bignell Loess. Holocene fluvial deposits occur as fill in a high terrace and as post-1,000-yr B.P. floodplain sediments.

Introduction

Purpose and scope of the investigation

In recent years, the Kansas Geological Survey has initiated a series of county-level mapping projects in Kansas. The primary goal of the program was to produce accurate, high-resolution maps of surficial geology in

selected counties (fig. 1). This report augments the geologic map of Russell County (Johnson and Arbogast, 1996), which was produced by the automated cartography facility of the Kansas Geological Survey. Although many advantages exist for storing the map digitally, the primary one is that the map can be easily revised without the




FIGURE 1. Index map of Kansas showing the location of Russell County and the status of county-scale geologic mapping; those counties without shading or patterns (e.g., Cheyenne) have older geologic maps.

complications associated with traditional cartography. All original project field materials (e.g., maps and aerial photography) are available in the Kansas Geological Survey archives.

Geologic mapping of Russell County was begun in the spring of 1991 and was completed in the early summer of 1992. During the spring of 1991, initial reconnaissance was conducted via remote sensing of aerial photography and field survey. Summer months were dedicated to detailed mapping of alluvium and Cretaceous bedrock, followed by mapping of Pleistocene loess in the fall. Subsequent work focused on areas with ill-defined contacts or of relative geologic complexity.

Location and nature of the study area

Russell County is located in north-central Kansas, bounded on the east by Lincoln and Ellsworth counties, on the south by Barton County, on the west by Ellis County, and on the north by Osborne County (fig. 1). It lies within the Smoky Hills physiographic province and is characterized by a fascinating diversity of landscapes, owing largely

to dissection of Cretaceous chinks, shales, and sandstones by the Smoky Hill River, Saline River, and their tributaries (fig. 2). Where extensive dissection has occurred, steep-sided valley walls are supported by resistant units (e.g., Fencepost Limestone) that are separated by less-resistant strata (e.g., Graneros Shale). Dissection of the Dakota Formation in the eastern third of the county has left a landscape of hoodoos where resistant, lenticular sandstone bodies (e.g., Rocktown channel sandstone) protect underlying mudstones. A mantle of late Quaternary loess of varying thickness caps most uplands in the county. In the larger valleys of Russell County, several cycles of alluvial cutting and filling during the Quaternary Period (Pleistocene and Holocene epochs) have resulted in deposition of gravel deposits and fine-grained alluvium which underlie terraces.

Previous geologic investigations

The first published research in Russell County was conducted by Hay (1889), who examined a lignite horizon in the Dakota Formation. Intensive research began in the




FIGURE 2. Drainage pattern for Russell County.

1920s, as exemplified by Rubey and Bass's (1925) description and map of Russell County geology. A subsurface correlation of rock units from Russell County to Marion County was completed by Bramlette (1925). In 1926, Bass included Russell County in a study of the petroleum and gas resources of western Kansas. That same year, Hedberg (1926) analyzed the effect of gravitational compaction on the structure of sedimentary rocks in and around Russell County.

Since the late 1920s, research in Russell County has largely focused on petroleum resources and the prominent Cretaceous shales and chinks of the area. In the context of petroleum production, Hintze (1928) discussed the discovery of the Gorham oil district, and Allan and Valerius (1929) studied the structure of the Fairport oil field. Chemical analyses of some oil-well waters in Russell, Ellis, and Trego counties was conducted by Runyon and Rankin (1940). In 1960, Landes et al. analyzed the

petroleum resources in basement rocks in and around Russell County. Riggs et al. (1963) conducted a petroleum engineering study of Hall-Gurney oil field in Russell County. Recently, a series of maps (e.g., Ross and Wong 1989a, 1989b, 1990a, 1990b) were constructed showing the oil and gas fields in and around Russell County.

The Cretaceous shales and chinks of north-central Kansas, including Russell County, have received considerable attention in the past 60 years. Studies of the general stratigraphy of the Colorado and Montana groups throughout north-central Kansas include those by Cobban (1949) and Hattin and Cobban (1977). A number of studies have focused on individual Upper Cretaceous formations. For example, characteristics of the Dakota Formation have been described by Franks (1967), Hattin (1967), and Siemers (1971a, 1971b). The Graneros Shale has been the focus of appreciable study. Hatfield (1961), for example, studied the paleoecology of the Graneros Shale in and around Russell County, and Hattin (1965a) described the stratigraphy of that unit. The nature of limestone beds in the Greenhorn Limestone was analyzed by Hattin (1971), who subsequently (1975) described the stratigraphy and depositional environment of the Greenhorn. The primary study of the Carlile Shale was conducted by Hattin (1962), who analyzed the stratigraphy of the Carlile in and around Russell County. The petrography, geochemistry, and economic utilization of the Fort Hays Chalk in Russell County and north-central Kansas was studied by Dubins (1947). Regarding the inclusion of all Cretaceous rocks in Russell County within the Colorado Group, Hattin (personal communication) suggests that the term Colorado Group be discontinued because the units are too lithologically diverse to be included within one group. As a result, the term Colorado Group is not used in this report.

Because the geology of north-central Kansas illicit such interest, several guidebooks have been published that include sites in Russell County. The first field guide, which focused on Pleistocene deposits, was produced by Frye et al. (1951). Guidebooks illustrating the nature of Cretaceous stratigraphy in and around Russell County include those by Hattin (1965b), Hattin and Cobban (1965), Nickel (1972), and Hattin and Siemers (1978).

In the last 15 to 20 years, interest in Russell County and the surrounding area has focused on salt dissolution and sinkhole evolution. For example, Burgat and Taylor (1972) described highway subsidence associated with salt dissolution. In 1977, Hansen studied dissolution in the Hutchinson Salt Member of the Wellington Formation near Russell. Knapp and Steeples (1981) and Knapp et al. (1989) conducted seismic investigations of collapse structures associated with salt dissolution.

Other research not cited here, such as that relating to ground water, has been conducted and reported for Russell

County. A listing is available in the *Bibliography of Kansas Geology, 1823–1984* (Sorensen et al., 1989, p. 388–389), and the *Bibliography of Kansas Geology, 1985–1989* (Sorensen, 1994, p. 147).

Methods of investigation

The geology of Russell County was mapped from stereopairs of black and white aerial photography (1:24,000) taken in January 1986 for statewide property reappraisal, 7½-minute topographic quadrangle maps, and field survey. Minimum thickness of any mappable unit was 1.8 m (6 ft), a criterion applicable primarily to loess and alluvium. Although no deep drilling or coring was done in association with the mapping, unit thicknesses beneath the solum were verified with hand and machine augering.

To assess the correlation between soil units and lithology, the Soil Survey of Russell County (Jantz et al., 1982) was consulted. After an initial field survey, it was determined that the relationship of soil units and bedrock geology was too poor and unpredictable to use the soil maps exclusively for mapping (table 1). The correlation between soil units and unconsolidated deposits, specifically gravel and loess, was sufficient, however, for preliminary mapping in the laboratory. Loess boundaries, for example, could be generally defined by those of the Harney and Holdredge soil map units, with subsequent field refinement.

Initial mapping focused on alluvial deposits because aerial photography permitted easy recognition and delineation of floodplains and terraces. Rocks of the Cretaceous System, the most time-consuming to map, were mapped next. The Fencepost Limestone, a widespread, resistant bed of chalky limestone capping the Greenhorn Limestone, was an important stratigraphic marker during the early stages of surficial bedrock mapping because it is easily detected on the aerial photography. We next determined the distribution of the Graneros Shale, which delineated the lower and upper contacts of the Greenhorn Limestone and Dakota Formation, respectively. Although boundaries of the Graneros were often hard to find due to its thinness and sparse exposure, its mapping was facilitated by the location of numerous farm ponds situated on the impermeable shale. Loess deposits were delineated in the final phase of mapping.

As mapping progressed, geologic contacts were first drawn on paper copies of the 7½-minute, 1:24,000-scale topographic quadrangles and were then transferred to planimetric Mylar base maps of the same scale. Subsequent digitizing and map production through computer-aided cartography using the GIMMAP software was conducted at the Kansas Geological Survey.

TABLE 1. Degree of correspondence between geologic map units and soil series associations.

Geologic unit	Soil series	Correspondence of oil distribution to geology ^a
Loess	Crete	good
	Harney	good
Ogallala Formation	Dorrance	good
Niobrara Chalk	Nibson ^b	fair
Carlile Shale	Bogue ^c	good
	Corinth ^d	fair
Greenhorn Limestone	Armo	fair
	Edalgo	fair
	Nibson ^e	fair
	Wakeen ^f	fair
Graneros Shale	Nibson ^e	poor
	Lancaster ^g	poor
Dakota Formation	Hedville	good
	Lancaster ^g	fair
High terrace (gravel)	Dorrance	good
	Wells	fair
Low terrace and floodplain	Detroit	fair
	Humbarger	fair
	Inavale	fair
	McCook	fair
	Munjor	fair
	Roxbury	fair

a. Scaling: Excellent: $\geq 80\%$ correspondence between geologic and soil-series map units. Good: ≥ 50 to $< 80\%$ correspondence. Fair: at least 50% correspondence. Poor: geologic unit is unmappable from the soil series map unit.

b. Soil series do not distinguish between the Smoky Hill and Fort Hays Members.

c. Soil series distinguishes the Blue Hill Shale Member from the Fairport Chalk Member.

d. Soil series distinguishes the Fairport Chalk Member from the Blue Hill Shale Member, but does not distinguish the Fairport Chalk Member from the Greenhorn Limestone.

e. Soil series does not distinguish the Greenhorn Limestone from the Carlile Shale nor the Graneros Shale.

f. Soil series does not distinguish the Greenhorn Limestone from the Carlile Shale.

g. Soil series does not distinguish the Dakota Formation from the Graneros Shale.

Stratigraphy of outcrops

Rock units cropping out in Russell County range in age from Lower Cretaceous to Holocene and include the Dakota Formation, Graneros Shale, Greenhorn Limestone, Carlile Shale, Niobrara Chalk, Ogallala Formation, and a variety of unconsolidated Quaternary deposits (loess and alluvium). Rock- and time-stratigraphic units germane to this study are represented in fig. 3.


Cretaceous System—Lower Cretaceous Series

Dakota Formation


Meek and Hayden (1862) first used the name Dakota Group to describe exposures of sandstone, clays of various

colors, and lignite beds near Dakota City, Dakota County, Nebraska. Plummer and Romary (1942) subsequently used the term Dakota Formation to include all deposits of dominantly continental and littoral origin lying between the transgressive deposits of the Kiowa Shale and the Graneros Shale.

The Dakota Formation crops out through much of western Kansas, where the formation has a thickness ranging from 200 ft to 300 ft (60 m to 90 m). In Russell County, the upper 100–150 ft (30–46 m) of the Dakota Formation is exposed (Hattin and Siemers, 1978). Six major facies have been delineated in this part of the Dakota (Hattin, 1965b; Siemers, 1971a, 1976; Hattin and Siemers, 1978) and represent a transition to shallow-marine deposits in an environmentally diverse, deltaic, and



CENOZOIC ERATHEM



UPPER CRETACEOUS CRETACEOUS SYSTEM

MESOZOIC ERATHEM

marginal marine setting at the onset of the Greenhorn transgression. In general, these facies consist of yellowish to reddish and greenish and brownish mottled mudstones, lignites, and ironstones interbedded and overlain by cross-bedded, elongate-trough-shaped sandstones (e.g., Rocktown channel sandstone) and flat-bedded, elongate-tabular-shaped sandstones. A variety of trace fossils such as *Planolites* and *Skolithos* are common in the Dakota Formation as well as such diverse, marginal marine macroinvertebrates as *Geltena subcompressa* and *Exogyra spooneri* (Hattin, 1967; Siemers, 1971b).

Although the Dakota Formation underlies all of Russell County, it is extensively exposed only in the eastern one-

half of the county. A variety of excellent exposures of the Dakota exist, including two (figs. 4 and 5) generally described by Hattin and Siemers (1978). On the geologic map of the county (Johnson and Arbogast, 1996), all major facies of the Dakota are included within the map unit **Kd**.

Cretaceous System—Upper Cretaceous Series

Graneros Shale

The Graneros Shale was named by Gilbert (1896) for exposures along Graneros Creek near Walsenburg, Colorado. The earliest definitive description of the Graneros Shale in Kansas, then called "the Bituminous Shale Horizon," was by Logan (1897) when he recorded general attributes such as thickness, color, and structure. The next year, Logan (1898) correlated this unit with the Graneros Shale of Colorado. Rubey and Bass (1925) extensively described the Graneros Shale in Russell County, but the most detailed study of the Graneros in Kansas was conducted by Hattin (1965a) who analyzed the stratigraphy of the unit throughout the Smoky Hills. The Graneros Shale is composed predominantly of fissile, noncalcareous, medium- to dark-gray shale that weathers to shades of gray and yellowish brown; it is approximately 30 ft (10 m) thick in Russell County (Rubey and Bass, 1925). Clay-ironstone and calcareous septarian concretions occur locally. At many localities, interbeds of calcareous and noncalcareous siltstone and sandstone lie in the middle part of the shale. In addition, several layers of bentonite and beds of skeletal limestone are found in the middle and upper parts of the Graneros. A variety of invertebrate fossil material, including *Inoceramus rutherfordi* and *Ostrea beloiti*, are common in the upper part of the formation.

The Graneros Shale crops out as a narrow formation between the overlying Greenhorn Limestone and underlying Dakota Formation in Russell County (Johnson and Arbogast, 1996). According to Rubey and Bass (1925), irregular variations in the thickness of the shale in Russell County may exceed 20 ft (4.5 m) due to intertonguing with the Dakota. The outcrop belt may be as much as 1 mile (1.6 km) wide on the northern side of stream valleys due to the regional northwest dip of the bedrock, but it is usually less than a quarter of a mile (0.4 km) throughout the county. Topographically, the Graneros usually forms low-angle slopes, except in steep-sided valleys where the shale is protected by resistant rocks at both contacts. Complete exposures of the Graneros Shale are limited in Russell County because dense sod or talus usually covers part of the unit. Hattin (1965a) described an exposure, however, in the NW sec. 35, T. 12 S., R. 14 W., along U.S. Highway 281, north of Russell (fig. 6). Another exposure of the Graneros, in the NE sec. 33, T. 12 S., R. 14 W., was created by recent slumping; in this exposure the contact with the overlying Greenhorn Limestone is nicely ex-




FIGURE 3. Classification of rocks in Russell County (after Baars, 1994).




FIGURE 4. Exposure of the upper part of the Dakota Formation located in a roadcut approximately 0.5 mile (0.6 km) from the Russell County line in the NW SE sec. 19, T. 13 S., R. 10 W., Lincoln County. Active fill of the Rocktown channel sandstone is represented in units 1a and 1b, with large trough features in unit 1a—1–3 ft (0.3–0.9 m) high and 5–15 ft (1.5–4.6 m) across—and relatively small crossbed sets in unit 1b, 0.5–1.5 ft (0.15–0.46 m). The unit 1 sandstone bed is truncated by carboniferous, silty claystone and shale of unit 2, a 25-ft (7.6-m)-thick deposit of abandoned-channel fill. Unit 3, which is 16 ft (4.9 m) thick, is a root-mottled, carbonaceous sandy siltstone of unknown origin. Unit 4 consists of flat-bedded sandstone that is 31 ft (9.5 m) thick. Reproduced from Hattin and Siemers (1978).

pressed (fig. 7). On the geologic map of the county (Johnson and Arbogast, 1996), the Graneros Shale is designated **Kgr**.

Greenhorn Limestone

In central Kansas, the Greenhorn Limestone outcrop extends northeastward from Ford County, through Russell County, into the northwestern corner of Washington County on the Nebraska border. The Greenhorn is unconformable on the Graneros Shale at most localities (Hattin, 1975). The first references and descriptions of rocks, later included within the Greenhorn Limestone, were made by Hayden (1872), Mudge (1876), and St. John (1883). In 1899, Logan described in detail the rocks of the Limestone Group, later to become the Greenhorn. The term “Greenhorn” was first used for Kansas rocks by Darton (1904) to describe the flaggy and chalky limestones of central Kansas. The earliest comprehensive studies of the Greenhorn were conducted by Rubey and Bass (1925) in Russell County and by Bass (1926) in western Kansas. In these reports, the fundamentals regarding thickness, lithology, important fossils (e.g., *Inoceramus prefragilis*, *Ostrea beloiti*), and stratigraphic correlations were established, as well as the subdivision of the Greenhorn into the Pfeifer, Jetmore, Hartland, and Lincoln members. The most intensive and extensive study of the stratigraphy and depositional environment of the Greenhorn, including the most detailed analysis of each member, was conducted by Hattin (1975).

The Greenhorn Limestone is approximately 100 ft (30 m) thick in Russell County. The Lincoln Limestone Member at the base of the Greenhorn is about 20 ft (6 m) thick (Rubey and Bass, 1925). Interbedded chalky shales and chalky limestones, both of which are light gray and weather to yellowish gray or yellowish tan, compose the bulk of the Lincoln. Also contained within the shales of the Lincoln are thin beds of bentonite and isolated lenses of skeletal limestone. Conformably above the Lincoln is the Hartland Member, which is about 35 ft (11 m) thick. Although this unit is typically referred to as the Hartland Shale, Hattin (personal communication) favors discontinuing the use of the term “Shale” because it is not a true shale. The contact between the Lincoln Limestone and the Hartland is at the top of a concentration of thin beds or lenses of skeletal limestone (Hattin, 1975). The Hartland is largely a gray, chalky shale that contains numerous bentonite seams and a few thin beds of chalky limestone. Conformably overlying the Hartland is the Jetmore Chalk Member, whose contact with the Hartland is placed at the base of the lowest of numerous, closely spaced, conspicuous hard beds of chalky limestone (Hattin, 1975). The Jetmore Member, which is approximately 20 ft (6 m) thick in Russell County, consists of chalky shale and 12 to 15 beds of chalky limestone; the uppermost bed of the member is the *Inoceramus* or Shellrock limestone. At the top of the Greenhorn is the Pfeifer Member, which has an average thickness of 20 ft (6 m) in Russell County (Rubey and Bass, 1925) and conformably overlies the Jetmore Member. Typically this unit is referred to as the Pfeifer




FIGURE 5. Exposure of the upper part of the Dakota Formation, illustrating the complex relationships of lithologic units, located in a roadcut approximately 0.5 mi (0.6 km) from the Russell County line in the NW SE sec. 19, T. 13 S., R. 10 W., Lincoln County. In general, unit 1 represents active-channel fill, while units 2–6 include complex floodplain and partially abandoned channel-fill deposits. Units 7–12 are likely deltaic deposits from a mixed fluvial and marginal-marine environment during the onset of the Graneros transgression. The discordant surface labeled 4 may represent faulting of unit 5 against units 1 and 2 before aggradation of unit 6. Reproduced from Hattin and Siemers (1978).

Shale, but Hattin (personal communication) suggests that the term “Shale” be dropped because the Pfeifer is not a true shale. In general, the Pfeifer consists of interbedded chalky shale and chalky limestone that accumulated during a maximum transgression. Three marker beds are contained within the Pfeifer. The lowermost is an unnamed bed of chalky limestone that directly overlies a thin seam of bentonite. An intermediate marker bed is a bentonite and granular-calcite unit known as the “sugar sand.” At the top of the Pfeifer is the well-known Fencepost Limestone, which is bluish gray and weathers to light tan.

The Greenhorn Limestone is widespread in Russell County with the outcrop belt ranging from less than one-

fourth mile (0.35 km) along bluffs to several miles (>1.5 km) in less dissected, interstream areas (Rubey and Bass, 1925). Intensive dissection of the Greenhorn, especially in the central part of the county, by tributaries in the southern uplands of major stream basins has resulted in steep canyon walls where members of the Greenhorn are clearly visible. An exposure described by Hattin (1975) on the east side of the Bunker Hill–Luray Road halfway along the western edge of sec. 18, T. 13 S., R. 12 W. (fig. 8) illustrates the nature of the unit. An exceptional exposure of the Greenhorn, one that shows all four members (fig. 9), is located in a tight meander bend of the Saline River in the NE sec. 33, T. 12 S., R. 14 W. Abandoned and active



FIGURE 6. Exposure of the Graneros Shale, located in the NW sec. 35, T. 12 S., R. 14 W., on U.S. Highway 281 north of Russell. Described by Hattin (1965a), the Graneros at this locality lies conformably on the Dakota Formation. A. Arbogast is kneeling on the contact.



FIGURE 7. The Graneros Shale–Greenhorn Limestone contact (rock hammer) in an exposure created by recent slumping (NE sec. 33, T. 12 S., R. 14 W.).



FIGURE 8. The Hartland Shale and Jetmore Chalk Members of the Greenhorn Limestone exposed in a roadcut, described by Hattin (1965b), on the east side of the Bunker Hill–Luray Road, halfway along the western edge of sec. 18, T. 13 S., R. 12 W. The arrow points to the approximate location of the Hartland-Jetmore contact.

Fencepost quarries, such as the one illustrated in fig. 10, are scattered on upland margins throughout the county. Because the rocks of each member of the Greenhorn are genetically similar (i.e., marine sediments that accumulated during a transgressive phase), the formation has not been subdivided on the geologic map of the county (Johnson and Arbogast, 1996); they are included within the map unit **Kgh**.

Carlile Shale

The Carlile Shale was named by Gilbert (1896) from exposures along the Arkansas River at Carlile Station southwest of Pueblo, Colorado. Subsequent modification

and subdivision of the Carlile Shale in Kansas was done by Logan (1899) and Bass (1926). Hattin (1962) conducted the most intensive study of the Carlile Shale, including a detailed analysis of each member. In Kansas, the Carlile crops out generally in two areas (Hattin, 1962): an extensive outcrop extending from Finney County northeastward to Washington County and a smaller area in northwestern Hamilton County. The Carlile Shale is approximately 300 ft (90 m) thick in Russell County, with the thickest outcrops in the northwestern corner of the county. It consists of three members. At the base of the Carlile is the Fairport Chalk Member, which is approximately 85 ft (55 m) thick. The middle member is the Blue Hill Shale Member, an approximately 195-ft (125-m)-thick



FIGURE 9. Extensive exposure of the Greenhorn Limestone along a meander bend of the Saline River in the NE sec. 33, T. 12 S., R. 14 W. In Russell County, the Greenhorn Limestone unconformably overlies the Graneros Shale, which is the dark unit adjacent to A. Arbogast. At this locality, all four members of the Greenhorn (Lincoln, Hartland, Jetmore, Pfeifer) are exposed.



FIGURE 10. A small fence-post quarry (abandoned) in the SE sec. 29, T. 11 S., R. 15 W. The Fencepost Limestone (arrow) is the uppermost bed of the Greenhorn Limestone. Early settlers of Russell County discovered that, once exposed, the Fencepost Limestone was very soft and could be easily worked. Upon subsequent weathering, the chalk hardened. Conformably overlying the Fencepost Limestone at this locality is the lower part of the Fairport Chalk Member of the Carlile Shale.




FIGURE 11. Map showing the location of sinkholes relative to Interstate 70 in secs. 2 and 3, T. 14 S., R. 15 W. in western Russell County (from Knapp et al., 1989). The tick marks along the axes formed by I-70 and the north-south section boundary represent 10-m intervals, which are numbered every 20 m.

unit of dark-gray, clayey, fissile shale. At the top of the Carlile is the Codell Sandstone Member, which has a maximum thickness of no more than 25 ft (7.6 m) and thins to less than 1 ft (0.3 m) (Rubey and Bass, 1925). On the geologic map of the county (Johnson and Arbogast, 1996), the Fairport and Blue Hill members were differentiated and mapped as **Kc** and **Kb**, respectively. Because only one exposure of the Codell Sandstone was found, the Codell is undifferentiated from the Blue Hill Shale Member and is included within the map unit **Kb**.

Fairport Chalk Member. Formerly the Fairport Chalky Shale, the Fairport Chalk Member is a complex of marine chalks and marls that lie conformably on the Greenhorn Limestone. Outcrops of the Carlile Shale are widespread in Russell County, with isolated remnants of the lower part on hilltops in the eastern third of the county and extensive outcrops in the western two-thirds of the project area. In the northwestern corner of Russell County, the entire thickness of the Fairport is exposed (Rubey and Bass, 1925).

Along Interstate 70, near the western county line, three sinkholes (Witt, Crawford, and Roubach) have recently formed in the surface of the Carlile Shale (fig. 11). Seismic investigations of the sinkholes indicates they have developed during the last 25 years due to significant dissolution

of the Hutchinson Salt member of the Permian Wellington Formation 400–500 m (1,300–1,600 ft) below the surface (fig. 12). Apparently, dissolution of the Hutchinson Salt Member is associated with saltwater disposal wells (fig. 11) within the Gorham oil field (Knapp and Steeples, 1981; Steeples et al., 1986; Knapp et al., 1989).

As it occurs in Russell County, the Fairport Member consists of bluish-gray to gray chalky shale, chalky limestone, and calcareous shale that weather yellowish gray and grayish orange; it is considerably more resistant than the overlying Blue Hill Shale Member. At the base of the Fairport Member, chalky limestone is more abundant and is identical to the upper beds of the Greenhorn Limestone, reflecting deposition during a maximum transgression (Hattin, 1962). Thin bentonite beds occur throughout the member. Important fossils include *Inoceramus* and *Pseudoperina*, fish scales and sharks' teeth. A section of the lower Fairport Member, measured by Hattin (1962) in the NE sec. 5, T. 13 S., R. 15 W., is shown in fig. 13.

Blue Hill Shale Member. Although a type section has never been designated, the name *Blue Hill Shale* was apparently derived from the Blue Hills of western Mitchell County, Kansas, by Logan (1897). In Russell County, the




FIGURE 12. Geologic profile of the Crawford sink in the SW sec. 2, T. 14 S., R. 15 W. (from Knapp et al., 1989).

Blue Hill Shale Member consists of a dark-gray clayey, blocky to fissile shale, that contains septarian concretions of calcium carbonate and clay-ironstone. The Blue Hill Member lies between the underlying chalky shale of the Fairport Member and the overlying Codell Member of the Carlile Shale. The Blue Hill Shale Member consists primarily of quartz-dominant terrigenous detritus eroded from surrounding Paleozoic and Precambrian terranes during a regressive phase (Hattin, 1962). Strikingly uniform throughout, the primary differences within the Blue Hill are the number, size, and composition of the calcium concretions and an increase in sand content toward the top of the member (Rubey and Bass, 1925). The Blue Hill Member crops out on the higher ridges in the northwestern corner of Russell County and is usually expressed topographically as broad, gentle slopes, but may be exposed locally on steep bluffs where the overlying, more resistant Fort Hays Limestone serves as a protective caprock (fig. 14). Recent slumping in the SE sec. 5, T. 11 S., R. 15 W., has exposed portions of the Blue Hill Member (fig. 15), but it may also be observed on the surface of unimproved roads (fig. 16).

Codell Sandstone Member. Bass (1926) named and described the Codell sandstone bed from a sandstone overlying the Blue Hill Shale Member in the vicinity of Codell, along the Saline River valley in Ellis County. The

Codell was elevated to member status by Dane and Pierce (1933), but was relegated to informal rank of zone within the Blue Hill Shale member by Moore et al. (1951). Informal rank persisted until the Codell was re-elevated to member status by Merriam (1957) and Jewett (1959).

As defined, the Codell Sandstone Member is a fine-grained, silty sandstone that is locally shaly. The lower contact with the Blue Hill Shale Member is conformable and gradational, whereas the upper contact with the Fort Hays Limestone is unconformable and abrupt. Fossils are rarely found in the Codell Sandstone. In Russell County, the total thickness of the Codell Sandstone is unknown. The only known exposure of the Codell recognized in this study, a result of recent slumping, is located in the SE sec. 5, T. 11 S., R. 15 W. (fig. 17).

Niobrara Chalk

Meek and Hayden (1862) named the Niobrara Chalk from exposures of chalky limestone and calcareous marl in the Niobrara River valley of northeastern Nebraska. The formation was later divided into the Fort Hays Limestone and Smoky Hill Chalk Members by Logan (1897). Subsequently, the stratigraphy and paleontology of the Niobrara in western and central Kansas were described and analyzed by a number of researchers (e.g., Bass, 1926; Elias, 1931; Moss, 1932; Russell, 1929; Cobban and Reeside, 1952; and Hattin, 1982). The Niobrara crops out only in the extreme northwestern corner of Russell County where a maximum of 100 ft (30 m) remains. The lower 45–50 ft (14–15 m) consists of the Fort Hays Limestone, and the remainder is the Smoky Hill Chalk Member (Rubey and Bass, 1925). On the geologic map of the county (Johnson and Arbogast, 1996), the two members are differentiated and mapped as **Knf** and **Kns**, respectively.

Fort Hays Limestone Member. Unconformably overlying the Blue Hill Shale is the Fort Hays Limestone Member. The Fort Hays Limestone is characterized by thick to very thick beds of resistant chalky limestone that is light gray to medium gray or light olive gray and that weathers yellowish gray, grayish or pale grayish orange, or almost white. Thin beds of bentonite and shale separate many of the limestone beds, which consist largely of homogeneous, micrograined carbonate rock that have been thoroughly bioturbated. Increased sand content at the base of the Fort Hays apparently reflects reworking of Codell sands at the onset of the Niobrara transgression (Hattin and Siemers, 1978). Macrofossils (e.g., *Inoceramus deformis*, *Inoceramus browni*) and trace fossils (e.g., *Teichichnus*, *Planolites*) are found in the member. In Russell County, the Fort Hays is well exposed at only two localities, a slump in the NW sec. 5, T. 11 S., R. 15 W. (fig. 17) and a quarry in the NW sec. 6, T. 11 S., R. 15 W. (fig. 18). The contact with the overlying Smoky Hill Chalk Member was



FIGURE 13. Roadcut in the lower part of the Fairport Chalk Member of the Carlile Shale, located in the NE sec. 5, T. 13 S., R. 15 W., described by Hattin (1962). The spade (arrow) rests on the Fencepost Limestone of the uppermost Greenhorn Limestone. The Fairport Chalk Member is similar to the Greenhorn Limestone in that alternating beds of chalk are separated by beds of marl and chalky shale.

difficult to locate precisely due to the lack of a suitable outcrops.

Smoky Hill Chalk Member. The Smoky Hill Chalk Member has been intensively studied by Hattin (1982). In Kansas, the Smoky Hill outcrop extends northeasterly 190 mi (306 km) from north-central Finney County to north-eastern Jewell County on the Nebraska border. The type area is in western Kansas and consists of a composite section from 12 localities that is about 596 ft (182 m) thick. The Smoky Hill Member lies conformably over the Fort Hays Limestone Member and is dominated by fecal-pellet-speckled shaly chalk, which is light olive gray to olive gray. More than 100 bentonite seams, many of which have weathered to iron oxide and gypsum, have been recognized in the Smoky Hill by Hattin; they are more abundant in the lower part of the member. Vertebrate and invertebrate fossil material is common in the chalk, with the most obvious fossils being *Inoceramus grandis* and *Pseudoperna congesta*. In the very northwestern corner of the county, the Smoky Hill consists of thin remnants that cap the higher hills held up by the Fort Hays Limestone. The best exposure occurs in a quarry located in the NW sec. 6, T. 11 S., R. 15 W. (fig. 19), which also contains the Fort Hays Limestone.

Tertiary System—Miocene Series

Ogallala Formation

The Ogallala Formation is a heterogeneous complex of massive to crossbedded alluvium deposited during the Miocene and early Pliocene by streams originating in the Rocky Mountain region. The Ogallala covers most of the High Plains and is the primary aquifer in the region. A major discontinuity separates the Ogallala from both the underlying Cretaceous deposits and overlying unconsolidated sediments of the Quaternary System in Russell County.

Darton (1899) first proposed the name Ogallala based on exposures near Ogallala Station in southwestern Nebraska. Subsequently, in a report on the geology of Hamilton and Kearny counties in southwestern Kansas, Darton (1920) identified a general type locality in the vicinity of Ogallala Station. Detailed studies of the Ogallala in Kansas, which included designation of a precise type locality near Ogallala Station in Nebraska, were conducted by Elias (1931, 1932, 1935, 1942). The Nebraska Geological Survey later classified the Ogallala as a group and subdivided it into four formations.



FIGURE 14. Typical landscape developed on the Blue Hill Shale Member of the Carlile Shale located in the SE sec. 5, T. 11 S., R. 15 W. The Blue Hill Shale Member crops out in the northwestern corner of Russell County. Topographically, the nonresistant shale is expressed as broad, gentle slopes, such as those in the foreground, but forms steeper slopes where capped by the resistant Fort Hays Limestone Member of the Niobrara Chalk, such as those on the horizon of the photograph.



FIGURE 15. Slumping of the Blue Hill Shale Member of the Carlile Shale in the NE sec. 5, T. 11 S., R. 15 W. Persistent, heavy rains in summer 1993, resulted in mass wasting of the Blue Hill at many localities where the shale is capped by the Fort Hays Limestone Member of the Niobrara Chalk on steeper slopes.



FIGURE 16. Outcrop of the Blue Hill Shale Member of the Carlile Shale in the roadcut and road surface along the SW boundary, secs. 5 and 8, T. 11 S., R. 15 W. The overlying Fort Hays Limestone Member of the Niobrara Chalk can be seen in the road at the top of the hill (arrow).

The age of the Ogallala has been somewhat controversial. Darton (1899) argued that the Ogallala was late Tertiary. Based on work in Wallace County, Elias (1931) believed that mammalian fauna in the Ogallala indicated a late Miocene to early Pliocene age. Boellstorff (1976) obtained a late Miocene age of 7.6 ± 0.7 Ma from volcanic glass shards (fission-track dating) collected at the Ogallala type section. In Texas, where both fluvial and eolian facies in the Ogallala have been observed, deposition was recognized at about 8.5 Ma (Gustavson and Winkler, 1988). According to Voorhies (1990), the onset of Ogallala time in Nebraska appears to correlate with the medial Barstovian land mammal age at about 14 Ma. Voorhies also obtained a range of fission-track ages in Nebraska from 13.6 ± 1.3 Ma to 5.0 ± 0.2 Ma, which suggest that a medial age of 9.5 Ma (14–4.5 Ma) is indicated for the Ogallala in the region.

Three members. The Kansas Geological Survey recognizes three members within the Ogallala Formation (Bayne and O'Connor, 1968), which have been differentiated and correlated by floral and faunal remains, volcanic ash petrology, and gross lithology. The lowermost member is the Valentine, which is confined to Cretaceous bedrock valleys, and consists of medium- to fine-grained sands and gravels that are greenish gray to pink and pale tan. Layers of volcanic ash and bentonite are common as are silica-

cemented sand and gravel, diatomaceous marl, and a diagnostic fossil grass seed (*Stipidium commune*).

Overlying the Valentine is the Ash Hollow Member, which consists of lithologically heterogeneous, generally coarse, gray to pink sand and gravel that crop out in bedrock valleys and on valley side slopes and divides. The Ash Hollow contains rich floral zones (e.g., *Krynitzkia coroniformis* seed zone), molluscan assemblages, and abundant volcanic ash beds (particularly in the lower half) that are cemented by silica locally to form erosion-resistant lentils of sandstone and are weakly cemented with carbonate at other localities to form mortar beds.

The uppermost member of the Ogallala is the Kimball, which consists largely of highly calcareous and predominantly gray fine- to medium-grained sand, silt, and clay that accumulated after bedrock valleys were filled by the Valentine and Ash Hollow Members. As a result, the Kimball consists of a thin, unconfined mantle of sediments on the landscape. Capping the Kimball is a dense pink to white pisolitic nodular caliche (calcrete) that is 3 ft (0.9 m) thick. The member has an observed thickness ranging from 2.5–44 ft (0.76–13 m) and is easily differentiated by distinctive plant fossils and molluscan assemblages that occur in the basal part.

The Ogallala Formation is neither a major nor pervasive geologic unit in Russell County. In fact, the correlation of



FIGURE 17. Outcrop of the Codell Sandstone in SE sec. 5, T. 11 S., R. 15 W., at an exposure created by recent slumping of underlying Blue Hill Shale Member. At this locality, approximately 2 m (6.5 ft) of the Codell is exposed. The lower contact of the Fort Hays Limestone Member of the Niobrara Chalk is also visible in the exposure.

the Ogallala in Russell County by Rubey and Bass (1925) and Frye and Brazil (1943) is extremely tentative, based on lithologic similarities and recognition of the formation to the north, south, and west of Russell County. Uncertainty exists because cemented sand and gravel in Russell County does not correlate in elevation and gradient with known Ogallala outcrops immediately to the west in Ellis County (Frye and Brazil, 1943). According to Frye and Brazil (1943), however, cemented sand and gravel deposits in the uplands of Russell County are undoubtedly older than unconsolidated Pleistocene sand and gravel deposits found at lower elevations, suggesting to them an uppermost Pliocene or lowermost Pleistocene age.

Although confusion remains concerning the age of cemented upland sand and gravel deposits in Russell County, their lithologic similarity with known Tertiary deposits to the west favors tentative correlation with the Ogallala Formation, at least until more is known. It is impossible, however, to determine positively which member of the Ogallala occurs in Russell County. As a result, the deposits are not differentiated on the geologic map of the county (Johnson and Arbogast, 1996) and are collectively designated **To**. Two areas of outcrop for the Ogallala occur in Russell County. The largest area is the very northwestern corner of the county in the vicinity of the Niobrara Chalk outcrops, where the Ogallala consists

of thin, isolated remnants high on valley walls. The best exposure of the Ogallala in this area is in a roadcut in the SE sec. 9, T. 11 S., R. 15 W. (fig. 20). An isolated exposure is in a gravel quarry in the NE sec. 13, T. 14 S., R. 14 W., southeast of Russell (fig. 21). At both localities, the Ogallala consists of massive to crossbedded sand and gravel, and grit and clay that are locally cemented by calcium carbonate. Clay balls were also recognized in the deposits, particularly in the northwestern outcrop area.

Quaternary System—Pleistocene Series

Classically, the Pleistocene has been divided on the basis of four glacial advances or stages: the Nebraskan, Kansan, Illinoian, and the Wisconsinan. Due to the complexity of the stratigraphic record and faulty correlations by early researchers, however, the traditional Pleistocene chronology has largely been abandoned in recent years. Many of the early stratigraphic correlations, for example, were made on the basis of a single volcanic ash marker bed, the Pearlette (Frye and Leonard, 1952; Schultz and Martin, 1970). In fact, three separate Pearlette ashes and three other ashes have been differentiated (table 2) in the central Great Plains on the basis of petrographic and chemical characteristics (Izett et al., 1970; Izett, 1981; Izett and Wilcox, 1982) and fission-track dating

TABLE 2. Fission-track dated volcanic ashes of major stratigraphic significance in the Central Plains (Nebraska, Kansas, and Oklahoma).

Ash	Alternate names	Source area	Representative age ($\times 10^6$ yr)
Pearlette O	Pearlette restricted, Lava Creek B, Cudahy, Hartford	Lava Creek Tuff, Member B, Yellowstone	0.610
Bishop	Mount Clare	Bishop Tuff	0.738
Tsankawa	—	Toledo & Valles caldera, NM	1.12
Pearlette S	Coleridge, Mesa Falls	Mesa Falls Tuff, Yellowstone	1.27
Guaje	—	Toledo & Valles caldera, NM	1.45
Pearlette B	Huckleberry, Ridge, Borchers	Huckleberry Ridge Tuff, Yellowstone	2.01

Sources: Boellstorff (1976), Izett (1981, 1982), Izett and Wilcox (1982), Richmond and Fullerton (1986), and Ward (1991).

(Boellstorff, 1973, 1974, 1976; Naeser et al., 1973; Carter et al., 1990; Ward et al., 1993).

An informal scheme of geologic divisions and associated chronology for the Quaternary of the United States has been assembled using various stratigraphic markers (e.g., Pearlette O volcanic ash, glacial tills, Matuyama-Bruhnes magnetic polarity reversal) and the marine ^{18}O record (fig. 22). The Pliocene-Pleistocene boundary was set at 1.65 Ma, in accordance with the corresponding radiometric-age estimate from the Virca section in Italy (Aguirre and Pasini, 1985). Richmond and Fullerton (1986) subdivided the Pleistocene into early, middle, and late, with the middle being further divided into early middle, middle middle, and late middle Pleistocene. Pleistocene time prior to the Illinoian is referred to as pre-Illinoian on the basis of the ^{18}O record, which indicates that ten glaciations (seven Pleistocene, three Pliocene) occurred prior to 302 ka rather than two, as proposed in classical theory. A more detailed discussion of the Pleistocene chronology in Kansas is found in Bayne and O'Connor (1968) and Johnson (1993).

The late Quaternary stratigraphic succession for Russell County is represented in fig. 23. Pre-Illinoian deposits are tentatively recognized in the county but are spatially isolated and limited to the upland margins of major valleys. Illinoian deposits occur more extensively but are also isolated in upland and valley contexts. Wisconsinan sediments are widespread, with appreciable surface expression on both the uplands and along stream valleys. Holocene deposits are very common and are recognized in both upland and alluvial settings.

Pre-Illinoian deposits

Isolated pre-Illinoian deposits of unconsolidated gravel, sand, and silt, with lentils of volcanic ash, are found throughout northwestern Kansas as terrace-fill remnants in major valleys. Originally, these nonglacial sediments were included within the Meade Formation and were thought to be of classical Kansan age. The Meade Formation was later subdivided into the Grand Island and Sappa Members (Frye and Leonard, 1952) based on stratigraphic nomenclature derived in Nebraska by Lugn (1934, 1935) and Condra et al. (1950). Subsequently, the name of Meade Formation was abandoned in favor of elevating the Grand Island and Sappa units to the formational level (Bayne and O'Connor, 1968).

Despite a lack of absolute age control (e.g., fission-track, thermoluminescence) validating the continued use of "Grand Island" and "Sappa" as formation names, their use persists. The Grand Island Formation, as it is defined, consists of gravel, sand, and minor amounts of silt approximately 20 ft (6.1 m) thick that are present in northwestern Kansas beneath a highly dissected terrace in major stream valleys. Conformably overlying the Grand Island is the Sappa Formation, which is composed of alluvial silt and small amounts of sand with lentils of Pearlette volcanic ash (Frye and Leonard, 1952).

Isolated deposits of unconsolidated sand, silt, and gravel, potentially correlating to the Grand Island and Sappa Formations, as they are presently defined, are located in Russell County along the Smoky Hill River, Saline River, and their major tributaries (e.g., Wolf Creek).



FIGURE 18. Gravel quarry in the NW sec. 6, T. 11 S., R. 15 W., exposing the upper part of the Fort Hays Limestone Member of the Niobrara Chalk. The Niobrara Chalk is confined to the northwestern four sections of Russell County.

These deposits were originally described by Logan (1897), who referred to them as the "Salt Creek gravel beds." Rubey and Bass (1925) recognized two gravel beds, an upper and a lower one, in the Saline River valley. The lower gravel bed, which corresponds to Logan's (1897) Salt Creek gravel beds, occurs consistently about 50 ft (15.3 m) above the modern floodplain; it is approximately 2 ft (0.6 m) thick. The upper gravel bed is found between 40–70 ft (12.2–21.4 m) above the modern floodplain and is about 10 ft (3 m) thick. Gravel beds similar to the upper gravel bed in the Saline valley were also recognized by Rubey and Bass (1925) along the Smoky Hill and its major tributaries. Frye and Brazil (1943), on the basis of fossil assemblage and terrace elevation, suggested that the gravel beds in Russell County correlate to the McPherson Formation farther downstream.

Although no absolute time control exists, upland gravel deposits in major stream valleys of Russell County are thought to be of pre-Illinoian age and may correlate to the Grand Island and Sappa Formations, if continued use of this nomenclature is indeed appropriate. On the geologic map of the county (Johnson and Arbogast, 1996), the Grand Island and Sappa Formations are undifferentiated and included within the map unit **Qg**. A typical example of one such gravel bed, exposed in the SE sec. 30, T. 14 S., R. 11 W. within the Smoky Hill River valley, is depicted in fig. 24.

Illinoian deposits

All Pleistocene deposits of Illinoian and younger age were originally included within the Sanborn Formation. As a result, the formation encompassed sediments from two glacial stages and substages as well as two prominent discontinuities, the Sangamon and Brady soils. The name "Sanborn" was derived from unconsolidated deposits in northwestern Cheyenne County, Kansas, by Elias (1931). Frye and Fent (1947) subsequently designated a Sanborn type section with subdivision into three members: the Loveland silt (Illinoian), the Peoria silt (Wisconsinan), and the Bignell silt (Holocene). A fourth member, the Crete sand and gravel, which underlies the Loveland silt, was recognized by Frye and Leonard (1949). Although the Sanborn Formation was originally thought to be a convenient term for mapping purposes, the term was abandoned in 1959 by the Kansas Geological Survey because the formation could be subdivided into discernable, mappable units and could be confused with the Sanborn Group, a thick siltstone exposed in Meade County, Kansas, which has been dated (fission-track dating) to 1.08 Ma (Carter and Ward, 1991). Instead, members of the Sanborn Formation were elevated to formational rank (Jewett, 1959), and phases were subsequently defined (Bayne and O'Connor, 1968).



FIGURE 19. Smoky Hill Chalk Member of the Niobrara Chalk exposed in a gravel quarry in the NW sec. 6, T. 11 S., R. 15 W. Only the lower 2 m (6.5 ft) of the Smoky Hill is exposed in Russell County, and the contact with the Fort Hays Limestone Member is approximate.

Crete Formation. The Crete sand and gravel is an alluvial deposit, probably reflecting several cut and fill sequences, that is presumably of Illinoian age. Originally, the Loveland unit included both eolian and alluvial sediments (Lugn, 1935), but Condra et al. (1947) introduced the name "Crete Formation" in Nebraska in order to differentiate the alluvium from the loess. Based on recent data from Nebraska, Wayne and Aber (1991) suggested that the names "Crete" and "Grand Island" may refer to the same sedimentary unit. As a result, they favored discontinuing use of the name "Crete Formation."

The Crete Formation, as it is defined in Kansas, is widely recognized in the north-central part of the state, where it

consists locally of gully fills and as terrace deposits on the north side of major river valleys (Frye and Leonard, 1952, p. 112). Typically, Crete lithology is consistent with that of the drainage basin within which the deposits occur. In Russell County, the Crete Formation is tentatively identified as underlying the Loveland Loess at the Beisel-Steinle site (Diekmeyer, 1994), a roadcut in the NW sec. 27, T. 14 S., R. 11 W. (fig. 25). In addition, it may be exposed in gravel pits along the Smoky Hill River, Saline River, and their major tributaries. Nevertheless, the Crete is not differentiated from other, presumably older, Pleistocene sand and gravel deposits in Russell County and is included within the map unit **Qg** (Johnson and Arbogast, 1996).



FIGURE 20. Exposure of cemented alluvial sand and gravel, tentatively assigned to the Ogallala Formation, in the uplands of the SE sec. 9, T. 11 S., R. 15 W. In this roadcut, the Ogallala unconformably overlies the Blue Hill Shale Member of the Carlile Shale (arrow).



FIGURE 21. Crossbedded and cemented alluvial sand and gravel, tentatively assigned to the Ogallala Formation, in a quarry in the NE sec. 13, T. 14 S., R. 14 W.

Established time divisions		Informal time divisions			Age (ka)	Oxygen marine isotope stages ²	
Period	Epoch						
Quaternary	Holocene				0	1	
	Pleistocene	Late Pleistocene	Wisconsinan	Late Wisconsinan	10 ^a /12 ^b	2	
				Middle Wisconsinan	24 ^b	3	
					59 ^b	4	
				Early Wisconsinan			a
							b
					c		
					d		
					e		
			Sangamon		50–123 ^c		
		Late Middle Pleistocene	Illinoian	Late Illinoian	130 ^b	6	
				Early Illinoian	190 ^b	7	
						8	
		Middle Middle Pleistocene	pre-Illinoian		302 ^d	9	
						10	
						11	
					12		
					13		
					14		
					15		
					16		
					17		
					18		
					19		
				20			
		⋮					
				36			
Tertiary	Pliocene				1,650 ^g	37	
						⋮	
						40	

FIGURE 22. Quaternary time scale. Modified from Richmond and Fullerton (1986, p. 6). Notes: (a) Pleistocene-Holocene boundary after Hopkins (1975). (b) Ages of the marine isotope boundaries taken from Martinson et al. (1987); values were derived using the Milankovich orbital tuning techniques. (c) The Sangamon-Wisconsinan (Interglacial-Glacial) boundary is yet uncertain and may be diachronic (Johnson et al., 1991; Curry and Follmer, 1992), 75 ka (Fulton, 1984, 1986), 100 ka (Terasmae and Dreimanis, 1976), and 122 ka (Richmond and Fullerton, 1986). The age of 116 ka has been adopted by Clark et al. (1993). (d) Ages of the marine isotope boundary interpreted by Richmond and Fullerton (1986) from graphic data presented by R. G. Johnson (1982). (e) K-Ar age of the Lava Creek tuff and Pearlette O volcanic ash bed (Izett and Wilcox, 1982). (f) Astronomical age of the Matuyama-Bruhnes magnetic polarity reversal derived by R. G. Johnson (1982) and adopted by Richmond and Fullerton (1986). (g) Age of the Pliocene-Pleistocene boundary at the Virca, Italy, section (Aguirre and Pasini, 1985).

Loveland Loess. The Loveland Loess is the most pervasive pre-Wisconsinan loess in the central United States. Shimek (1909) first identified the Loveland in exposures along the eastern bluff of the Missouri River northeast of Loveland, Iowa. Subsequently, Lugin (1935) recognized and described an eolian and alluvial component to the Loveland in Nebraska. Condra et al. (1950) later elevated the alluvial component to formational rank, naming it the Crete Formation. Recently, the Loveland Loess has been described throughout the Missouri, Mississippi, and Ohio River basins (e.g., Reed and Dreeszen, 1965; Ruhe, 1969; Willman and Frye, 1970; Ruhe and Olson, 1980) and has been recognized as far south as Arkansas and Mississippi (McCraw and Autin, 1989).

Loveland Loess consists of carbonate-enriched eolian silt that accumulated during the Illinoian Stage between

about 300 ka and 130 ka (fig. 22). The Loveland Loess is yellowish brown or reddish brown, with increased red hues toward the top of the formation due to development of the Sangamon Soil. Several paleosols of variable development, including the Sangamon Soil, have been recognized in the Loveland (Frye and Leonard, 1954; Feng, 1991; Feng et al., 1994a, 1994b). Four zones of carbonate enrichment, occurring at about 410–360, 330–290, 250–200, and 130–95 ka, were interpreted by Feng et al. (1994a, 1994b) to be soil-forming intervals associated with interglacial stages, the latter two of which would be associated with the Illinoian Stage. The thickest accumulations of Loveland Loess in Kansas are in the north-central part of the state and measure up to 49 ft (15 m). Elsewhere it is typically less than 12 m (39 ft) thick (Johnson, 1993).

In Russell County, the Loveland Loess was not differentiated from other loess formations, but it is recognized at

Time stratigraphic units		Age (ka)	Rock and pedostratigraphic units	
QUATERNARY SYSTEM	HOLOCENE SERIES	0	fluvial deposits with soils	
		5		
		10		
	PLEISTOCENE SERIES		Bignell Loess	
			Brady Soil (geosol)	
		20	Peoria Loess	fluvial deposits
		50	Gilman Canyon Formation (loess and geosol)	
		130	Sangamon Soil	
		190	Loveland Loess alluvial sand and gravel (Crete Formation?)	
		1650	loesses (?) alluvial sand and gravel (Grand Island and/or Sappa Formations?)	

FIGURE 23. Late Quaternary stratigraphic succession in Russell County. Modified from Johnson (1993).

the Beisel-Steinle site (fig. 25). The absolute age of the Loveland is unknown for Russell County, but Oviatt et al. (1988) reported thermoluminescence (TL) ages of 136 ka and 130 ka from the upper part of Loveland loess near Milford in northeastern Kansas. At the Barton County sanitary landfill, P. Maat and W. C. Johnson (unpublished data) obtained an age of 212 ka from a pre-Sangamon soil zone in Loveland Loess, whereas Feng (1991) dated the upper Loveland Loess at 69 ka.

Sangamon Soil. The Sangamon Soil is a strongly developed paleosol or complex of paleosols distributed throughout the midcontinent, including the states of Indiana (Hall, 1973; Ruhe et al., 1974; Ruhe and Olson, 1980), Illinois (Bushue et al., 1974; Follmer, 1979), Iowa (Simonson, 1941; Ruhe, 1956, 1969), Nebraska (Schultz and Stout, 1945; Thorpe et al., 1951), and Kansas (Frye and Leonard, 1952). Initially recognized by Leverett (1899) to differentiate between the Illinoian and Wisconsinan deposits, the Sangamon is characterized by intense oxidation (vivid to pale reddish-brown colors),

deep leaching, and high clay accumulation. Because of apparent time transgressiveness and the diachronous nature of the Sangamon Soil, the age of the soil is not precisely known. Estimates indicate, however, that it ranges from approximately 135 ka to 75 ka in age (e.g., Forman 1990; Forman et al., 1992; Johnson et al., 1990; Maat and Johnson, 1996).

The Sangamon Soil probably should be considered a pedocomplex, with several soils representing significant temporal and environmental variability welded together, rather than a single soil that developed under unique environmental conditions (Schultz and Tanner, 1957; Fredlund et al., 1985; Morrison, 1987). From his research in Brown County, Schaetzl (1986) observed that the Sangamon appears to be a strongly developed Ultisol or Mollisol.

Originally known in Kansas as the "soil in the Sanborn formation" (Hibbard et al., 1944) and the Loveland soil (Frye and Fent, 1947), the Sangamon has been extensively studied in the northeastern part of the state (e.g., Frye and



FIGURE 24. Alluvial sand and gravel of Pleistocene age in a quarry located on the north side of the Smoky Hill River in the SE sec. 30, T. 14 S., R. 11 W. The gravel bed unconformably overlies the Dakota Formation at this locality.

Leonard, 1949, 1952; Caspall, 1970; Schaetzl, 1986) but has been recognized elsewhere (Bayne and O'Connor, 1968). Recently, in a study of the Sangamon Soil at Barton County, Kansas, Feng et al. (1994a, 1994b) concluded from chemical and physical characteristics that the soil formed in a warm, moist climate, resulting in strong chemical weathering. In Russell County, the Sangamon Soil has been identified at the Beisel-Steinle site (fig. 25) and is assumed to exist elsewhere in the county. It is not, however, differentiated from the other late Pleistocene loesses and is included within the map unit **QI** on the geologic map of the county (Johnson and Arbogast, 1996).

Wisconsinan deposits

Since it is the most recent glacial episode, the Wisconsinan has the greatest chronostratigraphic resolution. Radiocarbon dating indicates that the Wisconsinan glaciation began 79,000 to 70,000 years ago. An early chronology, based on deposits in the state of Illinois, recognized five substages in the Wisconsinan: the Altonian (70,000–28,000 yr B.P.), Farmdalian (28,000–22,000 yr B.P.), Woodfordian (22,000–12,500 yr B.P.), Twocreekan (12,500–11,000 yr B.P.), and Valderan (11,000–5,000 yr B.P.) (Willman and Frye, 1970; Frye and Willman, 1973). In Kansas, Frye and Leonard (1965) used the Brady Soil as

a chronostratigraphic marker to divide pre-Bradyan time (Altonian, Farmdalian, Woodfordian, and Twocreekan substages) from post-Bradyan time (Valderan substage).

Subsequent research in type areas of Illinois and Wisconsin has placed the age of the Sangamon-Wisconsinan boundary at about 50 ka (McKay, 1979; Leigh, 1991, 1994; Curry and Folmer, 1992). The boundary appears, however, to be time transgressive (Johnson et al., 1991), which accounts for the lack of consistency in the limited chronologic data. In fig. 22, the Wisconsinan chronology has been subdivided so as to include Early, Middle, and Late Wisconsinan divisions, with the stage extending from about 116 ka to 10 ka (marine isotope stages 5d through 2; Clark et al., 1993). Wisconsinan rock-stratigraphic units recognized in Russell County consist of the Gilman Canyon Formation (loess) and the Peoria Loess.

Gilman Canyon Formation. The Gilman Canyon Formation, a middle Wisconsinan loess and valley fill, was first recognized in Nebraska by Reed and Dreeszen (1965). It is the apparent chronostratigraphic equivalent to the Pisgah Formation in western Iowa (Bettis, 1990, Forman et al., 1992) and the Roxana Silt (Leigh and Knox, 1993, Leigh, 1994), which has been recognized in an area extending from Minnesota and Wisconsin to Arkansas. In Kansas and Nebraska, the upper part or all of the Gilman



FIGURE 25. The Beisle-Steinle site (Diekmeyer, 1994) in the NW sec. 27, T. 14 S., R. 11 W. Poorly exposed in the roadcut are four late Quaternary stratigraphic units, from bottom to top: alluvial sand and gravel of the Crete Formation (A), Loveland Loess (L), Sangamon Soil (S), and Gilman Canyon Formation (G).

Canyon Formation is usually silty, leached of calcium carbonate, commonly bioturbated, and dark colored as a consequence of heavy enrichment of pedogenic organic carbon. The Gilman Canyon Formation was once thought to be the attenuated A horizon of the Sangamon Soil (Thorpe et al., 1951; Reed and Dreeszen, 1965), but radiocarbon time control and stratigraphic information indicate that it is a geosol or a composite geosol (Johnson, 1993) because it is a laterally traceable, mappable, pedostratigraphic unit with a consistent time-stratigraphic position (Morrison, 1965; North American Commission on Stratigraphic Nomenclature, 1983, p. 865).

In Illinois, radiocarbon ages based on organic materials from within early Wisconsinan loess range from 40,000 to 31,000 yr B.P. Based on these ages, McKay (1979) extrapolated that loess deposition began about 45,000 yr B.P. Radiocarbon and TL ages from loess lying directly above the Sangamon Soil in Iowa range from 35 ka to 30 ka (Forman, 1990; Forman et al., 1992). Radiocarbon ages from the middle Wisconsinan Roxana silt in the Upper Mississippi River valley indicate that the loess unit was deposited between 50 ka and 27 ka (Leigh, 1991, 1994; Leigh and Knox, 1993).

Radiocarbon ages from the Gilman Canyon Formation range from approximately 38,000 yr B.P. near the base to

about 20,000 yr B.P. at the top (May and Souders, 1988; Johnson et al., 1990; Johnson, 1993). The age of 38 ka is derived from a sample taken from the base of the Gilman Canyon geosol, not the base of the formation. Consequently, the age of the basal part of the formation is perhaps close to 50 ka, which would agree with that of the Roxana silt (Leigh, 1991; Leigh and Knox, 1993). In Nebraska and Kansas, the Gilman Canyon Formation has been dated at 54 localities (11 of which are in Kansas).

Limited textural data and description of the Gilman Canyon Loess at the Buzzard's Roost type section was provided by Reed and Dreeszen (1965, p. 62): "Upper 12 inches (31 cm) is medium dark gray, slightly humic, silt; middle 1 inches (1.1 m) is dark brownish-gray, humic, soil-like silt; entire thickness is noncalcareous . . . 5 feet 9 inches (1.8 m)." The bimodal distribution of the humus indicates that two periods of relative stability and low accumulation are separated by a period of increased accumulation. As a result, the Gilman Canyon is often expressed as one or more cumelic A horizons. May and Souders (1988) recognized three distinct organic zones in an expanded valley-fill section in Nebraska and two organic zones have been recognized at the Eustis Ash Pit in south-central Nebraska by Johnson et al. (1993). These observations indicate that the Gilman Canyon accumulated



FIGURE 26. Upper part of the Beisle-Steinle site (Diekmeyer, 1994) in the NW sec. 27, T. 14 S., R. 11 W. Exposed in the upper part of the site are four late Quaternary stratigraphic units—from bottom to top, Gilman Canyon Formation (G), Peoria Loess (P), Brady Soil (Br), Bignell Loess (Bi)—and modern surface soil (M).

at a rate sufficient for continuous pedogenesis (<0.08 mm/yr), but with a variable intensity.

Due to the lack of exposures, the Gilman Canyon Formation has been positively identified at only one locality in Russell County, the Beisel-Steinle site (figs. 25, 26). Two peaks exist in the organic matter curve, indicating that two distinct episodes of soil formation, separated by a brief period of increased deposition, occurred (fig. 27). Texturally, the Gilman Canyon Loess consists largely of silt, with lesser, but equal, percentages of sand and clay. Two magnetic parameters, susceptibility and frequency dependence, have been employed as surrogates of weathering and soil formation. The bimodal organic matter

distribution is corroborated by frequency dependence, an indicator of pedogenically produced fine sediments. These observations are consistent with data previously noted from the type locality (Reed and Dreeszen, 1965) and the Eustis Ash Pit of southwestern Nebraska (Johnson et al., 1993).

Four radiocarbon ages were determined from samples collected in the Gilman Canyon Loess at the Biesel-Steinle site (table 3). Based on the position of the lowest sample, soil development began shortly before about 32,800 years ago. The age of $22,590 \pm 270$ yr B.P. on the uppermost part of the formation indicates approximately when Gilman Canyon time ended. A radiocarbon age of $34,340 \pm 110$ yr

TABLE 3. Radiocarbon ages from the Beisel-Steinle site (reported in Diekmeyer, 1994).

Depth (m)	Lab number	Uncorrected age (yr B.P.)	$\delta^{13}\text{C}$ (‰)	Corrected age (yr B.P.)
1.45–1.50	Tx-8183	3,030 \pm 50	–11.5	3,250 \pm 50
2.40–2.45	Tx-8186	11,270 \pm 110	–20.1	11,350 \pm 110
3.70–3.75	Tx-8184	22,440 \pm 260	–15.9	22,590 \pm 270
4.25–4.30	Tx-8185	27,400 \pm 430	–14.4	27,580 \pm 440
about 4.2	Tx-7709	27,220 \pm 480	–14.2	27,380 \pm 490
4.95–5.00	Tx-8187	32,620 \pm 1060	–14.1	32,780 \pm 1080

B.P. (Tx-7710) was obtained from an exposure of the Gilman Canyon soil developed within alluvium of the Saline River valley in the eastern part of adjacent Ellis County (W. C. Johnson, unpublished data), indicating that an alluvial facies of the Gilman Canyon Formation may also be preserved in Pleistocene terraces of the Saline River valley in Russell County.

Carbon isotope fractionation analyses have proved to be of use in determining past vegetation and associated climatic conditions. The carbon composition ($^{13}\text{C}/^{12}\text{C}$) of the organic matter produced by plants is highly correlated with the type of photosynthetic pathway: C_3 pathway plants (cool, moist adapted) have an average $\delta^{13}\text{C}$ value of -27‰ (parts per thousand), while C_4 pathway plants (warm, dry adapted) have an average value of -14‰ (Krishnamurthy et al., 1982). Accordingly, the time series of $\delta^{13}\text{C}$ left in the loess by the plants (namely grasses) should be a proxy record of climate. Proxy data for vegetation and climate associated with humate-derived $\delta^{13}\text{C}$ values obtained from loess samples collected at the Biesel-Steinle site (fig. 27). These data suggest that the plant ecology of the Gilman Canyon Loess was dominated by C_4 -type grasses, reflecting a relatively warm, possibly dry climate (Johnson, 1993). Opal phytolith data (Fredlund et al., 1985; Fredlund and Jaumann, 1987; Johnson et al., 1994) and isotopic data (Fredlund, 1993; Johnson et al., 1994) from sites located elsewhere in the region indicate that a panicoid-dominated (C_4) grassland existed in an environment suitable for moist, temperate-adapted tall grasses during accumulation of Gilman Canyon Loess.

Although recognized at only one site in Russell County, the Gilman Canyon Formation is assumed to be present elsewhere in the county, specifically on broad interfluvies where it is buried by Peoria Loess. The Gilman Canyon Formation is not differentiated from other loesses in Russell County and is included within the map unit **QI** on the geologic map of the county (Johnson and Arbogast, 1996).

Peoria Loess. The name "Peoria" was first proposed by Leverett (1899) for deposits of an interglacial period that separates the Iowan and Wisconsinan glacial stages. When Alden and Leighton (1917) demonstrated that the Peoria is younger than the Iowan, the name "Peoria" became associated with a loess rather than a weathering interval. Several names have been used for post-Farmdalian loess in the central United States. Ruhe (1983), for example, preferred the term "late Wisconsin loess" because of correlation uncertainties from one region to another.

The Peoria Loess is typically calcareous, massive, light-yellowish-tan to buff silt that usually overlies the Loveland Loess or an approximate equivalent of the Gilman Canyon Formation. Ruhe (1983) noted three major characteristics of the Peoria, i.e., that it thins downwind of the source area, systematically fines in particle size downwind from the source area, and is time transgressive. Regarding the latter, Ruhe (1969) found that the age of the soil beneath the Peoria Loess decreases from 24,500 yr B.P. near the Missouri River to about 19,000 yr B.P. eastward across southwestern Iowa. Similarly, the base of the loess decreases in age from 25,000 to 21,000 yr B.P. along a transect in Illinois (Kleiss and Fehrenbacher, 1973). Ages from the top of the loess range from 12,500 yr B.P. in Illinois (McKay, 1979) to 14,000 yr B.P. in central Iowa (Ruhe, 1969). Deposition of the Peoria Loess was not continuous, as evinced by stratigraphic breaks such as that marked by the Jules soil in Illinois (Frye and Willman, 1973; Frye et al., 1974; Ruhe, 1976; McKay, 1979) and the soil zones recognized in Iowa (Daniels et al., 1960; Ruhe et al., 1971).

In Kansas, the Peoria Loess is a reddish, yellowish, or tan-buff homogeneous, massive, locally fossiliferous, variably calcareous coarse silt to very fine sand to medium to fine silt and clay (Frye and Leonard, 1952). The source of the silt is not certain. In a review of available data, Welch and Hale (1987) concluded that loess deposits in




FIGURE 27. Stratigraphy, percent organic matter, particle size, and magnetic susceptibility ($10^{-8} \text{ m}^3/\text{kg}$) and frequency dependence (%) of the Beisle-Steinle site in the NW sec. 27, T. 14 S., R. 11 W.

Kansas were not derived from a single source, but rather from a combination of three sources: glacial-outwash river floodplains, present sand dune fields, and erosion of the Ogallala Formation. In Phillips County, the Peoria contains moderate but variable amounts of sand with

increasing amounts of clay. According to Johnson (1993), this trend may have resulted from increased clay influx from the southwest when atmospheric circulation patterns shifted as the Laurentide ice sheet diminished (COHMAP members, 1988). Thickness of the Peoria ranges from 100




FIGURE 28. Changes with depth in humate-derived carbon isotope values ($\delta^{13}\text{C}$) determined from the backhoe trench excavated at the Biesel-Steinle site.

ft (30.5 m) along the Missouri River valley to 2 ft (0.6 m) in isolated patches elsewhere. Where accumulation is less than 2 ft (0.6 m), the Peoria is considered to be unrecognizable because it is incorporated into the existing surface soil.

Although visible soil zones such as those recognized in Illinois (Frye and Willman, 1973; Frye et al., 1974; Ruhe, 1976; McKay, 1979) and Iowa (Daniels et al., 1960; Ruhe et al., 1971) have not been recognized in Kansas, lenses of plant remains, reflecting former stable surfaces, suggest that Peoria Loess deposition was discontinuous in the region. Remains of *Picea* (cf. *glauca*), which indicate a cool, moist environment, are common. Radiocarbon ages from these materials indicate two temporal clusters of deposition: 18–17 ka and 14–13 ka. The earliest interval occurred during the last glacial maximum, while the latter represents major deglaciation (Ruddiman, 1987). Atmospheric data obtained from Greenland ice-cores indicate a significant decrease in dust at about 13 ka (Paterson and Hammer, 1987), which may have resulted in relative surface stability, soil formation, and tree establishment in the central Great Plains (Johnson, 1993).

In Russell County, the Peoria Loess is undifferentiated from the other loesses and is included within the map unit **Q1** on the geologic map of the county by Johnson and Arbogast (1996). The only exposure where a significant thickness of Peoria Loess is recognized is the Beisle-

Steinle site (fig. 25, 26). At this site, the Peoria consists dominantly of silt which was derived locally and from long-distance transport from the Platte River valley of Nebraska. In addition, organic matter decreases significantly from the underlying Gilman Canyon Formation, indicating that deposition of the Peoria was relatively continuous and rapid. The relatively low values for magnetic susceptibility and frequency dependence, a function of accumulation rates exceeding pedogenic rates, verify the relatively unweathered nature of the Peoria loess (fig. 27). Carbon isotope values (fig. 28) are smaller, indicating the C_3 plant environment characteristic of the cool, moist climate of the Late Wisconsinan. The Peoria is probably the most widespread loess in the county, however, with thick, relatively continuous deposits likely on broad interfluvies, especially those sloping to the south. Elsewhere, the Peoria probably consists of discontinuous patches that mantle heavily dissected upland terrain.

Quaternary System—Holocene Series

As defined, the Holocene comprises the last 10,000 years and began at a time of dramatic environmental change associated with disintegration of the Laurentide ice sheet. Although the Holocene is generally considered to be a diachronous geochronometric boundary with localized environmental change (e.g., Watson and Wright, 1980), recent research on the regional and subcontinental scale (e.g., Johnson and Martin, 1987; Johnson and Logan, 1990; and Johnson and May, 1992) has documented major pedogenesis, the first since Gilman Canyon time, in both alluvial and eolian/upland settings at about 10 ka.

Until recently, the Kansas Geological Survey recognized the Recent Stage of the Pleistocene Series (Bayne and O'Connor, 1968), which was defined as the last 5,000 years or the time since the end of the Valderan Substage of the Wisconsinan Stage. But the nature of the late Quaternary stratigraphic record in Kansas (Johnson, 1993) indicates that use of Recent Stage be abandoned in favor of Holocene (series status), in accordance with practice of the U.S. Geological Survey (Cohee, 1968). In a recent revision of the Kansas stratigraphic succession, the Kansas Geological Survey adopted Holocene as the name for the last ten thousand years (Baars, 1994).

Brady Soil. The Brady Soil was first named and described by Schultz and Stout (1948) at the Bignell Hill type locality, a roadcut exposure of loess and associated soils in the south valley wall of the Platte River of southwestern Nebraska. There, the Brady is developed within uppermost Peoria Loess and is overlain by the Holocene Bignell Loess. Subsequently, the soil name was adopted by researchers in Kansas (Frye and Fent, 1947; Frye and Leonard, 1949, 1951; Frye et al., 1949). The Brady Soil has been recognized in northeastern Kansas (Frye and Leonard, 1951; Caspall, 1970, 1972) but is regionally extensive only in the northwestern and west-central parts

of the state where it occurs discontinuously. Where the overlying Bignell Loess is not visibly present and the surface has not been eroded during the Holocene, the Brady is expressed as an exposed paleosol, i.e., it has pedogenically integrated any post-Bradyan loess fall.

As defined, the Brady Soil is usually dark gray to gray brown and better developed than the overlying surface soil within the Bignell. The Brady can be strongly weathered both physically and chemically (Feng et al., 1994a, 1994b), resulting in a well developed Bt horizon and concentrations of illuviated carbonate in the C horizon. Recent radiocarbon dating of the Brady in both alluvial and upland environments indicates that pedogenesis occurred from about 10,500 yr B.P. to as recently as 8,500 yr B.P. (Johnson, 1993; Johnson et al., 1993).

The Brady is recognized only at the Beisel-Steinle site (fig. 26, 27). Texturally, the Brady contains significantly more clay than the Peoria Loess, which probably reflects in situ weathering of the Brady during pedogenesis. As expected, organic matter increases markedly in the Brady from the underlying Peoria Loess. The Brady Soil is expressed in both magnetic parameters: susceptibility defines the upper part of the solum, whereas frequency dependence probably articulates the textural B horizon of the lower solum (fig. 27). Carbon isotope values (fig. 28) increase progressively during Brady time, reflecting the shift from C₃-type grasses and possibly trees to C₄-type grasses characteristic of the region today.

The Brady Soil has been recently dated at several sites in Kansas and Nebraska (Johnson, 1993). From these radiocarbon data, it appears that Brady pedogenesis occurred between about 10.5 ka and 8.5 ka, although the chronology requires appreciable refinement. The Beisel-Steinle site is only the third site in Kansas to provide radiocarbon ages for the Brady Soil (table 3). The age of $11,350 \pm 110$ yr B.P. was derived from a sample collected immediately below the Brady Soil and probably predates initiation of Brady pedogenesis by about 1,000 years. A sample collected at the perceived top of the Brady dated at $3,250 \pm 50$ yr B.P. This was originally anticipated to date to about 9,000 yr B.P. but was much younger due to the incorporation of younger or modern carbon from the adjacent surface soil, which has apparently welded to the Brady Soil.

Although the Brady Soil is recognized at only one locality in Russell County, it undoubtedly exists elsewhere in the county, most often on interfluvies adjacent to large valleys where it is buried by thin deposits of Bignell Loess. In addition, research elsewhere in the Kansas River basin (Johnson and Martin, 1987; Johnson and Logan, 1990) indicates that the contemporaneous alluvial soil is probably common in the larger stream valleys of Russell County, namely those of the Saline and Smoky Hill rivers. In Russell County, the Brady Soil is not differentiated from the loesses and is included on the geologic map within the map unit **Ql** (Johnson and Arbogast, 1996).

Bignell Loess. As with the Brady Soil, the Bignell Loess was first named and described by Schultz and Stout (1945) at the Bignell Hill type locality of the Platte River valley. Seldom more than 5 ft (1.5 m) thick, the Bignell is typically a gray or yellow-tan massive, calcaerous silt. Bignell Loess is relatively well weathered, probably because it was derived from the Brady Soil (Feng, 1991). Positive identification of this loess can not be made without the underlying Brady Soil, even though the Bignell is commonly less compact, more friable, and darker colored than the Peoria Loess. The Bignell Loess is not ubiquitous; it occurs most often as discontinuous deposits on the south side of adjacent present-day valleys. Radiocarbon and TL ages from the type section in Nebraska, the Speed roadcut in Phillips County, and other sites in the region indicate that the Bignell Loess is younger than 8,000 yr B.P. (Johnson, 1993; Maat and Johnson, 1996).

The Bignell Loess is not differentiated from the other loesses on the geologic map of Russell County (Johnson and Arbogast, 1996) and is included within the map unit **Ql**. Like the Brady Soil, the Bignell Loess was recognized only at the Beisel-Steinle site (fig. 26). The textural and organic composition of the Bignell Loess at the site (fig. 27) indicates that, compared to underlying units, the percentage of sand significantly increases in the Bignell. Organic matter also increases dramatically, which reflects the incorporation of surface soil in the loess. Magnetic susceptibility indicates that the Bignell Loess is less weathered than the Brady Soil and modern surface soil, but it is more weathered than the Peoria Loess due to one of its probable sources, the eroding Brady Soil surface (fig. 27). Although the Bignell was recognized only at this site, it probably occurs elsewhere in the county as discontinuous patches on the broad interfluvies on the south side of major stream valleys.

Fluvial deposits. Holocene alluvial deposits are pervasive in stream valleys of Russell County. They are found in two topographic positions: (1) as a major component of the fill beneath a prominent, high terrace (overlying basal late Wisconsin fill) and (2) in bottomlands below the high terrace and in smaller valleys where no high terrace is present. At localities where the Holocene alluvium comprises at least the upper component of the high terrace (fig. 29), it is mapped as **Qt** on the geologic map of the county. Where younger deposits occur (figs. 29, 30), they are designated **Qal** (Johnson and Arbogast, 1996).

Although much research (e.g., Mandel, 1988; Mandel et al., 1991; Johnson and Martin, 1987; Johnson and Logan, 1990; Martin, 1992) concerning the absolute age of Holocene alluvial deposits has been conducted in the Kansas River basin, of which streams in this study are a part, detailed information concerning the chronology of fluvial events during the last 10,000 years in Russell County has only recently emerged. A study associated with



FIGURE 29. Holocene alluvium in the Saline River valley in the NE sec. 33, T. 12 S., R. 14 W. The arrow left of center in the photograph points to a cutbank exposing fill beneath the high terrace (mapped as **Qt** on the geologic map) that extends to the left. In the middle ground, the arrow points to late Holocene alluvium (**Qal** on geologic map) adjacent to the stream channel.



FIGURE 30. Channel way, floodplain, and low terraces of the Smoky Hill River in the NW sec. 18, T. 12 S., R. 14 W. Bedrock exposed in the valley wall to the right is the Dakota Formation.




FIGURE 31. Wolf Creek basin with towns, major tributaries, and location of study sites.

an archeological investigation was conducted by May (1986) along the Saline River in the vicinity of Wilson Reservoir. May recognized three terraces, at 16 m (52 ft), 8 m (26 ft), and 3 m (10 ft) above the former channel in the upper end of the reservoir, with multiple alluvial fills being recognized in the two lower terraces. A radiocarbon age of $5,090 \pm 60$ yr B.P. (Beta-14135) was obtained on a buried soil in one fill, whereas in a younger fill a buried soil yielded an age of $1,940 \pm 70$ yr B.P. (Beta-14136). In a study of Hebrer's salt marsh upstream of Wilson Lake, floodplain alluviation and paludification began about 10 ka as indicated by a basal radiocarbon age of $9,280 \pm 210$ yr B.P. (Tx-6111) (Norman, 1990).

The most detailed reconstruction of Holocene fluvial behavior in Russell County was conducted in Wolf Creek basin (Arbogast, 1992; Arbogast and Johnson, 1994). Located primarily in northeastern Russell County, but including parts of southeastern Osborne County and northwestern Lincoln County (fig. 31), Wolf Creek basin has an area of approximately 163 km^2 (100 mi^2). Major tributaries of Wolf Creek include East Fork Wolf Creek, Coon Creek, and West Fork Wolf Creek.

Results obtained from Wolf Creek basin serve as a proxy for the late Quaternary alluvial chronology in the larger stream valleys of Russell County. Four terraces and valley fills were identified in Wolf Creek basin (fig. 32). The uppermost terrace, T-4, is an unpaired fill-top terrace

that varies in height from 27 to 18 m (88 to 59 ft) above the floodplain. Best observed at the Schoen gravel quarry in the SE sec. 18, T. 12 S., R. 10 W. (fig. 33), Lincoln County, this terrace is underlain by gravel fill (Fill I) that contains lenses of coarse silt up to 1 m (3.2 ft) thick and is mapped by Johnson and Arbogast (1996) as **Qg**. As noted earlier, the absolute age of upland gravel deposits in Russell County is unknown.

The T-3 terrace is an unpaired strath terrace, cut on the Dakota Formation 5–6 m (16–20 ft) above the modern floodplain; it is best preserved at the Dick site in the SE sec. 25, T. 11 S., R. 12 W. (fig. 34). Mapped by Johnson and Arbogast (1996) as **Kd** because of the underlying Dakota Formation, the T-3 was probably created during the time when entrenchment elevated the T-4.

The pervasive alluvial surface in Wolf Creek basin is the T-2, a paired fill-top terrace occupying approximately 95% of the valley floor. Mapped as **Qt** on the geologic map of the county (Johnson and Arbogast, 1996), this terrace extends 3–5 m (10–16 ft) above the modern floodplain and is underlain by silty early and late Holocene fills (Fills II and III, respectively). Both Fills II and III are exposed in the upper reaches of the basin and were analyzed in detail at the Naylor site (fig. 31). In the trunk valley of Wolf Creek and the lower reaches of the major tributaries, only Fill III is exposed; samples were analyzed from the Paschal and Schoen sites (fig. 31) for




FIGURE 32. Generalized stratigraphic cross section of Wolf Creek basin illustrating the relationship of the terraces (T) and floodplain with the alluvial fills in the basin.

radiocarbon age determination and for textural and chemical analyses.

The Naylor site is a 5.5-m (18-ft)-high cutbank exposure beneath the T-2 in the upper reaches of the East Fork Wolf Creek in the NW sec. 20, T. 10 S., R. 11 W., Osborne County (fig. 35). An unconformity separates Fills II and III, and three buried soils are present, one in Fill II and two in Fill III. Sediments separating the buried soils consist of well-laminated and poorly sorted coarse silt and fine sand. Each of the buried soils recognized at the site are Entisols, consisting of A-AC-C horizon sequences. A radiocarbon age of $6,770 \pm 100$ yr B.P. (Tx-6914) was obtained from the soil exposed in Fill II. The oldest of the two buried soils in Fill III, which is slightly truncated, yielded an age of $1,880 \pm 60$ yr B.P. (Tx-6962) whereas an age of $1,460 \pm 60$ yr B.P. (Tx-6961) was obtained from the uppermost, buried late Holocene soil.

Fill underlying the T-2 terrace was sampled in the upper reaches of the trunk valley of Wolf Creek at the Paschal site, a 9.5-m (31-ft)-high section of late Holocene silt and silt loam deposits located in the SW sec. 8, T. 11 S., R. 12 W., Russell County (fig. 36). At the base of the exposure, between 9.51 m and 7.93 m (31 ft and 26 ft) below the surface, is a sequence of gravel lenses separated by layers

of organic-rich silt, the thickest of which produced an age of $5,350 \pm 110$ yr B.P. (Tx-7078). Three buried soils, each consisting of A-AC-C horizons (Entisol), are also contained within Fill III at this locality. As at the Naylor site, the deposits separating the buried soils at the Paschal site are well laminated and poorly sorted. Radiocarbon ages of $1,830 \pm 70$ (Tx-7077), $1,510 \pm 80$ (Tx-7076), and $1,290 \pm 70$ yr B.P. (Tx-7075) were obtained from the lower, middle, and upper buried soils, respectively. A charcoal age of $2,060 \pm 160$ yr B.P. (Beta-2162), which correlates well with the age of the lowermost soil, was obtained from sediments just beneath the lowest buried soil in a cutbank exposure 2.5 km (1.5 mi) upstream (Johnson and Martin, 1987).

Close-interval sampling of T-2 fill in the lower reaches of the trunk valley of Wolf Creek was conducted at the Schoen cutbank site, located in the NW sec. 17, T. 12 S., R. 10 W., Lincoln County (fig. 37). At this locality, about 6.5 m (21 ft) of late Holocene fill is exposed beneath the T-2. Detailed analyses of Fill III indicate a stratigraphy and texture similar to that of the Paschal site, i.e., deposits of silt and silt loam that are separated by laminated, medium to fine sands and silt. A deposit of organic-rich silt at the base of the Schoen site yielded an age of $2,970 \pm 80$ yr B.P.



FIGURE 33. View of the T-4 terrace, underlain by Fill I, at the Schoen gravel pit in the SE sec. 18, T. 12 S., R. 10 W., Lincoln County. Fill I consists of Pleistocene gravel (mapped as **Qg** on the geologic map) that contains lenses of coarse silt.



FIGURE 34. View of the T-3 terrace at the Dick site in the SE sec. 25, T. 11 S., R. 12 W. During intensive valley entrenchment, a strath terrace was cut on the Dakota Formation (arrow) along the valley margins of Wolf Creek.




FIGURE 35. View of the Naylor site in the NW sec. 20, T. 10 S., R. 11 W., Osborne County, that shows the East Fork Wolf Creek, the T-2 terrace, and 5.5 m of alluvial fill. The fill is generally silt and silt loam in texture, poorly sorted, and well laminated. At the base of the exposure is nearly 1.5 m (4.9 ft) of early Holocene fill (Fill II), buried by 4.0 m (13.1 ft) of late Holocene alluvium (Fill III). Radiocarbon dating of the upper 5 cm (2 in) of the 4Ab, 3Ab, and 2Ab horizons yielded ages of $6,770 \pm 110$ (Tx-6914), $1,880 \pm 60$ (Tx-6962), and $1,460 \pm 60$ (Tx-7076) yr B.P., respectively.

(Tx-6795). Three buried soils, each an Entisol with A-AC-C horizons, were recognized at the Schoen site. Ages of $1,750 \pm 70$ (Tx-6959) and $1,250 \pm 60$ yr B.P. (Tx-6960) were obtained on the lowermost and uppermost buried soil, respectively. Although the intermediate soil was not dated, its stratigraphic position and morphologic similarity to the intermediate buried soil dated at the Paschal site indicate an age of about 1,500 yr B.P.

The youngest alluvial landforms in Wolf Creek basin are found in the T-1 complex, consisting of four ill-defined surfaces, and the modern floodplain (T-0). These surfaces

are less than 1,000 years old and formed as the channels of Wolf Creek basin became entrenched in conjunction with limited lateral migration. Similar entrenchment has been widely recognized (e.g., Johnson and Martin, 1987; May, 1989; Hall, 1990; Martin, 1992) in stream systems throughout the central Great Plains. Figure 38 illustrates the relationship of the T-1 complex with the T-2 at the Paschal site in plan view. A north-south transect at the Paschal site illustrates morphostratigraphic relationships of the main valley (fig. 39). The uppermost terrace in the T-1 complex is a fill-strath terrace cut on the 4Ab horizon in

Fill III alluvium. The remaining surfaces of the T-1 complex and T-0 are fill-top terraces on Fill IV, which consists of 3.5–5 m (11–16 ft) of gravel and silt, mapped as **Qal** by Johnson and Arbogast (1996). The modern

floodplain, or T-0, is poorly developed in Wolf Creek basin, consisting of isolated, narrow surfaces adjacent to the present stream channel where Fill IV is a thin veneer of silt and fine sand overlying coarse gravel.

Recent geologic history

Frye and Leonard (1952) outlined the geologic history of the Pleistocene, the period during which the present landscape of Russell County was developed. No major changes in the reconstruction are apparent from this study, but some subtle details, such as the absolute ages of depositional and erosional events have emerged. Specifically, the late Quaternary stratigraphic record, including the late Wisconsinan and Holocene, is better resolved because younger deposits are better preserved and datable by the radiocarbon dating technique.

During the Miocene and Pliocene, the existing landscape of Russell County was partially mantled by Rocky Mountain alluvium. Due to conflicting evidence, it is unclear whether deposits remaining from this time are part of the Ogallala Formation, which has been identified to

the west, north, and south of Russell County, or reworked Tertiary sediments. In spite of this uncertainty, remnants of alluvial sand and gravel, especially in the northwest part of the county, indicate that streams functioned as high-energy, braided systems during this time.

At some time during the early Pleistocene, after aggradation of stream channels with a Rocky Mountain sediment load, a period of major erosion, stream-system development, and entrenchment occurred in Russell County. No sediments from this time period have been positively identified owing to their antiquity, unconsolidated nature, and probable high topographic position. The present drainage pattern in Russell County was apparently established no later than late pre-Illinoian. This observation is supported by the stratigraphic association of



FIGURE 36. View of the Paschal site in the SW sec. 8, T. 11 S., R. 12 W., showing the T-2 terrace and about 9.5 m (31.2 ft) of late Holocene fill (Fill III). The fill at this locality consists of three sedimentary units, largely silt and silt loam in texture, that are poorly sorted and well laminated. At the base of the exposure are alternating layers of coarse gravel and silt. Radiocarbon-age determination on an organic-rich silt layer 8.6 m (28.2 ft) below the surface (arrow) yielded an age of $5,350 \pm 110$ (Tx-7078). Radiocarbon ages of $1,830 \pm 70$ (Tx-7077), $1,510 \pm 80$ (Tx-7076), and $1,290 \pm 70$ (Tx-7075) yr B.P. were obtained on the upper 5 cm (2 in) of the 4Ab, 3Ab, and 2Ab horizons, respectively.




FIGURE 37. The Schoen site in the NW sec. 17, T. 12 S., R. 10 W., Lincoln County, with Wolf Creek, the T-2 terrace and approximately 6.5 m (21.3 ft) of alluvial fill. The fill at this locality is late Holocene in age and is composed of three sedimentary units that are poorly sorted and well laminated. Radiocarbon-age determinations on the upper 5 cm (2 in) of an organic-rich silt layer at the base of the exposure and the 4Ab and 2Ab horizons yielded ages of $2,970 \pm 80$ (Tx-6795), $1,750 \pm 70$ (Tx-6959), and $1,250 \pm 60$ (Tx-6960) yr B.P., respectively.

pre-Illinoian and younger alluvial deposits with terraces and by the deposition of upland loess deposits. Late pre-Illinoian deposits, possibly correlating to the Grand Island and Sappa Formations, may be preserved on valley side walls of the Smoky Hill River, Saline River, and their major tributaries (e.g., Wolf Creek).

Illinoian time brought renewed incision in Russell County, removing much of the earlier deposits both laterally and vertically. Wolf Creek, for example, entrenched at least 17 m (56 ft) below the base of the pre-Illinoian fill. During one or more periods of valley

aggradation during the Illinoian, deposition of sand and gravel (possibly correlative with the Crete Formation) occurred in the major stream valleys whereas the fine fraction, transported by the wind, accumulated on the uplands as the Loveland Loess. Several brief periods of landscape stability, resulting in a series of paleosols, probably occurred during Illinoian loess deposition. The most intensive period of landscape stability, resulting in the extremely well-developed Sangamon Soil, took place as deposition of Loveland Loess terminated. Gaps in the stratigraphic record are too extensive to precisely recon-

struct the sequence of events in Russell County during the early and early middle Wisconsin, although a period of regional, post-Sangamon erosion has been documented (Johnson, 1993).

During the late middle and late Wisconsin (latest Altonian and Farmdalian), deposition of a thin loess mantle (Gilman Canyon Formation) probably occurred throughout Russell County on the uplands, valley side slopes, and bottomlands. Deposition of the loess was sufficiently slow that pedogenesis could incorporate new accumulations. Potentially, two or more distinct periods of landscape stability occurred, but individual A horizons have been obscured through bioturbation.

Late Wisconsin (Woodfordian) time was a period of large-scale Peoria Loess deposition in Russell County when the Gilman Canyon Formation was slowly and conformably buried. Terminal ages on the Gilman Canyon and basal A horizon ages from the Brady Soil indicate that Peoria Loess accumulated between 20 ka and 10 ka, although reduction in the depositional rate may have occurred at 18–17 ka and 14–13 ka as suggested by regional botanical, pedologic, and geomorphic evidence. Streams in Russell County entrenched to a level beneath the present floodplain from about 13 ka to shortly before 10.5 ka, when a well-developed soil, chronologically



FIGURE 38. Aerial photograph of the T-2 terrace (mapped as *Qt* on the geologic map) and T-1 terrace complex (mapped as *Qal*) near the Paschal site in sec. 8, T. 11 S., R. 12 W. In this reach of the stream, approximately 25 percent of T-2 fill has been removed by lateral erosion in conjunction with late Holocene stream entrenchment, leaving isolated fragments of the T-2 terrace.




FIGURE 39. Cross section of T-2 (mapped as *Qt* on the geologic map), T-1 complex, and T-0 (floodplain) (T-1 and T-0 mapped as *Qal*) at the Paschal site in SW sec. 8, T. 11 S., R. 12 W. The T-2 terrace is underlain by Fill III. The T-1 terrace complex consists of at least four ill-defined surfaces created by stream entrenchment and lateral channel migration in the last 1,000 years. The uppermost surface in the T-1 terrace complex is a cut-terrace on Fill III alluvium, while the remaining surfaces are fill-terraces underlain by Fill IV alluvium that caps Fill III.

equivalent to the upland Brady Soil, formed as it did elsewhere in the central Great Plains.

Except for deposition of a thin, discontinuous cover of Bignell Loess on the uplands, slumping in steep valley walls, and isolated sinkhole formation, the majority of Holocene landscape change in Russell County has been confined to stream systems. Reconstruction of Holocene cut-and-fill sequences elsewhere in the Kansas River basin indicates that Wolf Creek basin in northeastern Russell County can be confidently used as a proxy for events in large streams elsewhere in the county, although subtle differences in the timing or magnitude of events may exist.

During the early Holocene, from approximately 10.5 ka to 6 ka, aggradation of floodplains in Russell County was widespread. Deposition was episodic, with a brief period of soil formation recognized in Wolf Creek basin at 6.7 ka. Sometime during the middle Holocene (6–5 ka), a major erosional event occurred that flushed most early Holocene alluvium from Wolf Creek. Evidence elsewhere in the Kansas River basin suggests that erosion during the middle Holocene was probably not as inten-

sive in the Saline and Smoky Hill River valleys. Aggradation of floodplains in Russell County was renewed at 5 ka and lasted episodically until approximately 1 ka. Evidence from Wolf Creek indicates that brief periods of floodplain stability occurred at 1.8 ka, 1.5 ka, and 1.2 ka in the county. Recent entrenchment to a depth of 5–9 m (16–29 ft) in the last 1 ka has elevated a prominent terrace throughout Russell County and resulted in a series of ill-defined landforms in the bottomlands adjacent to the stream channel.

Acknowledgments

We are grateful to the many landowners who provided access to their land and information concerning local geology, soils, landforms, and associated historical events. Funding for radiocarbon dating was provided by grants from The University of Kansas General Research Fund and Sigma Xi. D. W. May of the University of Northern Iowa and D. E. Hattin of Indiana University kindly served as manuscript reviewers and provided many valuable comments.

References

- Aguirre, E., and Pasini, G., 1985, The Pliocene-Pleistocene boundary: Episodes, v. 8, p. 116–120
- Alden, W. C., and Leighton, M. M., 1917, The Iowan drift—a review of the evidence of the Iowan stage of glaciation: Iowa Geological Survey, Annual Report, v. 26, p. 49–212
- Allan, T. H., and Valerius, M. M., 1929, Fairport oil field, Russell County, Kansas; *in*, Structure of Typical American Oil Fields: Association of American Petroleum Geologists, Tulsa, Oklahoma, v. 1, p. 35–48
- Arbogast, A. F., 1992, Late Quaternary evolution of a small basin in the Kansas River system—Wolf Creek: Kansas Geological Survey, Open-file Report, no. 92-8, 152 p.
- Arbogast, A. F., and Johnson, W. C., 1994, Climatic implications of the late Quaternary alluvial record of a small drainage basin in the central Great Plains: Quaternary Research, v. 41, p. 298–305
- Baars, D., 1994, Classification of rocks in Kansas: Kansas Geological Survey, stratigraphic chart
- Bass, N. W., 1926, Geologic investigations in western Kansas: Kansas Geological Survey, Bulletin 11, 96 p.
- Bayne, C. K., and O'Connor, H. G., 1968, Quaternary System; *in*, The Stratigraphic Succession in Kansas, D. E. Zeller, ed.: Kansas Geological Survey, Bulletin 189, p. 59–67
- Bettis, E. A., III, ed., 1990, Holocene alluvial stratigraphy and selected aspects of the Quaternary history of western Iowa: Midwestern Friends of the Pleistocene, University of Iowa, Quaternary Studies Group Contribution 36, 197 p.
- Boellstorff, J. D., 1973, Fission-track ages of Pleistocene volcanic ash deposits in the central Plains, U.S.A.: Isochron/ West, v. 8, p. 39–43
- _____, 1974, Fission-track ages of Pearlette family ash beds—comment: Geology, v. 2, p. 21
- _____, 1976, The succession of late Cenozoic ashes in the Great Plains—a progress report: Kansas Geological Survey, Guidebook Series 1, p. 37–71
- Bramlette, M. N., 1925, A subsurface correlation of the stratigraphic units from Russell County to Marion County, Kansas: Kansas Geological Survey, Bulletin 10, pt. 2, p. 87–93
- Burgat, V. A., and Taylor, W. K., 1972, Highway subsidence caused by salt solutioning (abstract): Association of English Geologists, Annual Meeting, Programs and Abstracts 15, p. 20
- Bushue, L. J., Fehrenbacher, J. B., and Ray, B. W., 1974, Exhumed paleosols and associated modern till soils in western Illinois: Soil Science Society of America, Proceedings, v. 34, p. 665–669
- Carter, B. J., and Ward, P. A., III, 1991, A prehistory of the Plains border region: Guidebook, South-central Friends of the Pleistocene, Agronomy Department, Oklahoma State University, Stillwater, 121 p.
- Carter, B. J., Ward, P. A., III, and Shannon, J. T., 1990, Soil and geomorphic evolution within the Rolling Red Plains using Pleistocene volcanic ash deposits: Geomorphology, v. 3, p. 471–488
- Caspall, F. C., 1970, The spatial and temporal variations in loess deposition in northeastern Kansas: Ph.D. dissertation, University of Kansas, Lawrence, 294 p.
- _____, 1972, A note on the origin of the Brady paleosol in northeastern Kansas: Proceedings of the Association of American Geographers, v. 4, p. 19–24
- Clark, P. U., Clague, J. J., Curry, B. B., Dreimanis, A., Hicock,

- S. R., Miller, G. H., Berger, G. W., Eyles, N., Lamothe, M., Miller, B. B., Mott, R. J., Oldale, R. N., Stea R. R., Szabo, J. P., Thorleifson, L. H., and J.-S. Vincent, 1993, Initiation and development of the Laurentide and Cordilleran ice sheets following the last interglaciation: *Quaternary Science Reviews*, v. 12, p. 79–114
- Cobban, W. A., 1949, Stratigraphy of the Colorado and Montana groups (Upper Cretaceous) of the central and northern Great Plains with descriptions of the Colorado *Scaphites*: Ph.D. dissertation, Johns Hopkins University, Baltimore, Maryland, 408 p.
- Cobban, W. A., and Reeside Jr., J. B., 1952, Correlation of the Cretaceous formations of the Western Interior of the United States: *Geological Society of America, Bulletin* 63, p. 1,011–1,044
- Cohee, G. V., 1968, Holocene replaces Recent in nomenclature usage of the U.S. Geological Survey: *American Association of Petroleum Geologists, Bulletin* 52, p. 852
- COHMAP members, 1988, Climatic changes of the last 18,000 years—observations and model simulations: *Science*, v. 241, p. 1,043–1,052
- Condra, G. E., Reed, E. C., and Gordon, E. D., 1947, Correlation of the Pleistocene deposits of Nebraska: *Nebraska Geological Survey, Bulletin* 15, 71 p.
- _____, 1950, Correlation of the Pleistocene deposits of Nebraska: *Nebraska Geological Survey, Bulletin* 15-A, 74 p.
- Curry, B. B., and Follmer, L. R., 1992, The last interglacial-glacial transition in Illinois—123–125 ka; *in*, *The Last Interglacial-glacial Transition in North America*, P. U. Clark and P. D. Lea, eds.: *Geological Society of America, Special Paper* 270, p. 71–88
- Dane, C. H., and Pierce, W. G., 1933, Geology and oil and gas prospects in part of eastern Colorado: U.S. Department of Interior Press, memo 72215, 8 p.
- Daniels, R. B., Handy, R. L., and Simonson, G. H., 1960, Dark-colored bands in the thick loess of western Iowa: *Journal of Geology*, v. 67, p. 114–119
- Darton, N. H., 1899, Preliminary report on the geology and water resources of Nebraska west of the One Hundred and Third Meridian: U.S. Geological Survey, 19th Annual Report, pt. 4, Hydrology, p. 719–785
- _____, 1904, Comparison of the stratigraphy of the Black Hills, Bighorn Mountains, and Rocky Mountain front range: *Geological Society of America, Bulletin* 15, p. 379–448
- _____, 1920, Description of the Syracuse and Lakin quadrangles: U.S. Geological Survey, Atlas Folio Serial 212, 10 p.
- Diekmeyer, E. C., 1994, Characterizations and paleoclimatic inferences from the post-Illinoian stratigraphic sequences at two central Great Plains sites: M.S. thesis, Department of Geography, University of Kansas, Lawrence, 84 p.
- Dubins, I. M., 1947, The petrography, geochemistry, and economic utilization of the Fort Hays chalk in Kansas: M.S. thesis, Department of Geology, University of Kansas, Lawrence, 108 p.
- Elias, M. K., 1931, The geology of Wallace County, Kansas: *Kansas Geological Survey, Bulletin* 18, 254 p.
- _____, 1932, Grasses and other plants from the Tertiary rocks of Kansas and Colorado: *University of Kansas, Science Bulletin* 20, no. 20, p. 333–367
- _____, 1935, Tertiary grasses and other prairie vegetation from the High Plains of North America: *American Journal of Science*, series 5, v. 29, p. 24–33
- _____, 1942, Tertiary prairie grasses and other herbs from the High Plains: *Geological Society of America, Special Paper (Regular Studies)* 41, 176 p.
- Feng, Zhao-dong, 1991, Temporal and spatial variations in the loess depositional environment of central Kansas during the past 400,000 years: Ph.D. dissertation, University of Kansas, Lawrence, 250 p.
- Feng, Zhao-dong, Johnson, W. C., Lu, Yan-chou, and Ward, P. A. III, 1994a, Climatic signals from loess-soil sequences in the central Great Plains, U.S.A.: *Palaeogeography, Palaeoclimatology, Palaeoecology* 110, p. 345–358
- Feng, Zhao-dong, Johnson, W. C., Sprowl, D. R., and Lu, Yan-chou, 1994b, Loess accumulation and soil formation in central Kansas, United States, during the past 400,000 years: *Earth Surface Processes and Landforms*, v. 19, p. 55–67
- Follmer, L. R., 1979, A historical review of the Sangamon soil; *in*, *Wisconsinan, Sangamonian, and Illinoian Stratigraphy in Central Illinois*: *Illinois State Geological Survey, Guidebook* 13, p. 79–91
- Forman, S. L., 1990, Chronologic evidence for multiple episodes of loess deposition during the Wisconsinan and Illinoian in the midcontinent, U.S.A. (abstract): *Geological Society of America, Abstracts with Programs*, v. 22, no. 7, p. A86
- Forman, S. L., Bettis, E. A., Kemmis, T. L., and Miller, B. B., 1992, Chronological evidence for multiple periods of loess deposition during the late Pleistocene in Missouri and Mississippi River valleys, United States—implications for the activity of the Laurentide Ice Sheet: *Palaeogeography, Palaeoclimatology, Palaeoecology*, v. 93, p. 71–83
- Franks, P. C., 1967, Petrology and stratigraphy of the Kiowa and Dakota formations (basal Cretaceous), north-central Kansas: Ph.D. dissertation, Department of Geology, University of Kansas, Lawrence, 312 p. (2 volumes)
- Fredlund, G. G., 1993, Paleoenvironmental interpretations of stable carbon, hydrogen, and oxygen isotopes from opal phytoliths, Eustis Ash Pit, Nebraska: *MASCA Research Papers in Science and Archaeology*, v. 10, p. 37–46
- Fredlund, G. G., and Jaumann, P. J., 1987, Late Quaternary palynological and paleobotanical records from the central Great Plains; *in*, *Quaternary Environments of Kansas*, W. C. Johnson, ed.: *Kansas Geological Survey, Guidebook Series* 5, p. 167–168
- Fredlund, G. G., Johnson, W. C., and Dort Jr., W., 1985, A preliminary analysis of opal phytoliths from the Eustis ash pit, Frontier County, Nebraska: *Nebraska Academy of Sciences, Institute for Tertiary-Quaternary Studies, TER-QUA Symposium Series*, v. 1, p. 147–162
- Frye, J. C., and Brazil, J. J., 1943, Ground water in the oil-field areas of Ellis and Russell counties, Kansas, with analyses by Howard Stoltenberg: *Kansas Geological Survey, Bulletin* 50, 104 p.
- Frye, J. C., and Fent, O. S., 1947, The late Pleistocene loesses of central Kansas: *Kansas Geological Survey, Bulletin* 70, pt. 3, p. 29–52
- Frye, J. C., and Leonard, A. B., 1949, Pleistocene stratigraphic sequence in northeastern Kansas: *American Journal of Science*, v. 247, p. 883–899
- _____, 1951, Stratigraphy of late Pleistocene loess of Kansas: *Journal of Geology*, v. 59, no. 4., p. 287–305

- _____. 1952, Pleistocene geology of Kansas: Kansas Geological Survey, Bulletin 99, 230 p.
- _____. 1954, Significant new exposures of Pleistocene deposits at Kirwin, Phillips County, Kansas: Kansas Geological Survey, Bulletin 109, pt. 3., p. 29–48
- _____. 1965, Quaternary of the southern Great Plains; *in*, The Quaternary of the United States—A Review Volume for VII Congress of the International Association for Quaternary Research, H. E. Wright Jr., and D. G. Frye, eds.: Princeton University Press, Princeton, New Jersey, p. 203–216
- Frye, J. C., and Willman, H. B., 1973, Wisconsinan climatic history interpreted from Lake Michigan lobe deposits and soils; *in*, The Wisconsinan Stage, R. F. Black, R. P. Goldthwait, and H. B. Willman, eds.: Geological Society of America, Memoir 136, p. 135–152
- Frye, J. C., Leonard, A. B., and Leonard, A. R., 1951, Western Kansas; *in*, Road Log, Pleistocene Field Conference of June, 1951, part 2: Pleistocene Field Conference, Guidebook 3, p. WK-1–WK-18
- Frye, J. C., Plummer, N., Runnels, R. T., and Hladik, W. B., 1949, Ceramic utilization of northern Kansas Pleistocene loesses and fossil soils: Kansas Geological Survey, Bulletin 82, pt. 3, p. 49–124
- Frye, J. C., Leonard, A. B., Willman, H. B., Glass, H. D., and Follmer, L. R., 1974, The late Woodfordian Jules soil and associated molluscan faunas: Illinois State Geological Survey, Circular 486, 11 p.
- Fulton, R. J., 1984, Summary—Quaternary stratigraphy of Canada: Geological Survey of Canada, Paper 84-10, p. 1–5
- _____. 1986, Quaternary stratigraphy of Canada; *in*, V. Sibrava, D. Q. Bowen, and G. M. Richmond, eds., Quaternary Glaciations in the Northern Hemisphere, Quaternary Science Reviews, v. 5, p. 207–209
- Gilbert, G. K., 1896, The underground water of the Arkansas Valley in eastern Colorado: U.S. Geological Survey, 17th Annual Report, pt. 2, p. 551–601
- Gustavson, T. C., and Winkler, D. A., 1988, Depositional facies of the Miocene-Pliocene Ogallala Formation, northwestern Texas and eastern New Mexico: *Geology*, v. 16, no. 3, p. 203–206
- Hall, R. D., 1973, Sedimentation and alteration of loess in southwestern Indiana: Ph.D. dissertation, Indiana University, Bloomington, 103 p.
- Hall, S. A., 1990, Channel trenching and climatic change in the southern Great Plains: *Geology*, v. 18, p. 342–345
- Hansen, T. J., 1977, Dissolution in the Hutchinson Salt Member of the Wellington Formation near Russell, Kansas: M.S. thesis, Department of Geology, Kansas State University, Manhattan, 61 p.
- Hatfield, C. B., 1961, Paleocology of the Graneros Shale (Upper Cretaceous) in Kansas: M.A. thesis, Department of Geology, Indiana University, Bloomington, 97 p.
- Hattin, 1962, Stratigraphy of the Carlile Shale (Upper Cretaceous) in Kansas: Kansas Geological Survey, Bulletin 156, 155 p.
- _____. 1965a, Stratigraphy of the Graneros Shale (Upper Cretaceous) in central Kansas: Kansas Geological Survey, Bulletin 178, 83 p.
- _____. 1965b, Upper Cretaceous stratigraphy, paleontology, and paleoecology of western Kansas: Geological Society of America, Field Conference Guidebook, Annual Meeting, Kansas City, Missouri, 69 p.
- _____. 1967, Stratigraphic and paleoecologic significance of macroinvertebrate fossils in the Dakota Formation (Upper Cretaceous) of Kansas; *in*, Essays in Paleontology and Stratigraphy—R. C. Moore Commemorative Volume, Curt Teichert and E. L. Yochelson, eds.: Lawrence, University Press of Kansas, Department of Geology, Special Publication 2, p. 570–589
- _____. 1971, Widespread, synchronously deposited, burrow-mottled limestone beds in Greenhorn Limestone (Upper Cretaceous) of Kansas: Kansas Geological Survey, Bulletin 202, pt. 2, 11 p.
- _____. 1975, Stratigraphy and depositional environment of Greenhorn Limestone (Upper Cretaceous) of Kansas: Kansas Geological Survey, Bulletin 209, 128 p.
- _____. 1982, Stratigraphy and depositional environment of Smoky Hill Chalk Member, Niobrara Chalk (Upper Cretaceous) of the type area, western Kansas: Kansas Geological Survey, Bulletin 225, 108 p.
- Hattin, D. E., and Cobban, W. A., 1965, Road logs; *in*, Upper Cretaceous Stratigraphy, Paleontology, and Paleoecology of Western Kansas—A Field Conference Guidebook for the Annual Meetings, D. E. Hattin: Kansas Geological Survey, [guidebook for] Geological Society of America, Annual Meeting, p. 29–47
- _____. 1977, Upper Cretaceous stratigraphy, paleontology, and paleoecology of western Kansas: *Mountain Geologist*, v. 14, no. 3–4, p. 175–217
- Hattin, D. E., and Siemers, C. T., 1978, Guidebook, Upper Cretaceous stratigraphy and depositional environments of western Kansas: Kansas Geological Survey, Guidebook Series 3, 102 p.
- Hay, R., 1889, Horizon of the Dacotah lignite: Kansas Academy of Sciences, Transactions, v. 11, p. 5–8
- Hayden, F. V., 1872, Sketch of the geological formations along the route of the Union Pacific Railway, eastern division; *in*, Final Report of the U.S. Geological Survey of Nebraska and Portions of the Adjacent Territories: U.S. Congress, 42nd, 1st session, House Executive Document 19, p. 66–69
- Hedberg, H. D., 1926, The effect of gravitational compaction on the structure of sedimentary rocks: *American Association of Petroleum Geologists, Bulletin* 10, no. 11, p. 1,035–1,072
- Hibbard, C. W., Frye, J. C., and Leonard, A. B., 1944, Reconnaissance of Pleistocene deposits in north-central Kansas: Kansas Geological Survey, Bulletin 52, pt. 1, p. 1–28
- Hintze, F. F., 1928, Discoverer of Gorum oil district, Kansas (discussion): *American Association of Petroleum Geologists, Bulletin* 12, no. 4, p. 443
- Hopkins, D. M., 1975, Time-stratigraphic nomenclature for the Holocene Epoch: *Geology*, v. 3, p. 10
- Izett, G. A., 1981, Volcanic ash beds—recorders of upper Cenozoic silicic pyroclastic volcanism in the western United States: *Journal of Geophysical Research*, v. 86, p. 10,200–10,222
- _____. 1982, The Bishop ash bed and some older, compositionally similar, ash beds in California, Nevada, and Utah: U.S. Geological Survey, Open-file Report 82-582, 44 p.
- Izett, G. A., and Wilcox, R. E., 1982, Map showing localities and inferred distributions of the Huckelberry Ridge, Mesa Falls, and Lava Creek ash beds (Pearlette family ash beds) of Pliocene and Pleistocene age in the western United States and southern Canada: U.S. Geological Survey, Miscellaneous Investigations, Map I-1325, 1 sheet, scale 1:4,000,000

- Izett, G. A., Wilcox, R. E., Powers, H. A., and Desborough, G. A., 1970, The Bishop ash bed—a Pleistocene marker bed in the western United States: *Quaternary Research*, v. 1, p. 121–132
- Jantz, D. R., Wehmueller, W., and Owens, H. D., 1982, Soil survey of Russell County, Kansas: U.S. Department of Agriculture Soil Conservation Service, 103 p.
- Jewett, J. M., 1959, Graphic column and classification of rocks in Kansas: Kansas Geological Survey, 1 sheet
- Johnson, R. G., 1982, Brunhes-Matuyama magnetic reversal dated at 790,000 yr B.P. by marine-astronomical correlations: *Quaternary Research*, v. 17, p. 135–147
- Johnson, W. C., 1993, Surficial geology and stratigraphy of Phillips County, Kansas, with emphasis on the Quaternary Period: Kansas Geological Survey, Technical Series 1, 66 p.
- Johnson, W. C., and Arbogast, A. F., 1996, Geologic map of Russell County, Kansas: Kansas Geological Survey, Map Series M-37, 1 sheet, scale 1:50,000
- Johnson, W. C., and Logan, B., 1990, Geoarcheology of the Kansas River basin, central Great Plains; *in*, *Archaeological Geology of North America*, N. P. Lasca and J. Donahue, eds.: Geological Society of America, Decade of North American Geology, Centennial Special Volume 4, p. 267–299
- Johnson, W. C., and Martin, C. W., 1987, Holocene alluvial-stratigraphic studies from Kansas and adjoining states of the east-central Plains; *in*, *Quaternary Environments of Kansas*, W. C. Johnson, ed.: Kansas Geological Survey, Guidebook Series 5, p. 109–122
- Johnson, W. C., and May, D. W., 1992, The Brady geosol as an indicator of the Pleistocene/Holocene boundary in the central Great Plains (abstract): American Quaternary Association, Programs and Abstracts, p. 69
- Johnson, W. C., Bozarth, S., and Diekmeyer, E. C., 1994, Paleoenvironmental reconstruction via opal phytolith and carbon isotope analyses of late Wisconsinan loess—geoarchaeological investigations on Fort Riley, Riley and Geary counties, Kansas: Kansas Geological Survey, Open-file Report 94-38, 60 p.
- Johnson, W. C., May, D. W., and Souders, V. L., 1990, Age and distribution of the Gilman Canyon Formation of Nebraska and Kansas (abstract): Geological Society of America, Abstracts with Programs, v. 22, p. A87
- Johnson, W. C., May, D. W., and Valastro, S., 1993, A 36,000-year chrono-, bio- and magneto-stratigraphic record from loess of south-central Nebraska (abstract): Association of American Geographers, 89th Annual Meeting, Abstracts with Programs, p. 115
- Johnson, W. H., Hansel, A. K., Follmer, L. R., and Curry, B. B., 1991, Later Quaternary temporal classification in Illinois—geochronologic or diachronic?: Geological Society of America, Abstracts with Programs, v. 23, no. 3, p. 19–20
- Kleiss, H. J., and Fehrenbacher, J. G., 1973, Loess distribution as revealed by mineral variations: Soil Science Society of America, Proceedings, v. 37, p. 291–295
- Knapp, R. W., and Steeples, D. W., 1981, Investigation of salt dissolution collapse using high-resolution Mini-SOSIE reflection seismology (abstract): *Eos*, v. 62, no. 45, p. 954–955
- Knapp, R. W., Steeples, D. W., Miller, R. D., and McElwee, C. D., 1989, Seismic-reflection surveys at sinkholes in central Kansas; *in*, *Geophysics in Kansas*, D. W. Steeples, ed.: Kansas Geological Survey, Bulletin 226, p. 95–116
- Krishnamurthy, R. V., Deniro, M. J., and Pant, R. K., 1982, Isotope evidence for Pleistocene climatic changes in Kashmir, India: *Nature*, v. 298, p. 640–641
- Landes, K. K., Amoroso, J. J., Charlesworth, L. J., Jr., Heany, F., and Lesperance, P. J., 1960, Petroleum resources in basement rocks: Association of American Petroleum Geologists, Bulletin 44, no. 10, p. 1,682–1,691
- Leigh, D. S., 1991, Origin and paleoenvironment of the upper Mississippi Valley Roxana silt: Ph.D. dissertation, University of Wisconsin, Madison, 186 p.
- _____, 1994, Roxana silt of the upper Mississippi Valley—lithology, source, and paleoenvironment: Geological Society of America, Bulletin 106, p. 430–442
- Leigh, D. S., and Knox, J. C., 1993, AMS radiocarbon age of the upper Mississippi Valley Roxana silt: *Quaternary Research*, v. 39, p. 282–289
- Leverett, F., 1899, The Illinois glacial lobe: U.S. Geological Survey, Monograph 38, 817 p.
- Logan, W. N., 1897, The Upper Cretaceous of Kansas: University Geological Survey of Kansas, v. 2, p. 202–234
- _____, 1898, Part VIII, The invertebrates of the Benton, Niobrara, and Fort Pierre Groups, University Geological Survey of Kansas, v., no 4, p. 433–583
- _____, 1899, Some additions to the Cretaceous invertebrates of Kansas: Kansas University Quarterly, v. 8, series A, p. 87–98
- Lugn, A. L., 1934, Pleistocene geology of Nebraska: Nebraska State Museum, Bulletin 41, v. 1, pt. 1, p. 319–356
- _____, 1935, The Pleistocene geology of Nebraska: Bulletin of the Nebraska Geological Survey, series 2, v. 10, 223 p.
- Maat, P., and Johnson, W. C., 1996, Thermoluminescence and new ^{14}C age estimates for late Quaternary loesses in south-western Nebraska: *Geomorphology* (in press)
- Mandel, R. D., 1988, Geomorphology of the Smoky Hill River valley at Kanopolis Lake, north-central Kansas; *in*, *An Archaeological and Geomorphological Survey of Kanopolis Lake, North-central Kansas*, L. J. Schmits, ed.: Environmental Systems Analysis, Inc., p. 49–72
- Mandel, R. D., Reynolds, J. D., Williams, B. G., and Wulfkuhle, V. A., 1991, Upper Delaware River and tributaries watershed—results of geomorphological and archeological studies in Atchison, Brown, Jackson, and Nemaha counties, Kansas: Kansas State Historical Society, Contract Archeological Publication 9, Topeka, Kansas, 130 p.
- Martin, C. W., 1992, Late Quaternary sedimentation and paleoenvironmental change in the Republican River basin, south-central Nebraska: *Quaternary Research*, v. 37, p. 315–322
- Martinson, D. G., Pisias, N. G., Hays, J. D., Imbrie, J., Moore, T. C., Jr., and Shackleton, N. J., 1987, Age dating and the orbital theory of the Ice Ages—development of a high-resolution 0 to 300,000-year chronology: *Quaternary Research*, v. 27, p. 1–29
- May, D. W., 1986, Geomorphology; *in*, *Along the Pawnee Trail—Cultural Resources Survey and Testing*, Wilson Lake, Kansas, D. J. Blakeslee, R. Blasing, and H. Garcia, eds.: U.S. Army Corps of Engineers (Kansas City District), report no. DACW41-85-C-0135, p. 72–86
- _____, 1989, Holocene alluvial fills in the South Loup valley, Nebraska: *Quaternary Research*, v. 32, p. 117–120
- May, D. W., and Souders, V. L., 1988, Radiocarbon ages for the Gilman Canyon Formation in Dawson County, Nebraska

- (abstract): Nebraska Academy of Sciences, Proceedings, p. 47–48
- McCraw, D. J., and Autin, W. J., 1989, Lower Mississippi Valley loesses: Field guide, 1989 Mississippi Valley Loess Tour, INQUA Commission on Loess and the North American Loess Working Group
- McKay, E. D., 1979, Stratigraphy of Wisconsinan and older loesses in southwestern Illinois; *in*, Geology of Western Illinois: Illinois State Geological Survey, Guidebook 13, p. 95–108
- Meek, F. B., and Hayden, F. V., 1862, Descriptions of new Lower Silurian (Primordial), Jurassic, Cretaceous, and Tertiary fossils, collected in Nebraska, etc.: Philadelphia Academy of Natural Sciences, Proceedings, v. 13, p. 415–447
- Merriam, D. F., 1957, Subsurface correlation and stratigraphic relation of rocks of Mesozoic age in Kansas: Kansas Geological Survey, Oil and Gas Investigations, v. 14, 25 p.
- Moore, R. C., Frye, J. C., Jewett, J. M., Lee, W., and O'Connor, H. G., 1951, The Kansas rock column: Kansas Geological Survey, Bulletin 89, 132 p.
- Morrison, R. B., 1965, Principles of Quaternary soil stratigraphy; *in*, Quaternary Soils, R. B. Morrison and H. E. Wright, Jr., eds.: INQUA VII Congress, Proceedings, v. 9, p. 1–69
- _____, 1987, Long-term perspective-changing rates and types of Quaternary surficial processes-erosion-deposition-stability cycles; *in*, Geomorphic Systems of North America, W. L. Graf, ed.: Geological Society of America, Decade of North American Geology, Centennial Special Volume 2, p. 167–176
- Moss, R. G., 1932, The geology of Ness and Hodgeman counties, Kansas: Kansas Geological Survey, Bulletin 19, 48 p.
- Mudge, B. F., 1876, Notes on the Tertiary and Cretaceous periods of Kansas: U.S. Geological and Geographical Survey of the Territories, F. V. Hayden, Bulletin 2, no. 3, p. 211–221
- Naeser, C. W., Izett, G. A., and Wilcox, R. E., 1973, Zircon fission-track ages of Pearllette family ash beds in Meade County, Kansas: Geology, v. 1, p. 187–189
- Nickel, Jr., G., 1972, Development and evaluation of field experience with a field guide to the study of Lower and middle Cretaceous rocks of Kansas in the Kanopolis and Wilson Reservoir area: M.S. thesis, Division of Physical Sciences, Emporia State University, Emporia, Kansas, 93 p.
- North American Commission on Stratigraphic Nomenclature, 1983, North American Stratigraphic Code: American Association of Petroleum Geologists, Bulletin 67, p. 841–875
- Norman, T. O., 1990, The natural history of Heberer's salt marsh, Russell County, Kansas: unpublished M.S. thesis, Fort Hays State University, Fort Hays, Kansas, 111 p.
- Oviatt, C. G., Karlstrom, E. T., and Ransom, M. D., 1988, Pleistocene loess, buried soils, and thermoluminescence dates in an exposure near Milford Lake, Geary County, Kansas (abstract): Geological Society of America, Abstracts with Programs, v. 20, no. 2, p. 125–126
- Paterson, W. S. B., and Hammer, C. U., 1987, Ice core and other glaciological data; *in*, North America and Adjacent Oceans During the Last Deglaciation, W. F. Ruddiman and H. E. Wright Jr., eds.: Geological Society of America, Decade of North American Geology, The Geology of North America, v. K-3, p. 91–109
- Plummer, N. V., and Romary, J. F., 1942, Stratigraphy of the pre-Greenhorn Cretaceous beds of Kansas: Kansas Geological Survey, Bulletin 41, pt. 9, p. 313–348
- Reed, E. E., and Dreeszen, V. H., 1965, Revision of the classification of the Pleistocene deposits of Nebraska: Nebraska Geological Survey, Bulletin 23, 65 p.
- Richmond, G. M., and Fullerton, D. S., 1986, Introduction to Quaternary glaciations in the United States of America; *in*, Quaternary Glaciations in the Northern Hemisphere, V. Sibrava, D. Q. Bowen, and G. M. Richmond, eds.: Quaternary Science Reviews, v. 5, p. 3–10
- Riggs, C. H., Huff, R. V., and Ward, D. C., 1963, Petroleum engineering study of the Hall-Gurney field, Russell County, Kansas: Production Monthly, v. 24, no. 7, p. 18–19, 22–25
- Ross, J. A., and Wong, R. K.-W., 1989a, Oil and gas fields, Beloit quadrangle in Kansas: Kansas Geological Survey, Map Series no. 18-2, 1 sheet, scale 1:250,000
- _____, 1989b, Oil and gas fields, Great Bend quadrangle in Kansas: Kansas Geological Survey, Map Series no. 18-7, 1 sheet, scale 1:250,000
- _____, 1990a, Oil and gas fields, Beloit quadrangle in Kansas: Kansas Geological Survey, Map Series no. 22-2, 1 sheet, scale 1:250,000
- _____, 1990b, Oil and gas fields, Great Bend quadrangle in Kansas: Kansas Geological Survey, Map Series no. 22-7, 1 sheet, 1:250,000
- Rubey, W. W., and Bass, N. W., 1925, The geology of Russell County, Kansas, with special reference to oil and gas resources: Kansas Geological Survey, Bulletin 10, pt. 1, p. 1–86
- Ruddiman, W. F., 1987, Synthesis—the ocean/ice sheet record; *in*, North America and adjacent oceans during the last deglaciation, W. F. Ruddiman and H. E. Wright, Jr., eds.: Geological Society of America, Decade of North American Geology, The Geology of North America, v. K-3, p. 463–478
- Ruhe, R. V., 1956, Geomorphic surfaces and the nature of soils: Soil Science, v. 82, p. 441–455
- _____, 1969, Quaternary landscapes in Iowa: Iowa State University Press, Ames, 255 p.
- _____, 1976, Stratigraphy of midcontinent loess, U.S.A.; *in*, Quaternary Stratigraphy of North America, W. C. Mahaney, ed.: Stroudsburg, Pennsylvania, Dowden, Hutchinson, and Ross, p. 197–211
- _____, 1983, Depositional environment of late Wisconsin loess in the midcontinental United States; *in*, Late Quaternary Environments of the United States—v. 1, The Late Pleistocene, S. C. Porter, ed.: Minneapolis, University of Minnesota Press, p. 130–137
- Ruhe, R. V., and Olson, C. G., 1980, Clay-mineral indicators of glacial and nonglacial sources of Wisconsinan loess in southern Indiana, USA: Geoderma, v. 240, p. 283–297
- Ruhe, R. V., Hall, R. D., and Canepa, A. P., 1974, Sangamon paleosols of southwestern Indiana, USA: Geoderma, v. 12, p. 191–200
- Ruhe, R. V., Miller, G. A., and Vreeken, W. J., 1971, Paleosols, loess sedimentation, and soil stratigraphy; *in*, Paleopedology—Origin, Nature, and Dating of Paleosols, D. H. Yaalon, ed.: Jerusalem, Israel, Hebrew University Press, p. 41–60
- Runyon, H. E., and Rankin, R., 1940, Chemical analyses of some oil-well waters of Russell, Ellis, and Trego counties, Kansas: Kansas Academy of Science, Transactions, v. 43, p. 235–241
- Russell, W. L., 1929, Stratigraphy and structure of the Smoky Hill Chalk in western Kansas: American Association of Petroleum Geologists, Bulletin 13, p. 595–604

- Schaetzl, R. J., 1986, The Sangamon paleosol in Brown County, Kansas: Kansas Academy of Science, Transactions, v. 89, p. 152–161
- Schultz, C. B., and Martin, L. C., 1970, Quaternary mammalian sequence in the central Great Plains; *in*, Pleistocene and Recent Environments of the Central Great Plains, W. Dort, Jr., and J. K. Jones, eds.: Lawrence, University Press of Kansas, p. 341–353
- Schultz, C. B., and Stout, T. M., 1945, Pleistocene loess deposits of Nebraska: American Journal of Science, v. 243, p. 231–244
- _____, 1948, Pleistocene mammals and terraces in the Great Plains: Geological Society of America, Bulletin 59, p. 553–591
- Schultz, C. B., Stout, T. M., and Tanner, L. G., 1957, Medial Pleistocene fossil vertebrate localities in Nebraska: University of Nebraska, State Museum, Bulletin 4, p. 59–81
- Shimek, B., 1909, Aftonian sands and gravels in western Iowa: Geological Society of America, Bulletin 20, p. 399–408
- Siemers, C. T., 1971a, Deltaic deposits of upper part of the Dakota Formation (Upper Cretaceous), central Kansas (abstract): American Association of Petroleum Geologists, Bulletin 55, no. 2, p. 364
- _____, 1971b, Stratigraphy, paleoecology, and environmental analysis of upper part of Dakota Formation (Cretaceous), central Kansas: Ph.D. dissertation, Department of Geology, Indiana University, Bloomington, 287 p.
- _____, 1976, Sedimentology of the Rocktown channel sandstone, upper part of the Dakota Formation (Cretaceous), central Kansas: Journal of Sedimentary Petrology, v. 46, no. 1, p. 97–123
- Simonson, R. W., 1941, Studies of buried soils formed from till in Iowa: Soil Science Society of America, Proceedings, v. 6, p. 373–381
- Sorensen, J. H., 1994, Bibliography of Kansas geology, 1885–1989: Kansas Geological Survey, Bulletin 234, 158 p.
- Sorensen, J. H., Johnsgard, S. J., and Wozencraft, C., 1989, Bibliography of Kansas geology, 1823–1984: Kansas Geological Survey, Bulletin 221, 418 p.
- St. John, O. H., 1883, Sketch of the geology of Kansas: Kansas State Board of Agriculture, Third Biennial Report, p. 571–599
- Steeple, D. W., Knapp, R. W., and McElwee, C. D., 1986, Seismic reflection investigations of sinkholes beneath Interstate Highway 70 in Kansas: Geophysics, v. 51, no. 2, p. 295–301
- Terasmae, J., and Dreimanis, A., 1976, Quaternary stratigraphy of southern Ontario; *in*, Quaternary Stratigraphy of North America, W. C. Mahaney, ed.: Stroudsburg, Pennsylvania, Dowden, Hutchinson, and Ross, p. 51–63
- Thorpe, J., Johnson, W. M., and Reed, E. C., 1951, Some post-Pliocene buried soils of central United States: Journal of Soil Science, v. 2, p. 1–22
- Voorhies, M. R., 1990, Vertebrate biostratigraphy of the Ogallala Group in Nebraska; *in*, Geologic Framework and Regional Hydrology—Upper Cenozoic Blackwater Draw and Ogallala Formations, Great Plains, T. C. Gustavson, ed.: Bureau of Economic Geology, Austin, Texas, p. 115–151
- Ward, P. A., III, 1991, Preliminary correlations of southern High Plains border ashes; *in*, A Prehistory of the Plains Border Region, B. J. Carter and P. A. Ward, III, eds.: Guidebook, South-central Friends of the Pleistocene, Department of Agronomy, Oklahoma State University, p. 65–72
- Ward, P. A., III, Carter, B. J., and Weaver, B., 1993, Volcanic ashes—time markers in soil parent materials of the southern Plains: Soil Science Society of America Journal, v. 57, p. 453–460
- Watson, R. A., and Wright, H. E., Jr., 1980, The end of the Pleistocene—a general critique of chronostratigraphic classification: Boreas, v. 9, p. 153–163
- Wayne, W. J., and Aber, J. S., 1991, High Plains and plains border sections in Nebraska, Kansas, and Oklahoma; *in*, Quaternary Nonglacial Geology—Conterminous U.S., R. B. Morrison, ed.: Geological Society of America, Geology of North America, v. K-2, p. 462–469
- Welch, J. E., and Hale, J. M., 1987, Pleistocene loess in Kansas—status, present problems, and future considerations; *in*, Quaternary Environments of Kansas, W. C. Johnson, ed.: Kansas Geological Survey, Guidebook Series 5, p. 67–84
- Willman, H. B., and Frye, J. C., 1970, Pleistocene stratigraphy of Illinois: Illinois State Geological Survey, Bulletin 94, 204 p.



Kansas Geological Survey
1930 Constant Avenue
The University of Kansas
66047-3726