



Rock Units and Densities (g/cc)		
	Bayfield Group and equivalents	2.30
	Oronto Group and equivalents	2.66
	Keweenaw mafic volcanic and intrusive rocks	2.90; 3.00
	0-10 Km: 2.90; 10-20 Km: 3.00	
	Barron Quartzite	2.30
	Felsic intrusions, undifferentiated	2.75; 2.80
	0-10 Km: 2.75; 10-20 Km: 2.80	
	Mafic intrusions, undifferentiated	2.90; 3.00
	0-10 Km: 2.90; 10-20 Km: 3.00	
	Proterozoic turbidites	2.70

	Unit	Int. Pol. ¹	Incl. Pol. ²	Decl. Pol. ³	Q ⁴	Rem ⁵
A-A'	1	155.25	61.4	30.9	2.39	R
	2	347.43	60.1	-54.6	0.90	N
	3	226.17	47.4	-58.5	0.80	N ⁶
	4	301.77	63.1	-50.3	1.19	N
B-B'	1	228.10	64.9	-45.3	1.48	N
	2	149.34	-61.3	76.5	0.30	R
	3	146.72	55.7	-58.7	0.60	N
C-C'	1	214.15	70.4	-25.9	3.55	N
	2	486.63	70.9	-22.7	4.14	N
	3	486.63	70.9	-22.7	4.14	N
D-D'	1	245.16	58.5	-55.0	0.78	N
	2	146.22	55.5	-58.4	0.59	N
	3	453.60	64.7	-44.7	1.47	N

- Notes: 1: Intensity of total polarization ($\times 10^{-2}$ SI)
 2: Inclination of total polarization (below horizontal).
 3: Declination of total polarization (+ eastward; - westward)
 4: Koenigsberger ratio.
 5: Direction of remanent magnetization
 N - Keweenaw normal; R - Keweenaw reverse.
 6: Remanent magnetization Rotated 20° to the horizontal.