

**KANSAS GEOLOGICAL SURVEY  
OPEN-FILE REPORT 2000-66**

Paleoenvironmental Reconstruction at Fort Riley, Kansas  
1998 Phase  
(05/01/98 - 12/31/98)

by

William C. Johnson

*Disclaimer*

The Kansas Geological Survey does not guarantee this document to be free from errors or inaccuracies and disclaims any responsibility or liability for interpretations based on data used in the production of this document or decisions based thereon. This report is intended to make results of research available at the earliest possible date, but is not intended to constitute final or formal publications.

Kansas Geological Survey  
1930 Constant Avenue  
University of Kansas  
Lawrence, KS 66047-3726

Paleoenvironmental Reconstruction  
at  
Fort Riley, Kansas

1998 Phase  
(05/01/98-12/31/98)

Technical Report

William C. Johnson, Ph.D.  
Department of Geography  
1 Jayhawk Blvd  
University of Kansas  
Lawrence, KS 66045-2121

*submitted to*

United States Army Construction Engineering Research Laboratories (USACERL)  
2902 Newmark Drive  
Champaign, Illinois 61821-1076

*in fulfillment of Contract No. DACA88-98-M-0135, Mod P00004*

February 1999

## TABLE OF CONTENTS

---

<b>INTRODUCTION</b>	3
Research at Fort Riley	3
Tasks of the 1998 Phase	4
<b>BACKGROUND TO THE RESEARCH</b>	4
Regional Late-Quaternary Loess Stratigraphy	5
Pre-Illinoian Stages	5
Illinoian Stage	5
<i>Loveland loess</i>	6
<i>Sangamon soil</i>	6
<i>Fort Riley</i>	7
Wisconsinan Stage	7
<i>Gilman Canyon Formation</i>	7
<i>Peoria loess</i>	9
<i>Fort Riley</i>	10
Holocene Series	10
<i>Brady soil</i>	10
<i>Bignell loess</i>	11
<i>Fort Riley</i>	12
<b>REGIONAL LATE-QUATERNARY ALLUVIAL STRATIGRAPHY</b>	12
Late-Wisconsinan Stage	12
Holocene Series	13
<b>PALEOCLIMATIC HISTORY OF THE WISCONSINAN AND HOLOCENE</b>	15
<b>STUDY LOCALITIES</b>	17
<b>FIELD AND LABORATORY METHODS</b>	17
Mapping Loess Distribution	17
Mapping Alluvial Units	19
Radiocarbon Dating	19
Procedures	19
<b>RESULTS</b>	20
Loess Distribution	20
Loveland and Peoria loesses	20
Bignell loess	20
Distribution and Ages of Alluvial Fills	23
Breakneck Creek	23
Onemile Creek	23
Pumphouse Creek	38
Forsyth Creek	38
Wildcat Creek	46
Locating Buried Archaeological Remains	46
Pumphouse Creek	51
Forsyth Creek	51
Wildcat Creek	56
<b>CONCLUSIONS</b>	64
<b>REFERENCES CITED</b>	67

---

## INTRODUCTION

Many years ago, archaeologists recognized that the preserved record of past cultural activities is evidence of, with varying degrees of accuracy, both the articulation of human groups and their environment and the effects of post-occupational processes on the material remains of those activities (e.g., Schiffer, 1976; Binford, 1981; Wood and Johnson, 1978). The database is not inherited in a pristine condition, however, in that the archaeological record is an imperfect mirror of past cultural activities that formed vulnerable, "fossil" configurations, which are in turn usually flawed by a variety of post-depositional processes, both cultural and natural.

We have to realize that the cultural record is limited by erosional destruction of part of the record. As Ager (1973, p.34) notes, "The sedimentary pile at any one place on the Earth's surface is nothing more than a tiny and fragmentary record of vast periods of Earth history." Bettis and Benn (1984), Gladfelter (1985), Johnson and Logan (1990), and a host of others have made the same observation, but in an archaeological context. At Fort Riley, one has to contend with a record of destructive cut and fill episodes within the valleys and with another of water and wind erosion and downslope displacement of upland loess.

Sampling methods for regional and local archaeological surveys have been articulated by many researchers (e.g., Redman, 1974; Plog, 1976; Warren and O'Brien, 1981). As a function of these traditional approaches, resulting studies generally failed to address directly the pervasive problem of erosion (removal/destruction) and deposition (burial), and how they may skew the visible archaeological record. In recent years, archaeologists and earth scientists have routinely joined ranks to address the issue of geomorphic factors that affect site distributions at all scales.

Geoarchaeological research on Fort Riley represents a high-resolution examination of the alluvial and upland eolian record, with the goal of identifying and mapping the remnants of the sedimentary record that contain the potential for cultural remains. Through the use of radiocarbon dating, alluvial surfaces and fills have been dated and extent of Holocene loess on the uplands approximated.

### Research at Fort Riley

Geoarchaeological research began with an overview study by D.L. Johnson (1992). On-site work by D.L. Johnson, an assistant and the contractor in June, 1993 involved further reconnaissance, subsurface exploration with a vehicular-mounted coring rig, and documentation and sampling of natural exposures. Eighteen localities, numbered 21-38 and from various landscape positions, were studied through sediment and soil cores. These sites represented various landscape positions within Fort Riley. Of the sites that were cored, all except 21, 26, and 33 extended to bedrock or to the residual soil developed on the bedrock; laboratory analyses were conducted on samples from the cores (D.L. Johnson, 1994). Additional study described the distribution of the loess mantle on the upland and refined the stratigraphy at the Sumner Hill locality, i.e., core site 21 (D.L. Johnson, 1996).

The first involvement by this investigator centered on paleoenvironmental research. Specifically, tests of the potential for stable isotope ratio analysis (SIRA) of carbon and opal phytolith analysis were conducted to provide proxy time series of climate for the base (W.C. Johnson

et al., 1994). Study was carried out on a core extracted from the Sumner Hill locality. Isotopic analysis ( $\delta^{13}\text{C}$ ) provided a time series that compares well with a composite of regional carbon isotope data, whereas the opal phytolith record was largely uninterpretable below about 3.5m depth, i.e., for the lower 8.5m of the core. On the basis of a recently derived chronostratigraphy of the site (D.L. Johnson, 1996; W.C. Johnson, 1998), the viable portion of the phytolith record does, however, include the time interval related to cultural occupation, i.e., the last 13,000 years or so.

A second part of the environmental reconstruction was designed to consider the record from several sites distributed over the reservation in upland and valley environments. The research design was constructed as to make coincident use of magnetic analysis, SIRA, and analysis of opal phytoliths. Data from this multiparametric approach were of extremely high quality and produced an extractable environmental record.

The next effort continued investigations on the upland loess sequences, but placed emphasis on the alluvial fills of the Republican and Kansas River valleys and selected tributary valleys. Magnetic, isotopic and phytolith studies were continued on the uplands, in conjunction with close-interval radiocarbon dating of selected loess sites. Studies of the alluvial fill focused on the definition and correlation of stratigraphy, combined with radiocarbon dating.

A last focus addresses representation of the stratigraphic and, to a lesser extent, paleo-environmental data in a geographic information system (GIS) context. Distributions of alluvial surfaces and fills and upland loess deposits are being rendered, and, from these, probabilities of burial for cultural materials from different periods are being generated. The first phase in this last focus was reported earlier (W.C. Johnson, 1998).

### **Tasks of the 1998 Phase**

Major requirements of this contract three tasks, all of which represent a continuation of research conducted in one or more earlier phases of the Fort Riley Geoarchaeological Project. Tasks were to (1) develop spatial data of dated landform sediment assemblages of Fort Riley in ArcView format; (2) collect limited sets of field data that address identified data gaps for completion of the paleoenvironmental model; and (3) complete the laboratory analysis of limited field data and integrate the accumulated data sets into a final technical report, including an ArcView geospatial database of dated landform sediment assemblages. Data from this phase were distilled into a series of maps, which consist of ArcView files graphically rendered with Freehand 8.0.

### **BACKGROUND TO THE RESEARCH**

The distribution, character, and origin of late Pleistocene and Holocene deposits on Fort Riley are, as elsewhere, of direct relevance to realizing and understanding the nature of the prehistoric cultural record. Upland loess and valley fills are the primary late-Quaternary deposits found on the military installation, and these contrasting depositional settings were resource-rich landscapes for prehistoric cultural activities. Because bottomlands offer a particularly rich cultural resource base, the alluvial fills contain a far higher concentration of cultural remains than the upland loess mantle. Considerable research has been conducted on these deposits throughout the region in recent years, with an emphasis on development of chronostratigraphies, but research results from

Fort Riley are making a major contribution to the database and to the understanding of paleoenvironmental conditions for the central Great Plains through the high-resolution approach that has been adopted.

A review of the geomorphology and late-Quaternary stratigraphy and climates are presented, as they relate to the taphonomy of the cultural record. An articulation of the regional loess stratigraphy is necessary for the appreciation of the late-Quaternary record preserved at Fort Riley. The stratigraphic and climatic discussion below is abstracted from previous reports.

### **Regional Late-Quaternary Loess Stratigraphy**

A very good empirical relationship exists between cold stages in the marine oxygen isotope record and documented times of glaciation (ice volume) in the United States (Richmond and Fullerton, 1986b). Given the close correspondence between the glacial record and marine isotope record and the fragmentary nature of the former, it follows that the nearly continuous loessal record should be an excellent terrestrial cognate of the marine isotope record. Therefore, the climatic and chronological record assembled for the marine sequence should match the loessal record well. The generally accepted model relating climate to the loessal record indicates that periods of stability and pedogenesis are usually associated with warm interglacials, and periods of significant loess accumulation coincide with the colder glacial times (Kukla 1977, 1987).

#### **Pre-Illinoian Stages**

Little is known of the pre-Illinoian loesses because far fewer exposures exist than of the Loveland (Illinoian) loess and certainly the Peoria (late-Wisconsinan) loess. Pedogenesis has been recognized in these early loesses, however. Zones of carbonate enrichment, occurring at about 410-360, 330-290, and 250-200 ka, within the Barton County sanitary landfill exposure were interpreted to be pedogenic in origin (Feng et al., 1994). These carbonate zones are likely analogous to the soils observed in the pre-Loveland loesses at the Eustis ash pit, Nebraska (Fredlund et al., 1985) and elsewhere in the region (Schultz and Martin, 1970; Frye and Leonard, 1951). The zones are temporally equivalent to the nonglacial, or warm marine isotope stages (Feng et al., 1994).

No pre-Illinoian Quaternary sediments have yet been recognized on Fort Riley as a result of this series of CERL-funded studies or are reported in the literature. It appears that major entrenchment and denudation preceded the Illinoian because exposures and cores to date have exposed only what is presumed to be Illinoian-age material overlying bedrock. For example, the Illinoian Sangamon soil, loess, and alluvium rest upon a strath, or bedrock-defended terrace at the Sumner Hill locality; conversely, middle Pleistocene loess rests on an erosion surface developed on limestone at the Bala site and sites 27 and 28.

#### **Illinoian Stage**

Stratigraphy associated with this stage within the region consists of the Loveland loess and Sangamon soil. The latter is developed within the upper part of the former; as a consequence, they are often outcropping together.

**Loveland loess.** The Loveland loess is the most widespread pre-Wisconsinan loess in the Midcontinent. Several investigators (e.g., Reed and Dreeszen, 1965; Ruhe, 1969; Willman and Frye, 1970; Ruhe and Olson, 1980) have described it throughout the Missouri, Mississippi and Ohio River basins. Further, it has been recognized south into Mississippi and Arkansas (McCraw and Autin, 1989). The Loveland has been far less studied (e.g., absolute chronology, geometry, mineralogical composition) than the Wisconsinan loesses, namely the Peoria.

The Loveland may be described as a yellowish-brown or reddish-brown eolian silt. Red hues increase toward the top of the formation due to development of the Sangamon soil within the uppermost Loveland. The thickest accumulations occur in the north-central part of the state: recorded thicknesses approach 15 m. A thinning in the loess occurs both southward and westward such that the distribution becomes discontinuous to the southwest. In Kansas, the Loveland is typically less than 10 m thick, but produces a very distinctive mark on the landscape via its variation in stratigraphy. It occurs on uplands and valley side slopes. As a result, the Loveland and its capping Sangamon soil are well expressed in exposures, particularly freshly cultivated fields.

The absolute age of Loveland loess in Kansas is largely uncertain, but recent work at sections exposed in a Geary County quarry in northeastern Kansas, the Barton and Pratt County sanitary landfills of central Kansas, and the Eustis ash pit of southwestern Nebraska provided the first absolute-age information on the Loveland beyond that carried out at the paratype section. Oviatt and others (1988) reported TL ages of 136 ka and 130 ka for the upper part of the presumed Loveland loess exposed in an abandoned quarry near the town of Milford, immediately west of Fort Riley. TL age data from this and others sites in Kansas indicates that the Loveland loess began accumulating sometime before 130 ka (Feng et al., 1994; Johnson and Muhs, 1996). Maat and Johnson (1996) derived a TL age of approximately 160 ka immediately below the Sangamon soil in Loveland loess at the Eustis ash pit.

**Sangamon soil.** This paleosol is strongly developed and occurs throughout the Midcontinent beneath deposits of the Wisconsinan glaciation and within deposits of the Illinoian glaciation or older deposits. The Sangamon soil has been recognized in Indiana (Hall, 1973; Ruhe et al., 1974; Ruhe and Olson, 1980), Illinois (Bushue et al., 1974; Follmer, 1979) where the type section is located (Follmer, 1978), Iowa (Simonson, 1941; Ruhe, 1956, 1969), Nebraska (Schultz and Stout, 1945; Thorpe et al., 1951) and Kansas (Frye and Leonard, 1952). In Kansas, the Sangamon soil is well expressed, occurring throughout the state. Although the soil has received considerable attention in northeastern Kansas (Frye and Leonard, 1949, 1952; Tien, 1968; Caspall, 1970; Bayne et al., 1971; Schaetzl, 1986), it has been recognized at many localities in the state (Bayne and O'Connor, 1968) and recently studied in central Kansas (Feng et al., 1994). Historically, it has been referred to as a "soil in the Sanborn formation" (Hibbard et al., 1944), the Loveland soil (Frye and Fent, 1947), and the Sangamon soil (Frye and Leonard, 1951). The color of the soil ranges from a vivid to pale reddish-brown, with a loss in color occurring westward. Regionally, the soil character varies according to parent material, local drainage and climate which prevailed at the time of pedogenesis. The soil occasionally contains sufficient clay to create a subtle bench on cultivated slopes. Schaetzl (1986) noted that the soil appears to have been a very strongly developed Ultisol or Mollisol.

The Sangamon soil was first used in a time-stratigraphic context to differentiate deposits of the Illinoian and Wisconsinan glacial stages (Leverett, 1899). An appreciable time span for regional

landscape stability and soil formation are suggested by oxidation, apparent deep leaching, and high clay accumulation. A major problem associated with the Sangamon soil is its diachronous upper and lower boundaries (Follmer, 1978, 1982, 1983). To further confuse the time element, the lower 1-2 m (3.3-6.6 ft) of the early Wisconsinan loess is typically weathered and forms a pedological continuum with the underlying Sangamon soil (Follmer, 1983), and early investigators mistakenly included the former in the Sangamon profile. The Sangamon should be considered a *pedocomplex* rather than a single soil which developed under a unique environmental condition (Schultz and Tanner, 1957; Fredlund et al., 1985; Morrison, 1987). It apparently represents two or more paleosols welded together to form a complex that reflects significant spatial and temporal variation in environmental conditions and an appreciable time span. Although laboratory data from exposures in central Kansas indicate the Sangamon soil was strongly weathered chemically, presumably under a warm, moist climate (Feng, 1991), recent data from the Eustis ash pit in south-central Nebraska indicate that the Sangamon soil is not very mature geochemically (Muhs and Johnson, 1996).

Because of apparent time transgressiveness, the age of the Sangamon soil is not precisely known. Follmer (1983) reported a radiocarbon age of  $41,700 \pm 1100$  yr B.P. on plant material from the top of the Sangamon in its type area in Illinois. Forman (1990a) reported TL ages of  $140 \pm 20$  and  $70 \pm 10$  ka from loess below the Sangamon soil at two separate sites in Iowa and Illinois, and concluded the Sangamon soil is diachronous and may consist of multiple soils. Feng (1991) and Feng and others (1994) reported a TL age of about 70 ka in the lowermost part of the Sangamon soil exposed in central Kansas and associate it with marine isotope stage 3. Although Richmond and Fullerton (1986b) assigned Sangamon time to 132-122 ka (isotope substage 5e), they acknowledged reported ages (relative and absolute) ranging from early Illinoian to middle Wisconsin. Basal ages on the overlying Gilman Canyon Formation from numerous locations in Kansas and Nebraska (Johnson, 1993a, b) provide a minimum age of about 45 ka for the Sangamon soil. Also, Forman (1990b) and Forman and others (1992) obtained TL and radiocarbon ages of 35-30 ka within the loess overlying the Sangamon soil at the Loveland paratype section in Iowa and the Pleasant Grove School section in Illinois.

**Fort Riley.** Thus far, investigations on the reservation have indicated that Illinoian age sediments are ubiquitous but relatively thin, and expressed primarily as the Sangamon soil developed within them. Best known expression of the Sangamon soil occurred at the Sumner Hill locality in a thin layer of Loveland loess overlying alluvium.

### **Wisconsinan Stage**

Stratigraphy associated with the Wisconsinan glacial period in the central Great Plains consists of two loess units and associated soils: Gilman Canyon Formation and Peoria loess.

**Gilman Canyon Formation.** The Gilman Canyon Formation, first recognized in Nebraska (Reed and Dreeszen, 1965), is a middle to early-late Wisconsinan (cf. Farndalian) loess. Equivalents of the formation have been recognized elsewhere: the Loveland loess is buried by the Roxana silt from Minnesota and Wisconsin to Arkansas and by the Pisgah Formation in western Iowa (Bettis, 1990). The Gilman Canyon of Nebraska and Kansas is typically dark in color, silty, leached of calcium

carbonate, and heavily enriched in organic carbon via pedogenesis (melanization). As noted above, the formation was once considered to be the attenuated A horizon of the Sangamon soil (Thorpe et al., 1951; Reed and Dreeszen, 1965).

Reed and Dreeszen (1965) provide limited textural data and description of the Gilman Canyon Formation at the type section. Their description within the columnar section at the Buzzard's Roost exposures states (p. 62): "Upper 12 inches [31 cm] is medium dark gray, slightly humic, silt; middle 1 foot 1 inch [33 cm] is light brownish-gray silt; basal 3 feet 8 inches [1.12 m] is dark brownish-gray, humic, soil-like silt; entire thickness is noncalcareous . . . 5 feet 9 inches [1.75 m]." Although all of these attributes described at the type section appear representative of the formation as observed in Nebraska and Kansas, the bimodal distribution of humus is curious: this suggests the existence of two periods of relative stability, or low accumulation rates, and an intervening period of accelerated accumulation rates. Consequently, the Gilman Canyon Formation often appears as one or more cumulic A horizons that are developed within a variably to noncalcareous loess, usually no more than 1.2 m thick. In a section revealing an expanded valley phase of the Gilman Canyon Formation, May and Souders (1988) recognized three distinct organic zones, each of which may represent a separate episode of pedogenesis. Two such zones have been recently observed by the author at the Eustis ash pit in south-central Nebraska. If two or more distinct periods of soil formation did indeed occur regionally, they are obscured at many localities, likely due to bioturbation. Overall, the formation reflects a sufficiently slow rate of loess fall (<0.08mm/yr) such that pedogenesis was operating more or less continuously, but with a decreased intensity at one or more times.

As expressed, the Gilman Canyon Formation is frequently overlain by 0.9-1.5 m of leached loess which is considered to be basal Peoria Formation. Correlative with the Gilman Canyon and overlying leached loess zone is the *Citellus* zone (a ground squirrel now recognized as the genus *Spermophilus*) of Nebraska (Condra and Reed, 1950). The leached zone is transitional between the well developed A horizon(s) in the Gilman Canyon and the calcareous Peoria loess above, and probably reflects a sufficiently slow accumulation of Peoria loess such that pedogenesis could keep pace only partially. A.B. Leonard (1951, 1952) supported the contention that the leached, or basal zone was slowly accumulating, early Peoria loess experiencing pedogenesis through inference that gastropods were originally present, but subsequently destroyed during weathering of the loess. Above the leached zone, the rate of accumulation of Peoria loess was sufficiently rapid (c. 0.6mm/yr) as to preclude any soil development.

Radiocarbon ages from the Gilman Canyon Formation range from approximately 40 ka at the base to 20 ka at the top (May and Souders, 1988; Johnson et al., 1993a). The basal age of 40 ka agrees well with the time set by Richmond and Fullerton (1986a) for the beginning of the late Wisconsin. While Nebraska has many dated locations forming an arcuate pattern around the eastern and southern sides of the Sand Hills, data from Kansas are relatively limited. The ages in Kansas do show, however, good agreement from the south-central to the north-central part of the state and with those from Nebraska.

Given the radiocarbon time control and stratigraphic information currently available for the Gilman Canyon Formation within Kansas and Nebraska, it appears that the associated soil is a *geosol*, i.e. a laterally traceable, mappable, pedostratigraphic unit with a consistent time-stratigraphic position (Morrison, 1965; North American Commission on Stratigraphic Nomenclature, 1983, p.

865). The entire formation may be considered a geosol, but, because of the possibility for the existence of two or more identifiable cumulic A horizons merged or welded together, it may ultimately be considered a *composite geosol*. On Fort Riley, the Gilman Canyon Formation is typically welded to the Sangamon soil below and to the Brady soil above.

**Peoria loess.** Leverett (1899) first proposed the name Peoria for an interglacial period between the Iowan and Wisconsin glacial stages. When Alden and Leighton (1917) demonstrated the Peoria was younger than Iowan, usage shifted to that of a loess, rather than a weathering interval. Within the Midcontinent, several names have been used for post-Farmdalian loess. Ruhe (1983) prefers to use the term "late Wisconsin loess" because of the uncertainties in stratigraphic equivalency from one region to another. The Peoria Formation is typically an eolian, calcareous, massive, light yellowish-brown silt that typically overlies the Loveland Formation or an approximate equivalent of the Gilman Canyon Formation.

Ruhe (1983) notes three major features of late-Wisconsinan (Peoria) loess: it thins downwind from the source area, decreases in particle size systematically away from the source area, and is strongly time-transgressive at its base. The latter feature is unresolved and results in correlation problems. Ruhe (1969) realized a decrease in the age of the soil under the loess from 24,500 yr BP near the Missouri River to about 19,000 yr BP eastward across southwestern Iowa. A decrease from 25,000 to 21,000 yr BP was noted for the base of the loess along a transect in Illinois (Kleiss and Fehrenbacher, 1973). The top of the loess also seems to be time-transgressive, ranging from about 12,500 yr BP in Illinois (McKay, 1979b) to 14,000 yr B.P. in central Iowa (Ruhe, 1969).

In Kansas, the Peoria is a reddish, yellowish, or tan buff color, homogeneous, massive, locally fossiliferous, variably calcareous, and ranges from coarse silt and very fine sand to medium to fine silt and clay (Frye and Leonard, 1952). Thicknesses vary from in excess of 30 m adjacent to the Missouri River valley to 0.6 m in discontinuous patches. Any accumulation less than 0.6 m is presumed unrecognizable in the field because it has become incorporated into the existing surface soil. Peoria loess typically rests conformably upon the Gilman Canyon Formation.

Despite the amount of attention given Peoria loess in Kansas, the source of the silt is not completely certain. Upon a review of the available data, Welch and Hale (1987) conclude that a single source was not likely for all loess deposits in Kansas, and that the loess was derived from a combination of three sources: glacial outwash river flood plains, present sand dune areas, and fluvial and eolian erosion of the Ogallala Formation. Research on trace element concentrations in loess (Johnson and Muhs, 1996) indicates, however, that the Platte River valley was the primary source, with secondary inputs from the major river valleys to the south (e.g., Republican, Smoky Hill, Solomon, Arkansas).

Although readily visible stratigraphic breaks such as the Jules soil recognized in Illinois (Frye and Willman, 1973; Frye et al., 1974; Ruhe, 1976; McKay, 1979a, b) and the soil zones in Iowa (Daniels et al., 1960; Ruhe et al., 1971) have not yet been identified in Kansas and adjacent Nebraska, evidence of one or more stable or vegetated surfaces is common. The only indication of soil development recognized is that of a Bt horizon in the Medicine Creek valley (May and Holen, 1993); interestingly, the soil has a probable Paleoindian association (May, 1991). The most common line of evidence for a discontinuity(ies) in Peoria loess deposition is that of plant remains, usually outcropping as lenses. Many of the age determinations were made from *Picea* remains, indicating

a cool, moist environment. Although radiocarbon data document the burial of vegetative material throughout the Woodfordian, two temporal clusters or modes of ages appear from the limited data: one 18-17 ka and another 14-13 ka. The former time interval represents the last glacial maximum and the latter the time of major deglaciation (Ruddiman, 1987). Interpreting ice core data from Greenland, Paterson and Hammer (1987) record a dramatic decrease in atmospheric dust content from about 13,000; this period of reduced atmospheric dust may relate to the time of relative surface stability and tree establishment. Regional geomorphic data also support the existence of a hiatus at this time. May (1989), identifies deposition of the Todd Valley Formation in the South Loup River of central Nebraska at about 14 ka, which is subsequently buried by loess. Further, Martin (1990) identifies entrenchment in the Republican River of south-central Nebraska at about 13 ka, after which valleys were filled with late Peoria Loess.

**Fort Riley.** The Gilman Canyon Formation sediments and soil were present at all sites sampled for magnetic and isotopic analyses, indicating that the formation has both upland (loess) and valley (alluvium) phases preserved at Fort Riley, as elsewhere in the region. Radiocarbon ages on the formation range from about 19ka to over 23ka. These relatively young ages represent the most recent period of Gilman Canyon time pedogenesis; the relatively unaltered loess and alluvium was not sampled for dating. Coring and backhoe trenching indicates that Peoria loess is typically relatively thin, except along the bluffs of the Kansas River valley, where it reaches thicknesses of ten or more meters.

### **Holocene Series**

The transition from the late Pleistocene into the Holocene, about 10 ka (Hopkins, 1975), was a time of dramatic environmental change and attendant stratigraphic discontinuities. A major feature in the loessal stratigraphic record is a well developed soil, the Brady. Approximately 9,000 years ago, the uplands again destabilized with the renewed influx of loess, which continued episodically throughout the Holocene. Holocene loess is termed the Bignell throughout the central Great Plains.

**Brady soil.** The Brady soil was first named and described by Schultz and Stout (1948) at the Bignell Hill type locality, an eolian sequence exposed along a road cut in the south valley wall of the Platte River of western Nebraska. The soil is developed within the Peoria loess and is overlain by the Bignell loess. The name was subsequently adopted by researchers in Kansas (Frye and Fent, 1947; Frye and Leonard, 1949, 1951). It is regionally extensive only in the northwestern and west central parts of Kansas, and even there it occurs discontinuously on the landscape. Frye and Leonard (1951) and Caspall (1970, 1972) recognized Brady development in northeastern and other parts of Kansas. Without the overlying Bignell loess, the Brady soil does not exist; the modern surface soil has incorporated post-Bradyan loess fall into its profile or may have developed in Peoria loess subsequent to the erosion of the Brady soil and Bignell loess. The Brady soil is typically dark gray to gray-brown and better developed than the overlying surface soil within the Bignell loess. Strong textural B horizon development and carbonate accumulation in the C horizon are typical, although it occasionally displays evidence of having formed under poorer drainage conditions than have associated surface soils (Frye and Leonard, 1951). Feng (1991) noted that the Brady soil, as

expressed in Barton County, is strongly weathered both physically and chemically.

The Brady is the first major geosol to occur in the stratigraphic record of the region since the Gilman Canyon soil 10,000 years earlier. Still, its age had been uncertain, even at the type locality, until recently. Dreeszen (1970, p.19) reported a radiocarbon age of  $9160 \pm 250$  (W-234) obtained in 1954 and another in 1965 of  $9750 \pm 300$  (W-1676), both from the type section but very likely contaminated by modern plant roots. Subsequently, Lutenecker (1985) reported an age of  $8080 \pm 180$  yr BP but provided no specifics other than that the source was the A horizon of the Brady soil at the type section. Better age control for the type section has since been secured by this investigator: radiocarbon ages of  $9,240 \pm 110$  (Tx-7425) and  $10,670 \pm 130$  (Tx-7358) yr BP were obtained on the upper and lower 5 cm, respectively, of the Brady A horizon.

The Brady soil has been recently dated at localities in Nebraska and Kansas. Souders and Kuzila (1990) obtained a radiocarbon age of  $10,130 \pm 140$  yr BP on the Brady soil occurring within the Republican River valley of south-central Nebraska. Sites along Harlan County Lake upstream from Naponee have yielded a number of ages, ranging from  $10,550 \pm 160$  to  $9,020 \pm 95$  yr BP, on exposures of the Brady soil (Cornwell, 1987; Johnson, 1989; Martin, 1990; Martin and Johnson, 1995). Two radiocarbon ages of  $9820 \pm 110$  (TX-7045) and  $10,550 \pm 150$  (TX-7046) yr BP have been derived from the upper and lower 5 cm, respectively, of the Brady A horizon exposed in Barton County, central Kansas (Feng, 1991).

Although it appears Brady pedogenesis occurred from about 10,500 to as recently as 8,500 yr BP, greater refinement of the Brady soil chronology is necessary, but present data clearly indicate it was a product of a major period of landscape stability at a time when widespread climatic shifts were occurring at the end of the Wisconsin. This was the first significant period of soil development since Gilman Canyon time, and represents the climate of the early Holocene. There is an isochronous alluvial soil found throughout the region which is particularly well expressed within the Kansas River basin (Johnson and Martin, 1987; Johnson and Logan, 1990). The two ages of  $8,274 \pm 500$  (C-108a) and  $9,880 \pm 670$  (C-471) yr BP determined from alluvial fill (Fill 2A) at archaeological sites Ft-50 and Ft-41 on Harry Strunk Lake in southwestern Nebraska (Schultz et al., 1951; Libby, 1955) were the first radiocarbon determinations on the Brady soil. The soil, occurring in both eolian and alluvial contexts, appears to qualify, based upon present radiocarbon data, as a *geosol*, like the Gilman Canyon Formation soil. Collectively, age data indicate a soil forming interval that lasted for 1,500 to 2,000 years.

Development of the Brady soil correlates well with indicators of regional climatic change. The fossil pollen record at Muscotah Marsh of northeastern Kansas indicates that spruce had essentially disappeared from the region by about 10,500 yr BP. As this decline occurred, deciduous tree species increased until about 9,000 years B.P., the time at which grassland expansion began (Grüger, 1973). On a hemispheric scale, the abrupt decrease in atmospheric dust noted in the Greenland ice core at 10,750 yr BP (Paterson and Hammer, 1987) reflects decreased loess deposition and possibly Brady-age pedogenesis associated with relative terrestrial stability. Further,  $^{18}\text{O}$  levels within the same core suggest rapid warming about 10,750 yr BP, with the characteristic Holocene temperature regime being established about 9,000 yr BP.

***Bignell loess.*** The Bignell loess was first described and named at the type locality in a bluff exposure on the south side of the Platte River valley southeast of North Platte, Nebraska (Schultz and Stout,

1945). It is typically a gray or yellow-tan, massive silt, calcareous and seldom more than 1.5 m (5 ft) thick. Although it is often somewhat less compact and more friable than the underlying Peoria loess, no certain identification can be made without the presence of the Brady soil. The Bignell loess does not form a continuous mantle on the Peoria; instead, it occurs as discontinuous deposits which are most prevalent and thickest adjacent to modern-day valleys, particularly the south side, and often within depressions on the Peoria surface. Feng (1991) speculates that the Bignell loess of central Kansas is relatively well weathered because it was derived from a preweathered source, the Brady soil surface, perhaps eolian and alluvial phases alike. This is consistent with the earlier interpretation derived in Nebraska that Bignell loess is at least partially comprised of re-worked Peoria loess (Condra et al., 1947, p. 33).

It appears from the radiocarbon ages obtained at the type section in Nebraska and the Speed road cut in northwestern Kansas that the Bignell loess can be no older than 8,000 to 9,000 yr BP. Snails collected by A.B. Leonard from the lower part of the Bignell in Doniphan County, northeastern Kansas, produced ages of  $12,500 \pm 400$  (W-231) and  $12,700 \pm 300$  (W-233) yr BP (Frye and Leonard, 1965). Because the shell material had absorbed an indeterminate amount of dead carbonate, Frye and others (1968) proposed an averaged age of approximately 11,000 years. Based upon the age data available for the Brady, the soil humate-derived ages are probably closest to reality.

**Fort Riley.** The Brady soil and Bignell loess are prominent elements of the late-Quaternary stratigraphy on the reservation. Together with the surface soil, they typically comprise the approximately upper 2m of most sites documented (D.L. Johnson, 1996), including the loess sites references in this report. Therefore, it appears that appreciable loess was deposited during the Holocene. This is not surprising given the proximal juncture of two large stream systems, which formed a significant loess source (dust from a relatively wide valley floor), and the situation of the sites on the north side of the river valley with prevailing southerly winds.

## **Regional Late-Quaternary Alluvial Stratigraphy**

### **Late-Wisconsinan Stage**

Much of the chronology of late-Wisconsinan landform evolution for the region was compiled in the 1940s and early 1950s, prior to the use of radiocarbon dating (e.g., Lugin, 1935; Schultz and Stout, 1945; Frye and Leonard, 1951), and focused to a large degree on the upland rather than valley deposits. Additionally, erosion has removed a large part of late-Quaternary record from most drainage basins in the central Great Plains (Knox, 1983). Accordingly, a comprehensive sedimentation and erosion chronology for the region during the late Wisconsin is lacking.

It is becoming increasingly apparent that entrenchment occurred in the channels of the Kansas River basin sometime during the late Wisconsin. A basal soil buried within the fill of both tributary and major stream valleys of the Kansas River basin has an age of 10,500-10,000 yr BP (Johnson and Martin, 1987; Johnson and Logan, 1990), thereby providing a minimum age on the entrenchment. May (1989) has radiocarbon dated the Todd valley fluvial sand of central Nebraska to about 14,000 years B.P., although Condra et al. (1950) had postulated a much earlier Wisconsinan

age. Martin (1990; 1993) recognized a late-Wisconsin fill in the Republican River valley that was largely removed through entrenchment about 13,000 yr BP. A radiocarbon age of 14,700 yr BP was obtained on spruce wood situated above cross bedded fluvial sand and gravel at the North Cove site located in that same reach of the Republican River valley (Wells and Stewart, 1987; Johnson, 1989). At the Prairie Dog Bay site in the Republican River valley, the stratigraphy and radiocarbon ages suggest down cutting before 11,800 yr BP (Martin, 1990, 1993). Speculation about the cause of this entrenchment centers on an increase in effective moisture as climatic conditions ameliorated towards the end of the Pleistocene. Spring deposits dating to this time at the North Cove site possibly formed during the increase in moisture (Johnson, 1989; Martin, 1990).

Brice (1964) studied alluvial fills and terraces in the valleys of the North Loup, Middle Loup, and South Loup rivers of central Nebraska, and identified two major terraces in the Loup valleys. The Kilgore terrace occurs as remnants 26 to 30 m above stream level along the South Loup River. Brice suggested that valley fill underlying the Kilgore terrace is Peorian (late Wisconsin) in age. The adjacent Elba Terrace, which stands 11 to 12 m above stream level, is the most prominent and extensive terrace in the main valleys and is primarily of Holocene age.

### **Holocene Series**

During the last decade, a great deal of attention has been focused on the development of alluvial chronologies in the central Great Plains, typically in connection with geoarchaeological investigations. As a consequence, this research has resulted in a number of studies and a sizable radiocarbon data base; well over 400 radiocarbon ages have been obtained from alluvium in Kansas and Nebraska (Johnson et al., 1996). Only a sampling of the many studies is presented below.

Much of the research in Nebraska has focused on the Loup River basin. Brice (1964) recognized two major terrace systems in the basin and obtained early Holocene radiocarbon ages of 10,500, 9,000, and 8,500 yr BP on fill beneath the lower of these terraces, the Elba. In a recent re-examination of the Elba terrace, May (1990, 1991) secured radiocarbon ages ranging from nearly 11,000 to 4,670 yr BP from the Cooper's Canyon area. On the South Loup River, May (1986, 1989, 1992) recognized four alluvial fills, with the oldest one dating between about 10,200 and 4,700 years B.P., thereby correlating temporally with the Elba terrace of the North Loup. Elsewhere in the basin, Ahlbrandt and others (1983) dated organic accumulations in alluvial sands at 8,410 yr BP from a site on the Dismal River.

In the Kansas River basin, alluvial geomorphic studies have a relatively long history, beginning in the 1950s. The first dating of alluvial stratigraphy on the Kansas River proper was done by Holien (1982), who obtained an age of 10,450 yr BP on a soil buried within lower Newman terrace fill at the Bonner Spring site. Subsequent radiocarbon dating of Newman fill at this locality (Johnson and Martin, 1987; Johnson and Logan, 1990) and others (Bowman, 1985) produced more early Holocene ages. The lower Holliday terrace has dated about 4,300 yr BP and younger (Johnson and Logan, 1990).

Many studies have been conducted elsewhere in the Kansas River basin on the many tributaries. Some of the first radiocarbon dating was carried out on samples collected from the Republican River basin by Schultz and others (1948): from varied locations they secured early to middle Holocene ages from buried soils. The most recent research in the basin was conducted by

Martin (1990, 1992) who concluded from dating various alluvial fills that the majority of the fill was deposited less than about 4,600 yr BP.

Several geoarchaeological studies were done in conjunction with cultural resource management projects focusing on federal impoundments within the Kansas River basin. Mandel (1987) in a study of the lower Wakarusa River recognized two terraces, the lower of which produced radiocarbon ages of about 2,900 yr BP and less. A study of the alluvial history of the Smoky Hill River in the vicinity of Kanopolis Lake (Mandel, 1988, 1992) revealed a striking absence of early and middle Holocene fill in small valleys, and middle Holocene fill in the main valleys and in alluvial fans.

In their study of Wolf Creek basin, Kansas, Arbogast and Johnson (1994) observed that alluviation of early-Holocene flood plains in this small basin was episodic, with at least one period of flood plain stability and soil formation about 6,800 yr BP. During the middle Holocene (ca. 6,500-5,300 yr BP), lateral erosion and entrenchment flushed most early-Holocene fill from the main valley of Wolf Creek and the lower reaches of its larger tributaries. Following the interval of mid-Holocene erosion, sediment accumulated on flood plains between 5,300 and 3,000 yr BP. Late Holocene alluviation was episodic, with intervening periods of flood plain stability and soil formation about 1,800, 1,500, and 1,200 yr BP.

A number of studies have been conducted in the Arkansas River basin area of south-central and southeastern Kansas. Mandel examined terraces and associated fills in the Neosho (Mandel, 1989, 1992, 1993) and Verdigris Rivers (Mandel, 1993), obtaining radiocarbon ages on fill to about 4,200 yr BP. The most extensive study in the Arkansas River basin was that of the Pawnee River basin by Mandel (1988, 1991, 1994). Two terraces were recognized in the higher order tributaries, with fill of the high terrace dating between about 10,000 and 5,000 yr BP, and that of the low terrace to 3,000 yr BP and younger. Of the three terraces present in the lower part of the system, the lowermost one has Holocene fill and the others are Pleistocene. Holocene valley fills in the Pawnee Basin appear to lack soil development from about 7,000 to 5,000 yr BP.

The alluvial record is temporally and spatially fragmented, i.e., the history of valley and stream evolution stored in alluvium is scattered and wrought with gaps. So, it is only by assembling this fragmentary information that one obtains a unified perspective on the record of stream evolution. Out of the many studies conducted in recent years, a pattern of change is emerging. Large stream valleys appear to contain, more or less, alluvial fill dating throughout the Holocene, whereas small stream valleys typically contain only fill dating in the late Holocene. This model has an intuitive basis in that the probability of survival of early and middle Holocene fill in smaller streams is greatly diminished by the limited storage capacity for alluvium and the relatively high stream gradients, large area in hillslope, and associated peaked flood waves. Exceptions to this pattern do, of course, exist (e.g., Lime Creek, Nebraska: May, 1996; Wolf Creek: Arbogast and Johnson, 1994), but are likely due to locally unusual valley width and other discernable factors.

A first approximation of this alluvial model was presented by Johnson and Martin (1987) in an examination of radiocarbon ages obtained from alluvial fill in the central Great Plains. In recent years, the model has evolved with a vastly expanded data base and been articulated recently by Mandel (1995). He noted that fill in small valleys appears to be less than 4,000 years old, and that the missing early and middle-Holocene record is frequently preserved at the lower end of small stream valleys as terrace fill or alluvial fans.

From the alluvial chronologies, it is obvious that regional synchronicity of stream behavior exists in the central Great Plains (Johnson and Martin, 1987; Johnson and Logan, 1990; Mandel, 1995). When erosion and sedimentation are considered in a stream hierarchical sense, patterns of coincidence appear, such as similar times of flood-plain stability and attendant soil formation. A frequency distribution of over 400 radiocarbon ages from alluvium of Kansas and Nebraska provides an indication of the synchronicity. The high frequencies of the last 5,000 years reflect the age of the alluvium in large and small streams, whereas those prior to about 8,000 represent the ages from the large valleys alone. Alluvial fans ages account for many ages within the 4,000 to 8,000 year range (Mandel, 1995). The greatest frequency of ages occurs about 1,200 yr BP, a time when pronounced low terrace stability and soil development occurred throughout the stream systems. Another notable feature of the distribution is that when the ages obtained from alluvial fans are not considered, very few alluvial ages fall within the 5,000 to 7000 yr BP period. This paucity of ages suggests little flood-plain stability and/or preservation of alluvium from that interval, which coincides with the Altithermal climatic episode. Stream activity of this dry period may have been characterized by rapid sedimentation, thereby precluding soil development, in response to low-frequency, high-intensity convectional storms (Knox, 1976, 1983).

Regional synchrony in Holocene fluvial behavior suggests that climatic fluctuation is the dominant external variable in stream systems (Wendland, 1982; Knox, 1976, 1983). Changes in climate during the Holocene were frequent and episodic (e.g., Wendland and Bryson, 1974; Kutzbach, 1985; COHMAP members, 1988), resulting in discrete periods of stream stability and instability (Knox, 1983).

The concept of a middle Holocene, or Altithermal (ca. 7,000-5,000 yr BP) cultural hiatus on the Great Plains has become well-entrenched within the archaeological literatures. Of the various theories put forth to explain the hiatus (Reeves, 1973), fluvial erosion or aggradation sufficient to dramatically alter the record for the region during the interval 7,000-5,000 yr BP is most pertinent (Johnson, 1987; Mandel, 1995). Some argued that the similarity in the alluvial stratigraphic record from eastern humid portions of the region to the more arid western areas, as well as with chronologies further afield indicates that regionally anomalous erosion and deposition do not explain the hiatus completely; rather, the increased dryness during the Altithermal was likely sufficient to reduce populations on the Plains (Wedel, 1961; Knox, 1978; Wendland, 1978). However, the rapidly expanding alluvial radiocarbon and stratigraphic data base for the region is indicating that much of the cultural record, namely that of the Archaic period, is buried, often deeply, or lost to erosion.

### **Paleoclimatic History of the Wisconsin and Holocene**

As the most recent glacial episode, the Wisconsin has the greatest chronostratigraphic resolution. However, existing knowledge of the climatic environment for this time interval is relatively limited and inconsistent for the central Great Plains. To date, either pollen records from peripheral areas or synthesis of various types of proxy data have been used to reconstruct climate history of the region.

The insolation record exhibits two relatively warm peaks (c. 50 and 30 ka) during marine isotope stage 3. Each peak is followed by a relatively minor and gradual decrease, culminating in the decrease to the glacial maximum (c. 18ka). In contrast to the gradual insolation changes, most

paleoclimatic records from this interval indicate rapid alternations of warm and cold events, which in frequency and timing appear to be unrelated to the Milankovitch forcing (Curry and Follmer, 1992). Dansgaard and others (1985) showed that rapid and extreme fluctuations in stable isotopes (signifying air-temperature differences of 5 °C) from two sites in Greenland appear to correlate over the past 50,000 years. These fluctuations are in phase with changes in CO<sub>2</sub> and dust content. Rapid and substantial temperature fluctuations recorded in ice core segments during stage 3 also correspond with oscillations in the North Atlantic marine sediment record of species abundance and ice rafting (Heinrich, 1988). Using accelerator <sup>14</sup>C ages on a planktonic polar species, Broecker and others (1988) identified four rapid climatic oscillation between 40 and 22 ka in the North Atlantic.

The regional upland vegetation, as inferred from the pollen record from Cheyenne Bottom in Kansas (Fredlund, 1995), appears to have been nearly treeless throughout the Farmdalian (ca. 30-24ka). The pollen record suggests that, although regional tree and shrub populations were higher and more diverse than in Holocene, they were a secondary component of the overall vegetative structure. Grassland-sage steppe dominated the regional uplands surrounding the Cheyenne Bottoms basin throughout the Farmdalian period. The rise in Cheno-Am pollen percentages and an influx of sand beginning at about 25 ka probably mark the rapid onset of a cycle of aridity. Immediately after the onset of aridity, the most noticeable changes are declining percentages in the Cheno-Am types and rising percentages of *Pinus*. The increase in *Picea* and other arboreal pollen may signal a climatic shift toward cooler climatic conditions at ca. 24 ka. Unfortunately, the Woodfordian substage of the Wisconsin is missing from the Cheyenne Bottoms record.

Limiting radiocarbon dates in North America indicate that glaciation commenced about 25-27 ka and thus allow less than 10,000 years for ice buildup prior to 18 ka (Andrews, 1987). The structure of deglaciation is uncertain. There is evidence supporting: (1) a smooth deglaciation model with fastest ice wastage centered on 11 ka; (2) a two-step deglaciation model with rapid ice wasting from 14 to 12 ka and 10 to 7 ka, and a mid-deglacial pause with little or no ice disintegration from 12 to 10 ka; and (3) a Younger Dryas deglaciation model with two rapid deglacial steps as in (2) above, interrupted by a mid-deglacial reversal with significant ice growth from 11 to 10 ka.

The data supporting the smooth deglaciation model are maps of Laurentide ice area based on radiocarbon-dated glacial deposits (Andrews, 1987). Although there are subtle suggestions of more rapid retreat at or near the time of the two steps mentioned above, these curves indicate a steady progressive retreat of North America ice, with significant oscillations in retreat rate only at local spatial scales. Some marine δ<sup>18</sup>O curves also show a smooth progressive decrease toward Holocene values.

The step deglaciation model is also supported by some marine δ<sup>18</sup>O records (Mix, 1987). In addition, the distinctive patterns of change in sea-surface temperature of the North Atlantic Ocean and in Greenland ice-core δ<sup>18</sup>O values also show abrupt step-like warmings at 10 ka and about 13 ka; these warmings might be associated with step-like decreases in Laurentide ice volume. Regionally integrated rates of pollen change in eastern and central North America also show a rapid change in centered on 13.7 and 12.3 ka. (Ruddimann, 1987).

The Younger Dryas (e.g., Osborn et al., 1995; Bard et al., 1993; Peteet et al., 1992) deglaciation model is suggested by the strong signal of sea-surface temperature cooling between 11 and 10 ka in the North Atlantic Ocean. At least early and perhaps all of Brady pedogenesis coincides with an abrupt and brief cool interval correlative with the classic Younger Dryas cold interval of the

North Atlantic region.

By the middle Holocene, drying had reached a maximum according to most studies. Northwest Texas was experiencing conditions of maximum temperatures, minimum precipitation, and eolian activity between 6000 and 4500 years B.P. (Holliday et al., 1983; Holliday, 1985; 1989; Johnson, 1987; Pierce, 1987). This episode coincides with  $\delta^{13}\text{C}$  values from soil organic matter from the same area revealing a shift from -23‰ in the early Holocene to -15‰ in the middle Holocene (Haas et al., 1986). These results were interpreted to represent a shift from cool-season  $\text{C}_3$  grasses to warm-season  $\text{C}_4$  grasses. Based on enriched  $\delta^{13}\text{C}$  values in soil carbonate from northwest Texas, Humphrey and Ferring (1994) also show a middle Holocene xeric episode, although the  $\delta^{18}\text{O}$  values from these same carbonates do not indicate a significant temperature change.

A noticeable shift back to cooler and/or wetter conditions was detected in many areas shortly after 5000 yr BP. The Great Bend Sand Prairie transformed to conditions much like present (Arbogast, 1995). According to Humphrey and Ferring (1994), the return to mesic conditions after 5000 yr BP was interrupted in north-central Texas by a brief warming and drying episode between 2000 and 1000 yr BP. Based on depositional environments, they concluded that cooler and wetter conditions returned after 1000 yr BP.

## **STUDY LOCALITIES**

During this phase, stratigraphic investigations focused on tributary alluvial fills and further definition of the Brady soil and Holocene loess accumulations on the uplands. Tributary valleys studied included those of Onemile, Breakneck, Pumphouse, Forsyth, Threemile, Seven Mile, and Wildcat Creeks. Only the lowermost parts of Threemile and Sevenmile were studied, as most of these systems is inaccessible due to the extent of the firing range. Investigations of the upland loess mantle consisted of further defining the chronology and distribution of the Bignell loess.



## **FIELD AND LABORATORY METHODS**

### **Mapping Loess Distribution**

Due to the size and restricted access to much of the installation, loess distribution was initially mapped using the Natural Resources and Conservation Service (NRCS) electronic soils maps, available from Data Access and Support Center (DASC), Kansas Geological Survey. Unfortunately, these data for Fort Riley are of average to poor quality: in an attempt to combine different soil phases into a single soil series, the developers of the electronic database have often crossed series and grouped unrelated soils into associations. As a result, the soils map exhibits some unusual and unnatural patterns (Fig. 1). General patterns of loess extent do emerge, but scrutiny of the existing file indicates that loess-derived soils are significantly under represented. Because of the unreliability of the electronic data, the original paper copies of the Riley and Geary county soil surveys were used during the field work on the uplands. Distribution of the Bignell loess was determined through coring, trenching, and topographic interpretation, whereas the electronic map was relied upon for distribution of the Peoria and Loveland loesses.

# Soils of Fort Riley Military Base

- Frbase
- Ad
- Be
- Bf
- Bk
- Bop
- Ca
- Cc
- Cf
- Ch
- Cr
- Cs
- Ct
- Dr
- Dw
- Eu
- Ga
- Gc
- Ge
- Gf
- Ha
- He
- Hm
- Hn
- Ho
- Ic
- Id
- Ie
- Iv
- Ka
- Kb
- Ke
- Kf
- Ks
- Lo
- Mk
- Mu
- Oc
- Pt
- Qua
- Ra
- Rd
- Re
- Sa
- Sc
- Sh
- Sm
- Sn
- So
- St
- Tn
- To
- Ts
- Tt
- Tu
- Tv
- Wat
- Wm
- Wn
- Wo
- Wr
- Ws
- w
- xxx



3 0 3 6 Miles

A horizontal scale bar with alternating black and white segments. It is marked with the numbers 3, 0, 3, and 6, representing miles.

Figure 1

## Mapping Alluvial Units

After first mapping alluvial landforms using stereoscopic coverage of black and white aerial photography, a foot survey was conducted of each valley in order to refine the reconnaissance maps. A subsequent foot survey was made of each valley, but this time from the channel bed or water's side to correlate fills exposed in channel banks with their overlying surfaces, and to search for exposed sections worthy of documentation and, in some instances, radiocarbon dating.

## Radiocarbon Dating

Radiocarbon dating has been a valuable tool in late-Quaternary studies for more than 40 years. The numerical age control on stratigraphy provided by radiocarbon dating permits determination of the timing of various erosional and depositional events and affords the opportunity to calculate the rate and magnitude of environmental and geomorphic change. In particular, radiocarbon dating has, in recent years, become critical to geoarchaeological research. Materials for radiocarbon dating are, however, limited in late-Quaternary deposits of the central Great Plains. Late-Pleistocene vegetative cover of the region has produced only scattered wood and charcoal in loess and alluvium, and grass cover of the Holocene resulted in even fewer datable macrofossils. Earth scientists and archaeologists frequently use organic carbon preserved in buried soils and sediments for radiocarbon age control. There is, however, considerable debate about the accuracy of ages determined from buried soils and sediments, despite general acceptance of radiocarbon dating. Some argue that resulting age determinations are fraught with problems (e.g., Polach and Costin, 1971; Geyh et al., 1983; Forman and Miller, 1989; Hammond et al., 1991), while others maintain that ages provide relatively reliable age control (e.g., Matthews and Dresser, 1983; Haas et al., 1986, Martin and Johnson, 1995). Radiocarbon dating of organic carbon in soils and sediments has become routine in recent years and is producing results comparable to those from charcoal and wood. Dating soils and sediments does, however, require greater care in collection, preparation, and interpretation.

## Procedures

Of the four one-gallon bags of material collected for each sample, one was held in reserve and three were prepared for submission using the following procedure:

- 1) disaggregating the sample
  - (a) placing each bag (e.g., one-gallon *Ziploc*) of the sample into a 20-quart stainless steel or aluminum pot; filling the pot with distilled water; disaggregating with a large perforated stainless steel spoon or specially modified electric mixer; covering and allowing the sample to stand overnight or longer for complete disaggregation (some samples required repeated stirring);
- 2) sieving to remove floating debris and sand-size particles:
  - (a) stirring and skimming off floating debris with 60-mesh sieve until clear;
  - (b) passing the sample through a 230-mesh sieve over the sink and discarding the water and sediment (sand and larger);

- 3) transferring the sample to beakers:
  - (a) siphoning the pots and flushing the sediment into the 4-liter beakers using a distilled water jet; filling the beakers to within 2 inches of the top;
- 4) oven drying:
  - (a) siphoning the supernatant from the beakers.
  - (b) covering the beakers with aluminum foil and placing the beakers in the oven at 100° C until the sediment has dried completely, forming a cake;
- 5) pulverizing:
  - (a) removing the dry cake from the beakers and pulverizing.
  - (b) wrapping aluminum foil and sealing in a *Ziploc* or *Whirl-pak* bag

After preparation at the University of Kansas, samples were submitted to the Radiocarbon Laboratory at the Illinois State Geological Survey for conventional radiocarbon dating. There, prior to burning for gas production and counting, the samples are treated by boiling in 2N HCl for one hour in order to remove CaCO<sub>3</sub> and any dolomite that might be present.

## **RESULTS**

### **Loess Distribution**


#### **Loveland and Peoria loesses**

Distribution of the loessal soils indicates that the primary remnant of the upland loess mantle exists in the western and northern part of the base (Fig. 2). Outcrop of the older Loveland loess occurs, however, in the southeastern part, on the eroded, upper slopes of the steep tributaries to the Kansas River valley, e.g., upper Breakneck, Forsyth, and Onemile creeks, as well as along the valley walls at the upper end of Milford Lake. Peoria loess, topographically and stratigraphically above the Loveland, covers a large expanse along the north-south trending drainage divide in the center of the base, an area referred to as the "Bowling Alley." Areas of Peoria loess not shown on the map, due to errors in the field mapping, occur along the bluff of the Kansas River valley, where thicknesses exceed ten meters in places.

#### **Bignell loess**

Determining the distribution of the Bignell loess is difficult in that it has no unique soil series associated with it. This is not surprising in that the Bignell loess is very similar to the Peoria loess and, where thin, it becomes completely involved in surface soil development. A general model for its distribution has emerged, however: the Bignell is typically found on the highest areas of major drainage divides and at scattered locations along the bluff of the Kansas River valley (Fig. 3). The distribution indicated in the northern part of the Bowling Alley has not been ground-truthed due to its inaccessibility (firing range). It has, however, been located by soil coring along the northern and northwestern parts adjacent to the range. Six sites (3 major divides, 3 bluff) have documented the existence and age of the Brady soil and overlying Bignell loess; data from each of these sites is

# Distribution of Loess Soils in Fort Riley Military Base



-  Non-Loess Soils
-  Loveland Loess
-  Peoria Loess
-  Bignell Loess




Figure 2

# Distribution of Bignell Loess in Fort Riley Military Base




Figure 3

available in W.C. Johnson, 1998, and in Table 1.

### **Distribution and Ages of Alluvial Fills**

Alluvial landforms of drainages located along the southern and northeastern periphery of the central firing range have been mapped and in most instances radiocarbon dated (Fig. 4). Alluvial surfaces of Breakneck, Pumphouse, Onemile, Forsyth, and Wildcat creeks have been mapped. Further, radiocarbon age control has been obtained on Pumphouse, Forsyth, lower Threemile and Sevenmile (south of the firing range), and Wildcat creeks, as well as on the Kansas and Republican rivers along the southern boundary of the base (Table 2).

Alluvial units defined on the maps associated with the stream systems discussed below include collectively the flood plain or T0; terrace 1 or T1, which is lowest in elevation; terrace 2 or T2, which is next highest; terrace 3 or T3, which is higher yet; and fans.

#### **Breakneck Creek**

Except for the east branch, Breakneck Creek system exhibits very little flood plain (T0) due to apparent recent entrenchment (Fig. 5). Further, the period of entrenchment and subsequent lateral migration associated with Terrace 1 (T1) was likely of short duration because of the modest amount of surface area created prior to entrenchment to the present flood-plain level. During T1 time the channel did, however, cut into the north side of the distal portion of the upper alluvial fan (Fig. 6). Terrace 2 (T2) clearly dominates the system from the confluence down to where its valley opens onto the Kansas River valley (Fig. 7). Although no radiocarbon ages have yet been obtained from fill in this system, experience elsewhere on Fort Riley indicates that the T2 fill could be up to 10,000 years old, but based on soil development, a maximum estimated age of 5,000 years may be the case in this drainage.

#### **Onemile Creek**

In contrast to Breakneck, Onemile Creek exhibits appreciable T0 throughout the system (Fig. 8). The prominence of flood plain in the upper reaches is typical of most small tributaries and develops because of the flashy, concentrated runoff that occurs in the upper watershed, i.e., these areas are characterized by active channels and frequent flushing of the valley fill. From the upper confluence to the lower valley limit, T1 is regularly distributed, occupying about one-third of the valley floor. The distal portion of the single large fan was apparently eroded during T1 lateral migration. T2 remnants have survived at scattered location along the main valley and into the north branch; most of these elements are at risk due to present channel migration. To facilitate an appreciation of the valley fill in the context of the entire landscape, Onemile Creek alluvial surfaces have been superimposed on the orthophotography (Fig. 9). The distribution of woodland defines the full drainage network of the system and its size relative to the Kansas River. The single alluvial fan is evident below a high-energy tributary, i.e., a drainage large enough to have sufficient runoff and sediment supply to build a fan, but not large enough to have the discharge to reach the channel in the main valley. Ages of the individual fill units are probably temporally related to those of the Breakneck Creek system.

Table 1. Upland (Loessal) Radiocarbon Ages

Site Name (no.)	Depth (cm)	Material Assayed	ISGS Number	Uncorrected Age	$\delta^{13}\text{C}$ (‰)	Corrected Age
Summer Hill						
Trench 2	215	humates	3164	19,020±650	-15.5	19,170±650
	235	humates	3163	21,440±590	-15.1	21,610±590
	255	humates	3160	21,430±740	-15.4	21,580±740
Trench 5	310	humates	3159	20,610±750	-16.0	20,760±750
	330	humates	3158	22,440±992	-17.0	22,560±990
	350	humates	3157	22,850±100	-17.2	23,000±1,100
Trench 6	150	humates	3101	6,000±130	-16.2	6,140±130
	170	humates	3140	7,640±220	-17.1	7,770±220
	200	humates	3142	11,100±310	-16.8	11,230±310
	240	humates	3131	11,840±270	-16.2	11,980±270
	260	humates	3143	13,080±310	-15.9	13,240±310
	280	humates	3146	13,310±430	-15.6	13,470±430
Trench 6a	90	humates	4174	4,740±70	-14.9	4,900±70
	130	humates	3666	5,960±120	-17.2	6,080±120
	170	humates	3665	8,520±140	-17.3	8,650±140
	210	humates	3731	11,300±220	-17.4	11,430±150
	225	humates	3663	15,290±220	-17.2	15,420±220
	256	humates	3750	14,690±240	-17.3	14,810±240
	295	humates	3749	17,320±310	-16.5	17,460±310
	330	humates	3748	17,260±300	-16.0	17,400±300
	344	humates	3747	17,040±300	-15.9	17,180±300
Lookout Hill						
Core 23	100	humates	3010	3,750±160	-16.0	3,890±160

Table 1. Upland (Loessal) Radiocarbon Ages (continued)

Site Name (no.)	Depth (cm)	Material Assayed	ISGS Number	Uncorrected Age	$\delta^{13}\text{C}$ (‰)	Corrected Age
Pumphouse Hill	80	humates	3777	3,200±70	-14.2	3,380±70
	105	humates	3776	4,250±80	-14.9	4,410±80
	130	humates	3165	5,660±180	-15.6	5,810±180
	170	humates	3728	7,210±130	-16.2	7,350±130
	235	humates	3729	12,190±250	-16.6	12,330±250
	265	humates	3730	16,280±220	-21.8	16,330±220
	290	humates	3167	18,770±450	-21.6	18,830±450
	290	humates	3727	21,670±670	-22.2	21,720±670
	330	humates	3166	22,910±850	-18.4	23,010±850
	340	humates	3618	33,210±2700	-18.0	33,300±2,700
Site 27	64	humates	3779	3,300±70	-15.7	3,450±70
	129	humates	3616	9,350±100	-20.5	9,420±100
	190	humates	3615	17,730±320	-21.1	16,790±320
	234	humates	3614	21,180±150	-18.2	21,290±150
Farnum Creek So (site 28)	18	humates	3058	900±130	-15.7	1,050±130
	23	humates	3005	850±100	-15.5	1,000±100
	28	humates	3006	1,570±110	-15.4	1,730±110
	147	humates	3008	14,050±770	-19.0	14,140±770
	172	humates	3055	13,990±470	-17.3	14,120±470
	177	humates	3057	19,540±1800	-17.6	19,700±1,800
	202	humates	3007	14,980±590	-17.3	15,110±590
Bala Cemetery						
	Core 19	humates <sup>1</sup>	2622	18,950±280	-17.3	19,070±280
	Trench	humates	3613	16,300±150	-22.6	16,340±150

Table 1. Upland (Loessal) Radiocarbon Ages (continued)

Site Name (no.)	Depth (cm)	Material Assayed	ISGS Number	Uncorrected Age	$\delta^{13}\text{C}$ (‰)	Corrected Age
Trench (con't)	194	humates	3612	21,170±140	-20.3	21,240±140
	220	humates	3611	25,310±320	-18.3	25,420±320
	253	humates	3610	28,010±280	-17.1	28,140±280

<sup>1</sup> sample not treated with 2N hot HCl

# Fort Riley Military Base Study Stream Systems





Figure 4

Table 2. Valley (Alluvial) Radiocarbon Ages

Site Name (no.)	Depth (cm)	Material Assayed	ISGS Number	Uncorrected Age	$\delta^{13}\text{C}$ (‰)	Corrected Age
KANSAS R.						
Airport Terrace						
Core 20	168	humates <sup>1</sup>	2623	19,900±450	-19.4	19,990±450?
Core 33	30	humates	3056	2,130±140	-15.6	2,280±140
	40	humates <sup>1</sup>	2996	1,910±70	-14.5	2,080±70
	163	humates <sup>1</sup>	2997	11,480±440	-20.4	11,550±440
	193-203a	humates	3053	14,240±480	-17.8	14,360±480
	193-203b	humates <sup>1</sup>	3054	17,690±1000	-17.7	17,800±1,000
	228-233	humates	3003	13,640±540	-18.3	13,750±540
	525-528	humates	3004	11,440±570	-16.5	11,580±570
FR1	90	humates	3650	2,170±70	-15.2	2,330±70
	103	humates	3649	2,240±70	-17.2	2,360±70
	116	humates	3648	1,320±70	-15.2	1,480±70
	136	humates	3662	1,460±70	-15.5	1,610±70
	136	humates	3662 replicate	1,500±70	-15.6	1,650±70
	173	humates	3660	2,740±70	-14.9	2,900±70
	201	humates	3658	3,680±70	-15.6	3,830±70
	275	humates	3601	4,950±80	-17.2	5,070±80
	345	humates	3734	5,950±70	-18.2	6,060±70
FR2	74	humates	3602	2,960±70	-16.0	3,100±70
	101	humates	4175	4,060±70	-16.2	4,200±70
	258	humates	3726	5,870±160	-19.6	5,960±120
	345	humates	4174	5,640±70	-18.2	5,750±70
	450	humates	4173	5,820±70	-18.6	5,920±70

Table 2. Valley (Alluvial) Radiocarbon Ages (continued)

Site Name (no.)	Depth (cm)	Material Assayed	ISGS Number	Uncorrected Age	$\delta^{13}\text{C}$ (‰)	Corrected Age
FR3	290	humates	3812	7,240±70	-17.1	7,370±70
	415	humates	4064	9,400±160	-19.4	9,490±160
FR5	104	humates	3707	1,500±70	-17.2	1,630±70
	390	humates	3657	5,510±80	-18.4	5,620±80
REPUBLICAN R.						
FR10	246	humates	3597	1,380±70	-16.6	1,520±70
	310	humates	3598	1,590±70	-15.7	1,740±70
	424	humates	3599	1,880±70	-16.6	2,010±70
	510	humates	3600	3,790±110	-19.0	3,880±110
WILDCAT CR.						
WC1 (FR7)	85	humates	3780	1,790±70	-16.7	1,920±70
	226	humates	3781	9,830±100	-17.0	9,960±100
	315	humates	3604	17,000±300	-22.7	17,040±300
	434	humates	3603	23,770±300	-19.2	23,860±300
	560	humates	3608	23,780±410	-18.6	23,890±410
	612	humates	3607	23,400±400	-19.2	23,500±400
WC4-T1	149	humates	4001	2,710±70	-16.3	2,850±70
	222	humates	4024	1,850±70	-19.3	1,940±70
WC4-T2	117	humates	4073	3,370±70	-15.9	3,520±70
	175	humates	4000	4,210±70	-18.6	4,310±70
	271	humates	4025	5,440±70	-16.7	5,570±70

Table 2. Valley (Alluvial) Radiocarbon Ages (continued)

Site Name (no.)	Depth (cm)	Material Assayed	ISGS Number	Uncorrected Age	$\delta^{13}\text{C}$ (‰)	Corrected Age
WC4-T3	113	humates	4171	1,260±70	-18.0	1,380±70
WC5-T1	126	humates	3998	1,600±70	-16.4	1,740±70
WC5-T2	322	humates	4072	24,280±150	-16.9	24,410±150
WC5-T6	88	humates	4172	3,480±70	-13.9	3,660±70
	359	humates	4034	10,150±80	-16.0	10,290±80
FORSYTH CR.						
FR4A	86	humates	3856	1,720±70	-15.7	1,870±70
	118	humates	3617	2,850±70	-17.1	2,980±70
	380	humates	3609	6,630±70	-17.7	6,740±70
FR4B	155	humates	3871	8,510±90	-17.0	8,640±90
	167	humates	3860	8,550±70	-17.5	8,680±70
	225	humates	9090 <sup>2</sup>	9,600±70	-19.6	9,690±70
	264	humates	3778	10,600±200	-19.2	10,690±200
	270	humates	3858	10,520±140	-19.8	10,600±140
	320	humates	9091 <sup>2</sup>	10,230±60	-17.4	10,350±60
	375	humates	3857	10,410±120	-17.5	10,530±120 h <sup>3</sup>
	380	bone (Bison sp.)	3605	4,120±70	-10.5	4,350±70
	537	humates	3953	10,520±80	-18.2	10,630±80
FC1-T1	160	humates	4038	1,200±70	-18.7	1,300±70

Table 2. Valley (Alluvial) Radiocarbon Ages (continued)

Site Name (no.)	Depth (cm)	Material Assayed	ISGS Number	Uncorrected Age	$\delta^{13}\text{C}$ (‰)	Corrected Age
FC2-T1	145	humates	4035	2,470±70	-18.2	2,580±70
	259	humates	4042	2,800±70	-16.2	2,950±70
FC3-T1	147	humates	4039	4,770±70	-14.9	4,940±70
FC3-T2	193	humates	4074	8,260±90	-16.7	8,400±90
FC4-T1	140	humates	4066	6,410±90	-15.2	6,570±90
FC5-T1	167	humates	4062	4,860±70	-13.6	5,040±70
	186	humates	4179	5,810±70	-15.0	5,970±70
FC5-T2	100	humates	4061	4,910±70	-14.9	5,080±70
	208	humates	4068	9,310±90	-17.1	9,430±90
	295	humates	4063	12,760±100	-21.0	12,830±100
FC6-T2	120	humates	4065	5,070±70	-14.1	5,250±70
	188	humates	4067	6,770±110	-15.8	6,920±110
	208	charcoal	4057	5,670±70	-25.8	5,660±70
	208	humates	4058	6,880±80	-19.2	6,970±80
	280	humates	4060	10,600±160	-18.0	10,710±160
PUMPHOUSE CR.						
PHC1-T4	161	humates	3997	1,080±70	-17.3	1,210±70
PHC1-T2	258	humates	4026	8,860±110	-15.8	9,010±110

Table 2. Valley (Alluvial) Radiocarbon Ages (continued)

Site Name (no.)	Depth (cm)	Material Assayed	ISGS Number	Uncorrected Age	$\delta^{13}\text{C}$ (‰)	Corrected Age
PHC2-T2	305	humates	4177	2,800±70	-16.5	2,940±70
	376	charcoal & ash	3999	2,450±70	-21.2	2,510±70 h <sup>3</sup>
THREEMILE CR.						
TMC1-T1	307	humates	4059	9,000±90	-15.6	9,150±90
TMC1-T2	130	humates	4056	6,690±80	-17.2	6,820±80
SEVENMILE CR.						
FR13	91	humates	3852	4,540±70	-14.4	4,710±70
	146	humates	3651	6,820±80	-15.5	6,980±80
FR14	380	humates	3716	8,820±160	-19.7	8,910±160 h <sup>3</sup>
FR15	105	humates	3853	4,840±70	-14.7	5,010±70
	306	humates	3712	8,780±130	-17.6	8,900±130 h <sup>3</sup>
FR16	155	humates	3861	2,820±70	-16.3	2,960±70

<sup>1</sup> samples not treated with 2N hot HCl

<sup>2</sup> Tx- (University of Texas Radiocarbon Laboratory)


<sup>3</sup> Age determined from hearth

# Soils of Fort Riley Military Base

Frbase

Frsols

- Ad
- Be
- Bf
- Bk
- Bop
- Ca
- Cc
- Cf
- Ch
- Cr
- Cs
- Ct
- Cx
- Dr
- Dw
- Eu
- Ga
- Gc
- Ge
- Gf
- Ha
- He
- Hm
- Hn
- Ho
- Ic
- Id
- Ie
- Iv
- Ka
- Kb
- Ke
- Kf
- Ks
- Lo
- Mk
- Mu
- Oc
- Pt
- Qua
- Ra
- Rd
- Re
- Sa
- Sc
- Sh
- Sm
- Sn
- So
- St
- Tn
- To
- Ts
- Tt
- Tu
- Tv
- Wat
- Wm
- Wn
- Wo
- Wr
- Ws
- w
- xxx



3 0 3 6 Miles



Figure 1

# Distribution of Loess Soils in Fort Riley Military Base




Figure 2

# Distribution of Bignell Loess in Fort Riley Military Base




Figure 3

# Fort Riley Military Base Study Stream Systems



Figure 4

# Breakneck Creek - Fort Riley Military Base




Figure 5

# Upper Breakneck Creek




Figure 6

# Lower Breakneck Creek




Figure 7

# Onemile Creek - Fort Riley Military Base




Figure 8

# Onemile Creek - Fort Riley Military Base





Figure 9

## **Pumphouse Creek**

This system is characterized by a broad upper valley limited by Moon Lake, and narrow, bedrock-controlled, canyon-like lower valley, both of which are obvious in the width of alluvial fill mapped (Fig. 10). Flood-plain development is modest and uniform throughout the system, indicating an equilibrium longitudinal profile and a relatively short period of flood-plain development. Like the flood plain, the distributions of both terraces are fairly regular throughout the system, with T2 dominating the total valley bottom area. Four small alluvial fans, located above Moon Lake, have relatively steep surfaces and merge abruptly with the terraces and flood plain (Fig. 11). Despite the narrow nature of the lower bedrock valley, a large area of terrace has survived (Fig. 12). Overall, the alluvial surfaces of Pumphouse Creek are extremely well defined.

Three locations within the system were selected for radiocarbon dating (Fig. 10): PHC2-T2 is a cutbank profile exposed in the T2 fill of lower Pumphouse Creek; it revealed three weakly developed buried soils, a hearth, and point bar deposit at the base. Two ages obtained from the site are similar (Table 2): charcoal and ash for the hearth dated to about 2,500 yr B.P., whereas the buried soil in which it was located dated to about 2,900 yr B.P. The disparity is probably due to the latter date being contaminated by redeposition of older carbon in the sediments within which the soil formed. PHC1-T2, also located in T2 fill provided a much older age of about 9,000 yr B.P.; the age is most likely accurate and representative of the fill age. Inspection of the T2 fill in the system indicates that it consisted of two episodes of cutting and filling, the latter of which cut into the 9,000-yr old fill and replaced part of it up to the same level. Most of the T2 fill in the main valley appears, however, to date to 9,000 yr B.P. The site in T1 fill, PHC1-T4, provided an age younger than either T2 fills, which indicates that T1 fill ranges from about 1,500 to 1,000 years in age.

## **Forsyth Creek**

Five different alluvial surfaces appear in the Forsyth drainage, and, with one exception, these are regularly distributed along the main valley (Fig. 13). T0 is developed to a small extent throughout, but T1 and T2 clearly dominate, with four distinct, large areas of coalescing fans. The largest area of T0 is located at the confluence of the two branches forming the main valley, a network position where such channel mobility is anticipated (Fig. 14). The Williston Point area, another valley confluence, still retains large terrace remnants, in particular the triangular-shaped unit of T2 wedged between the main channel and the tributary (Fig. 15). The largest terrace remnants are found in lower Forsyth, at its confluence with Threemile Creek (Fig. 16), where all five alluvial units appear.

Ten sites, all from backhoe trenching, were investigated in the Forsyth Creek valley (Fig. 13). FR 4B, a T2 site at the confluence, produced radiocarbon age ranges about 11,000 to 8,000 yr B.P., whereas FC6-T2, a site in that same area, produced ages from about 11,000 to 5,200 yr B.P. (Table 2). The existence of an older uppermost age at the former site may be due to the stripping ( $\leq 1\text{m}$ ) taking place during development of the power line corridor, i.e., the younger buried soil was mechanically removed. Four T2 sites in the Williston Point area indicated fill ages that range from about 5,000 to 12,000 yr B.P., as does T2 site FC4-T1 upstream. T2 fill ranges in age from over 12,000 to about 5,000 yr B.P. T1 fill was dated at three sites, FR4A, FC1-T1, and FC2-T1, all of which indicate ages less than 5,000 to about 1,000 yr B.P. The older age of 6,740 yr B.P. from FR4A probably reflects contamination from older carbon since this was not a soil that was dated, but rather a silt drape from a flood, which would have transported older, recycled organic (plant-derived) and

# Pumphouse Creek - Fort Riley Military Base




Figure 10

# Moon Lake - Pumphouse Creek




Figure 11

# Lower Pumphouse Creek




Figure 12

# Forsyth Creek - Fort Riley Military Base




Figure 13

# Upper Forsyth Creek




Figure 14

# Williston Point - Forsyth Creek




Figure 15

# Confluence of Forsyth and Threemile Creeks




Figure 16

inorganic (rock-derived) carbon.

### **Wildcat Creek**

The strange, incomplete appearance of the pattern of alluvial fills in Wildcat Creek is a function of adherence to the base boundary extending down the valley axis (Fig. 17). Even with this incomplete mapping, the dominance of the T2 in the main valley is obvious: T0 and T2 are poorly developed throughout the mapped reach (and beyond). The pattern of T1 fill indicates that a limited amount of lateral migration occurred then, prior to entrenchment to the present level. The recent nature of the entrenchment is apparent: although the channel is highly meandering and characterized by actively eroding cutbanks, little of the T1 and adjacent T2 has been removed. Coalescing fans are common, especially in the upper reaches of the main valley (Fig. 18). In fact, fans dominate the Little Arkansas Creek; due to the coarse and active nature of the fans, the channel has, in one area, occupied a straight course for a long time, i.e., no terrace fill has been deposited (Fig. 19). The valley bottom in the lowermost part of the valley appear as a series of stair steps: T0, T1, T2, and fan deposits are all in close proximity to one another and have distinct scarps at their contacts (Fig. 20).

Radiocarbon ages from three different fills provide the greatest temporal range of the study stream systems. Sites WC1 and WC5-T2 are located on alluvial fans, and yielded basal ages in excess of 24,000 yr B.P., yet the uppermost age at WC1 was only about 1,900 yr B.P. Ages from the sites of T2 fill (WC4-T2, WC5-T6) indicate an age range from over 10,000 yr B.P. to as young as 3,000 yr B.P. Accordingly, T1 sites (WC4-T1, WC4-T3, and WC5-T1) ranged from less than 3,000 yr B.P. to about 1,000 yr B.P.

### **Locating Buried Archaeological Remains**

Because the Fort Riley project is geoarchaeologically focused, the ultimate question posed by this and earlier phases of the investigation concerns the preservation of cultural materials, particularly in a buried context. Investigations of cultural material located at or near the land surface on Fort Riley have demonstrated that aboriginal peoples made use of the entire landscape, evidence that was frequently preserved.

Since the archaeological record is concentrated in the last 12,000 years (Table 3), our investigations have centered on deposits of that antiquity. On the uplands, this time period is represented by the Brady soil, developed in terminal Peoria loess, and by overlying Bignell loess, deposited periodically during approximately the last 9,000 years. Consequently, the distribution of such lithostratigraphic and pedostratigraphic units on the uplands dictates where the potential for buried cultural material exists. Aboriginal activity was concentrated in valleys of the central Great Plains because of the relatively rich resource base, and Fort Riley is a prime place to observe this concentration. Alluvial fills range in age from pre-12,000 BP to the present, and most fills contain soils, evidence of stable flood-plain environments.

Through a process of separating the various individual alluvial surfaces from one another, the areas with certain fill age attributes can be appreciated in the context of the potential for burial of remains from particular cultures. This has been done for three systems representing different drainage sizes: Pumphouse, Forsyth, and Wildcat creeks. A synthesis of ages of fill and cultural

# Wildcat Creek - Fort Riley Military Base




Figure 17

# Upper Wildcat Creek




Figure 18

# Little Arkansas Creek - Wildcat Creek




Figure 19

# Lower Wildcat Creek




Figure 20

Table 3. Central Great Plains Cultural Chronology

Paleoindian	12,000 - 9000 BP 10,050 - 7050 BC
Early-Middle Archaic	9000 - 5000 BP 7050 - 3050 BC
Late Archaic	5,000 - 2,000 BP 3050 - 50 BC
Early-Middle Woodland	2,000 - 1,500 BP 50 BC - AD 450
Late Woodland, Late Prehistoric, and Protohistoric	1,500 - 250 BP AD 250 - 1700

associations indicates similarities among the small (Pumphouse), medium (Forsyth) and large (Wildcat) valleys (Table 4). Youngest cultural material (Late Prehistoric to Protohistoric and Historic) is associated with the flood-plain deposits. Because these deposits are being rework so extensively, it is unlikely that any archaeological material will be in situ, however; conversely, cultural material predating the age of the flood plain may be redeposited in the fill, e.g., a bark beater (Late Woodland?) was found in with the gravel lag on a channel point bar in Wildcat Creek. T1 fill in Forsyth and Wildcat creeks is Late Archaic to Late Woodland in age, whereas the T1 fill of Pumphouse Creek appears to have been deposited in a very brief interval, about 500 years. T2 fill clearly dates from the Paleoindian era into the Archaic, with Pumphouse Creek again being a bit different from the other two systems. Only Forsyth Creek has any T3 recognized, and the aerial extent is extremely limited. Because the T2 is inset, the T3 is of presumed pre-Paleoindian age, and any associated archaeological materials would be located within the upper few tens of centimeters. Alluvial fans merit special attention because of their aerial extent in some systems and extreme range in age of their fill, i.e., pre-Paleoindian to Early Woodland, where dated in Wildcat valley.

### **Pumphouse Creek**

Individual depictions of T0, T1, T2, and fan extent (Figs. 21-24) suggest that the majority of the valley fill has the potential to contain buried sites. Late Woodland-age fills are extensive (T1), but the older Paleoindian and Archaic-age fills are most extensive. Cutbank exposures of cultural features, including the hearth dated to the Late Archaic, attest to the presence of buried cultural materials in these fills.

### **Forsyth Creek**

The relative area of flood plain is greater in Forsyth than Pumphouse, but it is still a small

# Pumphouse Creek - Potential Subsurface Cultural Materials

## Late Prehistoric to Protohistoric




Figure 21

# Pumphouse Creek - Potential Subsurface Cultural Materials

## Late Woodland




Figure 22

# Pumphouse Creek - Potential Subsurface Cultural Materials

## Paleoindian to Late Woodland




Figure 23

# Pumphouse Creek - Potential Subsurface Cultural Materials

## Paleoindian to Early Woodland




Figure 24

# Forsyth Creek - Potential Subsurface Cultural Materials

## Late Prehistoric to Protohistoric




Figure 25

Table 4. Buried Cultural Associations and Estimated<sup>1</sup> Threshold Ages<sup>2</sup> for Alluvial Fills

Fill	Pumphouse Creek	Forsyth Creek	Wildcat Creek
T0	<i>Late Prehistoric - Protohistoric (≤1,000)</i>	<i>Late Prehistoric - Protohistoric (≤1,000)</i>	<i>Late Prehistoric - Protohistoric (≤1,000)</i>
T1	<i>Late Woodland  (1,500-1,000)</i>	<i>Late Archaic - Late Woodland (5,000-1,000)</i>	<i>Late Archaic - Late Woodland (3,000-1,000)</i>
T2	<i>Paleoindian - Late Woodland (&gt;9,000-1,500)</i>	<i>Paleoindian - Middle Archaic (13,000-5,000)</i>	<i>Paleoindian - Late Archaic (11,000-3,000)</i>
T3	-	<i>None (all surficial) (&gt;13,000)</i>	-
Fans	<i>Paleoindian - Early Woodland (?)</i>	<i>Paleoindian - Early Woodland (?)</i>	<i>Paleoindian - Early Woodland (24,000-1,800)</i>

<sup>1</sup> from <sup>14</sup>C ages and presumed sedimentation rates

<sup>2</sup> years B.P.

part of the total valley fill (Fig. 25). A sizable area is involved in the T1 and T2 fill distributions, particularly the latter (Figs. 26, 27). The large area of T2 fill at the confluence with Threemile Creek, dating to Paleoindian and Archaic, very likely contains buried cultural material and sites due to its resource-rich location. A similar confluence area in lower Sevenmile Creek, containing a large expanse of T2 fill, produced Early Archaic hearths in three of four backhoe trenches (Table 2). T3 fill should have only surficial cultural material, and that small remnant has been heavily modified by road construction (Fig. 28). Although undated in Forsyth Creek, the alluvial fan fill appears, from color and stratigraphic position, to be ages similar to those in Wildcat Creek where temporal determinations have been made, i.e., pre-Paleoindian to Woodland (Fig. 29).

### Wildcat Creek

Due to recent, region-wide entrenchment, Wildcat Creek has little expression of flood plain (Fig. 30), but older fills abound. T1 surfaces and fills are common throughout, including the Little Arkansas Creek valley (Fig. 31). T2 fills are, however, ubiquitous, implying that a large potential

# Forsyth Creek - Potential Subsurface Cultural Materials

## Late Archaic to Late Woodland




Figure 26

# Forsyth Creek - Potential Subsurface Cultural Materials Paleoindian to Middle Archaic




Figure 27

# Forsyth Creek - Potential Subsurface Cultural Materials

## All Surficial




Figure 28

# Forsyth Creek - Potential Subsurface Cultural Materials

## Paleoindian to Early Woodland




Figure 29

# Wildcat Creek - Potential Subsurface Cultural Materials

## Late Prehistoric to Protohistoric




Figure 30

# Wildcat Creek - Potential Subsurface Cultural Materials

## Late Archaic to Late Woodland




Figure 31

# Wildcat Creek - Potential Subsurface Cultural Materials

## Paleoindian to Late Archaic




Figure 32

# Wildcat Creek - Potential Subsurface Cultural Materials

## Paleoindian to Early Woodland




Figure 33

exists for the presence of buried Paleoindian and Archaic materials in the main valley (Fig. 32). Because of the relatively large extent and age range of alluvial fans in the creek system valley, it is also likely that potential sites are buried in these deposits as well; in fact, the higher and better drained fans may have been preferred surfaces during times of frequent flooding (Fig. 33).

## CONCLUSIONS

A model of culture period-landform associations provides a general framework within which to consider the taphonomy of prehistoric settlement, which to a large degree influences the patterns perceived. This study has dealt with the loess mantle of the uplands and the alluvial surfaces and fills of the valleys. Although much of the cultural record is lost or disturbed, some remains exposed on the surface or within the upper few centimeters, but much more of it is buried, often deeply. Through a combination of age determination, stratigraphic correlation, and landform mapping, the probability of locating buried archaeological remains has been articulated in a cultural-specific fashion.

This phase of the USACERL-funded Fort Riley project represents a continuation of geoarchaeological investigations aimed at producing a high-resolution model of late-Quaternary landscape change and potentials for burial of prehistoric cultural materials. Specific tasks and activities included:

1) *Development of spatial data of dated landform-sediment associations in an ArcView format.* Maps within this report are the representation of the relationships established between the age of sedimentary bodies and their distribution, with a perspective on distribution of prehistoric cultural remains.

2) *Collection of limited field data addressing data gaps for completion of the paleoenvironmental model.* These activities consisted of (a) mapping valley landforms through coring and trenching, (b) mapping Bignell loess through coring, and (c) collecting additional samples for radiocarbon dating.

3) *Completion of the laboratory analyses of limited field data.* Laboratory analyses consisted of (a) mapping landforms with stereoscopic black and white photograph coverage, (b) analyzing soil and sediment cores, and (c) preparing and submitting soil and sediment samples for radiocarbon dating.

The next phase (2/99-1/00) of the research will continue to develop and refine the GIS-based model of landform development and cultural material distribution. Specifically, the tasks will include (1) refinement of the 2-D landscape evolution model, (2) development of an ACCESS database incorporating dig permit data, and (3) preparation of 3-D visualization of landform sediment assemblages and military training disturbances in a GIS environment.

## REFERENCES CITED

- Ager, D.V., 1973, the nature of the stratigraphical record. MacMillan, 114 p.
- Ahlbrandt, T. S., Swinehart, J. B. and Maroney, D. G., 1983, The dynamic Holocene dune fields of the Great Plains and Rocky Mountain Basins, U.S.A., *in* Brookfield M. E. and Ahlbrandt, T. S. (ed.), *Eolian Sediments and Processes*, pp. 379-406. Elsevier, Amsterdam.
- Alden, W.C., and Leighton, M.M., 1917, The Iowan drift, a review of the evidence of the Iowan stage of glaciation: Iowa Geological Survey, v. 26, p. 49-212.
- Andrews, J. T., 1987, The Late Wisconsin glaciation and deglaciation of the Laurentide Ice Sheet, *in* Ruddiman, W. F., and Wright, H. E., Jr., *North America and adjacent oceans during the last deglaciation*. Geological Society of America, *The Geology of North America*, v. K-3, p. 13-37.
- Arbogast, A.F., 1995, Paleoenvironments and desertification on the Great Bend Sand Prairie in Kansas: unpublished Ph.D. dissertation, University of Kansas, Lawrence, 385 p.
- \_\_\_\_\_, and Johnson, W. C., 1994, Evolution of a small basin in the central Great Plains and climatic associations. *Quaternary Research* v. 41, p. 298-305.
- Bard, E., Arnold, M., Fairbanks, R. G., and Hamelin, B., 1993,  $^{230}\text{Th}$ - $^{234}\text{U}$  and  $^{14}\text{C}$  ages obtained by mass spectrometry on corals: *Radiocarbon*, v. 35, p. 191-199.
- Bayne, C.K., Davis, S.N., Howe, W.B., and O'Connor, H.G., 1971, Regional Pleistocene stratigraphy: Kansas Geological Survey, Special Distribution Publication 53, p. 4-8.
- \_\_\_\_\_, and O'Connor, H.G., 1968, Quaternary System, *in* Zeller, D.E., ed., *The Stratigraphic Succession in Kansas*: Kansas Geological Survey Bulletin 189, p. 59-67.
- Bettis, E.A., III and Benn, D.W., 1984, An archaeological and geomorphological survey in the central Des Moines River valley, Iowa: *Plains Anthropologist* 29: 211-227.
- Binford, L.R., 1977, General introduction, *in* Binford, L.R., ed., *For Theory building in archaeology*: Academic Press, p. 1-10.
- Bowman, M.W., 1985, A disparity in the rate of lateral channel cutting between two reaches of the Kansas River: unpublished masters thesis, University of Kansas, 205 p.
- Brice, J. C., 1964, Channel patterns and terraces of the Loup Rivers in Nebraska: U.S. Geological Survey, Professional Paper 422:D, 41 p.
- Bushue, L.J., Fehrenbacher, J.B., and Ray, B.W., 1974, Exhumed paleosols and associated modern

till soils in western Illinois: Soil Science Society of America Proceedings, v. 34, p. 665-669.

Caspall, F.C., 1970, The spatial and temporal variations in loess deposition in northeastern Kansas: unpublished Ph.D. dissertation, University of Kansas, Lawrence, 294 p.

\_\_\_\_\_, 1972, A note on the origin of the Brady paleosol in northeastern Kansas: Proceedings of the Association of American Geographers, v. 4, p. 19-24.

COHMAP members, 1988, Climatic changes of the last 18,000 years: observations and model simulations: Science, v. 241, p. 1043-1052.

Condra, G.E., Reed, E.C., and Gordon, E.D., 1947, Correlation of the Pleistocene deposits of Nebraska: Nebraska Geological Survey Bulletin 15, 71 p.

\_\_\_\_\_, and Reed, G.E., 1950, Correlation of the Pleistocene deposits of Nebraska (Revised): Nebraska Geological Survey Bulletin 15A, 74 p.

Cornwell, K.J., 1987, Geomorphology and soils [Chap. 4], in Adair, M.J., and Brown, K.L., eds., Prehistoric and historic cultural resources of selected sites at Harlan County Lake, Harlan County, Nebraska: U.S. Army Corps of Engineers, Kansas City District, p. 29-46.

Curry, B. B., and Follmer, L. R., 1992, The last interglacial-glacial transition in Illinois: 123-25 ka, in Clark, P. U., and Lea, P. D., eds., The Last Interglacial-Glacial Transition in North America: Geological Society of America Special Paper 270, p. 71-88.

Daniels, R.B., Handy, R.L., and Simonson, G.H., 1960, Dark colored bands in the thick loess in western Iowa: Journal of Geology, v. 67, p. 450-458.

Dansgaard, W., Clausen, H. B., Gundestrup, N., Johnsen, S. J., and Rygner, C., 1985, Dating and climatic interpretation of two deep Greenland ice cores: American Geophysical Union Geophysical Monograph, v. 33, p. 71-76.

Dreeszen, V.H., 1970, The stratigraphic framework of Pleistocene glacial and periglacial deposits in the Central Plains, in Dort, W., Jr., and Jones, J.S., Jr., eds., Pleistocene and Recent Environments of the Central Great Plains: University Press of Kansas, Lawrence, Department of Geology Special Publication 3, p. 9-22.

Feng, Z-D, 1991, Temporal and spatial variations in the loess depositional environment of central Kansas during the past 400,000 years: unpublished Ph.D. dissertation, University of Kansas, 250 p.

\_\_\_\_\_, Johnson, W.C., Sprowl, D.R., and Lu, Y-C., 1994, Loess accumulation and soil formation in central Kansas, United States, during the past 400,000 years: Earth Surface Processes and Landforms, v. 19, p. 55-67.

Follmer, L.R., 1978, The Sangamon soil in its type area: a review, in Quaternary Soils, Mahaney, W.C., ed.: Norwich (England), Geo Abstracts Limited, p. 125-165.

\_\_\_\_\_, 1979, A historical review of the Sangamon soil, in Wisconsinan, Sangamonian, and Illinoian stratigraphy in central Illinois, Illinois State Geological Survey, Guidebook 13, p. 79-91.

\_\_\_\_\_, 1982, The geomorphology of the Sangamon surface: its spatial and temporal attributes, in *Space and time in geomorphology*, Thorn, C.E., ed.: Allen and Unwin, p. 117-146.

\_\_\_\_\_, 1983, Sangamon and Wisconsinan Pedogenesis in the Midwestern United States, in Porter, S.C., ed., *Late-Quaternary Environments of the United States, Volume 1. The Late Pleistocene*: University of Minnesota Press, Minneapolis, p. 138-144.

Forman, S.L., 1990a, Chronologic evidence for multiple episodes of loess deposition during the Wisconsinan and Illinoian in the mid-continent, U.S.A. (abst.): *Geological Society of America Abstracts with Programs*, v. 22, n. 7, p. A86.

\_\_\_\_\_, 1990b, Thermoluminescence and radiocarbon chronology of loess deposition at the Loveland paratype, Iowa, in Bettis, E.A., III, ed., *Holocene Alluvial Stratigraphy and Selected aspects of Quaternary History of Western Iowa*: Iowa City, Midwestern Friends of the Pleistocene, University of Iowa Quaternary Studies Group Contribution, No. 36, p. 165-172.

\_\_\_\_\_, and Miller, G.H., 1989, Radiocarbon dating of terrestrial organic matter, in S.L. Forman, ed., *Dating Methods Applicable to Quaternary Geologic Studies in the Western United States*: Utah Geological and Mineral Survey Miscellaneous Publication 89-7, pp. 2-9.

\_\_\_\_\_, Bettis, E.A., Kemmis, T.L., and Miller, B.B., 1992, Chronological evidence for multiple periods of loess deposition during the late Pleistocene in Missouri and Mississippi River valleys, United States--implications for the activity of the Laurentide Ice Sheet: *Palaeogeography, Palaeoclimatology, Palaeoecology*, v. 93, p.71-83.

Fredlund, G.G., 1995, Late Quaternary pollen record from Cheyenne Bottoms, Kansas: *Quaternary Research*, v. 43, p. 67-79.

\_\_\_\_\_, Johnson, W.C., and Dort, W., Jr., 1985, A preliminary analysis of opal phytoliths from the Eustis ash pit, Frontier County, Nebraska: Nebraska Academy of Sciences, Institute for Tertiary-Quaternary Studies, TER-QUA Symposium Series, v. 1, p. 147-162.

Frye, J.C., and Fent, O.S., 1947, The late Pleistocene loesses of central Kansas: *Kansas Geological Survey Bulletin 70* (1947 rept. stud.), part 3, p. 29-52.

\_\_\_\_\_, and Leonard, A.B., 1949, Pleistocene stratigraphic sequence in northeastern Kansas, *American Journal of Science*, v. 247, p. 883-899.

\_\_\_\_\_, and Leonard, A.B., 1951, Stratigraphy of late Pleistocene loesses of Kansas: *Journal of Geology*, v. 59, no. 4, p. 387-305.

\_\_\_\_\_, and Leonard, A.B., 1952, Pleistocene geology of Kansas: *Kansas Geological Survey Bulletin 99*, 230 p.

\_\_\_\_\_, and Leonard, A.B., 1965, Quaternary of the southern Great Plains, *in* Wright, H.E., Jr. and Frey, D.G., eds., *The Quaternary of the United States—a review volume for VII Congress of the International Association for Quaternary Research*: Princeton University Press, Princeton, N.J., p. 203-216.

\_\_\_\_\_, and Willman, H.B., 1973, Wisconsinan climatic history . interpreted from Lake Michigan lobe deposits and soils; *in* R.F. Black, R.P. Goldthwait, and H.B. Willman, eds., *The Wisconsinan Stage*: Geological Society of America Memoir 136, p. 135-152.

\_\_\_\_\_, Leonard, A.B., Willman, H.B., Glass, H.D., and Follmer, L.R., 1974, The late Woodfordian Jules soil and associated molluscan faunas: Illinois State Geological Survey, Circular 486, 11 p.

\_\_\_\_\_, Wilman, H.B., and Glass, H.D., 1968, Correlation of Midwestern loesses with the glacial succession, *in* Schultz, C.B. and Frye, J.C. eds., *Loess and Related Eolian Deposits of the World: Proceedings 7th Congress, International Association of Quaternary Research*, Boulder-Denver, Colorado, August 14-September 19, 1965, University of Nebraska Press, Lincoln, p. 3-21.

Geyh, M.A., Roeschmann, G., Wijmstra, T.A., and Middeldorp, A.A., 1983, The unreliability of  $^{14}\text{C}$  dates obtained from buried sandy podzols. *Radiocarbon*, v. 25, p. 409-416.

Gladfelter, B.G., 1977, Geoarchaeology—the geomorphologist and archaeology: *American Antiquity* 42: 519-538.

Grüger, J., 1973, Studies on the late Quaternary vegetation history of northeastern Kansas: Geological Society of America, Bulletin, v. 84, p. 237-250.

Guillet, B., Faivre, P., Mariotti, A., and Khobzi, J., 1988., The  $^{13}\text{C}/^{12}\text{C}$  ratios of soil organic matter as a means of studying the past vegetation in intertropical regions: examples from Columbia (South America): *Palaeogeography, Palaeoclimatology, and Palaeoecology*, v. 65, p. 51-58.

Haas, H., V. Holliday, and R. Stuckenrath, 1986, Dating of Holocene stratigraphy with soluble and insoluble organic fractions at the Lubbock Lake archaeological site, Texas: an ideal case study: *Radiocarbon*, v. 28(2A), p. 473-485.

Hall, R.D., 1973, Sedimentation and alteration of loess in southwestern Indiana: unpublished Ph.D. dissertation, Indiana University, 103 p.

Hammond, A.P., Goh, K.M., Tonkin, P.J., and Manning, M.R., 1991, Chemical pretreatments for improving the radiocarbon dates of peat and organic silts in a gley podzol environment: Grahams Terrace, North Westland: *New Zealand Journal of Geology and Geophysics*, v. 34, p. 181-194.

Heinrich, H., 1988, Origin and consequence of cyclic ice rafting in the northeast Atlantic Ocean

during the past 130,000 years: *Quaternary Research*, v. 29, p. 143-152.

Hibbard, C.W., Frye, J.C., and Leonard, A.B., 1944, Reconnaissance of Pleistocene deposits in north-central Kansas: *Kansas Geological Survey Bulletin 52* (1944 rept. stud.), part 1, p. 1-28.

Holien, C.W., 1982, Origin and geomorphic significance of channel-bar gravel of the lower Kansas River: unpublished masters thesis, University of Kansas, 128 p.

Holliday, V.T., 1985, New data on the stratigraphy and pedology of the Clovis and Plainview sites, southern High Plains: *Quaternary Research*, v. 52, p. 388-402.

\_\_\_\_\_, 1989, Middle Holocene drought on the Southern High Plains: *Quaternary Research*, v. 31, p. 74-82.

\_\_\_\_\_, Johnson, E., Haas, H., and Stuckenrath, R., 1983, Radiocarbon ages from the Lubbock Lake site, 1950-1980: Framework for cultural and ecological change on the Southern High Plains: *Plains Anthropologist*, v. 28, p. 165-182.

Hopkins, D.M., 1975, Time-stratigraphic nomenclature for the Holocene Epoch: *Geology*, v. 3, p. 10.

Humphrey, J.D., and Ferring, C.R., 1994, Stable isotopic evidence for latest Pleistocene and Holocene climatic change in North-central Texas: *Quaternary Research*, v. 41, p. 200-213.

Johnson, D.L., 1992, Geomorphological survey and Geoarchaeological overview of Fort Riley, Riley and Geary Counties, Kansas: report to U.S. Army Construction Engineering Laboratory, Champaign, Illinois.

\_\_\_\_\_, 1994, Geoarchaeological research on Fort Riley, Kansas: a dynamic paleoenvironmental model of late Quaternary landscape evolution: report to U.S. Army Construction Engineering Research Laboratory, Champaign, Illinois.

\_\_\_\_\_, 1996, Geomorphological, pedological and geoarchaeological study of Fort Riley, Kansas: Unpublished report to U.S. Army Construction Engineering Research Laboratory, Champaign, Illinois.

Johnson, E., 1987, Paleoenvironmental overview, in *Lubbock Lake: Late Quaternary Studies on the Southern High Plains*, Johnson, E., ed.: Texas A&M University Press, College Station, p. 90-99.

Johnson, W.C., 1989, Stratigraphy and late-Quaternary landscape evolution, in *Archaeological investigations at the North Cove site, Harlan County Lake, Harlan County, Nebraska*, Adair, M.J., ed.: Lawrence, University of Kansas Office of Archaeological Research, p. 22-52.

\_\_\_\_\_, 1993a (ed.), *Second International Paleopedology Symposium Field Excursion: Kansas Geological Survey Open-File Report No. 93-30*.

\_\_\_\_\_, 1993b, Surficial geology and stratigraphy of Phillips County, Kansas, with emphasis on the Quaternary Period: Technical Series 1, Kansas Geological Survey, Lawrence.

\_\_\_\_\_. 1998, Paleoenvironmental processes and archaeology at Fort Riley, Kansas: report to U.S. Army Construction Engineering Research Laboratories.

\_\_\_\_\_, Bozarth, S., and Diekmeyer, E., 1994, Paleoenvironmental reconstruction via opal phytolith and carbon isotope analyses of late-Wisconsinan loess: geoarchaeological investigations on Fort Riley, Riley and Geary Counties, Kansas: Kansas Geological Survey Open-file Report 94-38.

\_\_\_\_\_, and Logan, B., 1990, Geoarchaeology of the Kansas River Basin, central Great Plains, *in* Lasca, N.P., and Donahue, J., eds., *Archaeological Geology of North America: Geological Society of America, Decade of North American Geology Centennial Special Volume 4*, p. 267-299.

\_\_\_\_\_, and Martin, C.W., 1987, Holocene alluvial-stratigraphic studies from Kansas and adjoining states of the east-central Plains, *in* Johnson, W.C., ed., *Quaternary environments of Kansas: Kansas Geological Survey Guidebook Series 5*, p. 109-122.

\_\_\_\_\_, May, D., Diekmeyer, E., Farr, M.R., and Park, K., 1993a, Stop 12 Eustis Ash Pit, *in* W.C. Johnson, ed., *Second International Paleopedology Symposium Field Excursion: Kansas Geological Survey Open-File Report No. 93-30*, p. 12/1-12/13

\_\_\_\_\_, May, D.W., and Mandel, R.D., 1996, A Database of alluvial radiocarbon ages from the central Great Plains (Kansas and Nebraska): *Current Research in the Pleistocene* 13:79-81.

\_\_\_\_\_, and Muhs, D.R., 1999, A revised chronology of loess exposed at the Barton County sanitary landfill, Kansas, based on recent thermoluminescence and radiocarbon ages: U.S. Geological Survey Professional Paper, in review.

Kleiss, H.J. and Fehrenbacher, J.G., 1973, Loess distribution as revealed by mineral variations: *Soil Science Society of America Proceedings*, v. 37, p. 291-95.

Knox, J.C., 1976, Concept of the graded stream, *in* Melhorn, W.N., and Flemal, R.C., eds., *Theories of Landform Development: State University of New York at Binghamton Publications in Geomorphology*, p. 169-198.

\_\_\_\_\_, 1978, Holocene alluvial chronologies and the Great Plains middle Holocene cultural hiatus (abst): *Geological Society of America, Abstracts with Program, North-Central Section*, v. 10, p. 259.

\_\_\_\_\_, 1983, Responses of river systems to Holocene climates, *in* Wright, H.E., Jr., ed., *Late Quaternary Environments of the United States, The Holocene: Minneapolis, University of Minnesota Press*, v. 2, p. 26-41.

Kukla, G.J., 1977, Pleistocene land-sea correlations, I, Europe. *Earth-Science Review*, v. 13, p. 307-374.

- \_\_\_\_\_, 1987, Loess stratigraphy in central China: *Quaternary Science Reviews*, v. 6, p. 191-219.
- Kutzbach, J.E., 1985, Modeling of paleoclimates: *Advances in Geophysics*, v. 28A, p. 159-196.
- Leonard, A.B., 1951, Stratigraphic zonation of the Peoria Loess in Kansas: *Journal of Geology*, v. 59, p. 323-332.
- \_\_\_\_\_, 1952, Illinoian and Wisconsinan molluscan faunas in Kansas: *Kansas University, Paleontological Contributions, Mollusca (Article 4)*, 38 p.
- Leverett, F., 1899, *The Illinois Glacial Lobe: U.S. Geological Survey Monograph 38*.
- Libby, W.F., 1955, *Radiocarbon dating: The University of Chicago Press*, 175 p.
- Lugn, A.L., 1935, The Pleistocene geology of Nebraska. *Bulletin of the Nebraska Geological Survey (2d Series)*, v. 10, p. 1-223.
- Lutenegger, A. J., 1985, Desert loess in the Midcontinent, U.S.A. (abst.): *First International Conference on Geomorphology, Manchester (England), Abstracts of Papers*, p. 378.
- Maat, P., and Johnson, W.C., 1996, Thermoluminescence and new <sup>14</sup>C age estimates for late Quaternary loesses in Southwestern Nebraska: *Geomorphology*, v. 17, p. 115-128.
- Mandel, R.D., 1987, Geomorphology of the Wakarusa River Valley, northeastern Kansas, *in* Logan, B., ed., *Archaeological Investigation in the Clinton Lake Project Area: National Register Evaluation of 27 Prehistoric Sites: U.S Army Corps of Engineers, Kansas City*, p. 20-34.
- \_\_\_\_\_, 1988, Geomorphology of the Smoky Hill River Valley at Kanopolis Lake, *in* Schmits, L.J., ed., *An Archaeological and Geomorphological Survey of Kanopolis Lake, North-Central Kansas: U.S. Army Corps of Engineers, Kansas City*, p. 49-72.
- \_\_\_\_\_, 1991, Geomorphology, *in* Schoen, C.M., ed., *Cultural resource investigations for the U.S. Highway 166 corridor: Kansas State Historical Society, Topeka*, p.8-59.
- \_\_\_\_\_. 1992, Soils and Holocene landscape evolution in central and southwestern Kansas, *in* Holliday, V.T., *Soils in Archaeology: Smithsonian Institution Press, Washington, D.C.*, p. 41-117.
- \_\_\_\_\_, 1993, Geomorphology, *in* Hawley, M.F., ed., *Cultural Resource Investigations for the U.S. Highway 166 Corridor: Kansas State Historical Society*, p. 24-75.
- \_\_\_\_\_, 1994, Holocene landscape evolution in the Pawnee River valley, southwestern Kansas: *Kansas Geological Survey, Kansas Historical Society Bulletin 236*, 117 p.
- \_\_\_\_\_, 1995, Geomorphic controls of the Archaic record in the central United States *in* Bettis, E.A.,

III, ed., *Archaeological Geology of the Archaic Period in North America*: Geological society of America special Paper 297, p. 37-66.

Martin, C.W., 1990, *Late Quaternary Landform Evolution in the Republican River Basin, Nebraska*: unpublished Ph.D. dissertation, University of Kansas, 289 p.

\_\_\_\_\_, 1993, Radiocarbon ages on late Quaternary loess stratigraphy of Nebraska and Kansas, central Great Plains, U.S.A.: *Quaternary Science Reviews*, v. 12, p. 179-188.

\_\_\_\_\_, and Johnson, W.C., 1995, Variation in radiocarbon ages of soil organic matter fractions from late Quaternary buried soils: *Quaternary Research*, v. 43, p. 232-237.

Matthews, J.A., and Dresser, P.Q., 1983, Intensive  $^{14}\text{C}$  dating of a buried paleosol horizon: *Geologiska Foreningens I Stockholm Forhandlingar*, v. 105, p. 59-63.

May, D.W., 1986, *Holocene alluviation, soil genesis, and erosion in the South Loup Valley, Nebraska*: unpublished Ph.D. Dissertation, Department of Geography, University of Wisconsin, Madison.

\_\_\_\_\_, 1989, Age and distribution of the Todd Valley Formation in the lower South Loup River valley (abstract): *Proceeding of the Nebraska Academy of Sciences*, p. 53.

\_\_\_\_\_, 1990, The potential for buried archaeological sites along the Fullerton and Elba canals, North Loup and Loup River valleys, Nebraska: Report submitted to the U.S. Department of the Interior, Bureau of Reclamation, Grand Island, Nebraska, 40 p.

\_\_\_\_\_, 1991, The stratigraphic context of fractured mammoth bones at the La Sena site (25FT177), Harry Strunk Lake (Medicine Creek Reservoir), Nebraska: report prepared for the Department of Anthropology, University of Nebraska, Lincoln, Nebraska, 25 p.

\_\_\_\_\_, 1996, personal communication, Department of Geography, University of Northern Iowa, Cedar Falls.

\_\_\_\_\_, and Holen, S.R., 1993, Radiocarbon ages of soils and charcoal in late-Wisconsinan loess in south-central Nebraska: *Quaternary Research*, v. 39, p. 55-58.

\_\_\_\_\_, and Souders, V.L., 1988, Radiocarbon ages for the Gilman Canyon Formation in Dawson County, Nebraska, (abs.): *Nebraska Academy of Sciences, Proceedings*, p. 47-48.

McCraw, D.J., and Autin, W.J., 1989, *Lower Mississippi Valley loesses*; Field Guide, 1989 Mississippi Valley Loess Tour, INQUA Commission on Loess and the North American Loess Working Group.

McKay, E.D., 1979a, Stratigraphy of Wisconsinan and older loesses in southwestern Illinois, *in* Geology of western Illinois: Illinois State Geological Survey, Guidebook 14, p. 37-67.

\_\_\_\_\_, 1979b, Stratigraphy of Wisconsinan and older loesses in southwestern Illinois, *in* Geology of Western Illinois: Illinois State Geological Survey, Guidebook 14, p. 37-67.

Mix, A.C., 1987, The oxygen-isotope record of glaciation, *in* Ruddiman, W.F., and Wright, H.E., Jr., eds., North America and Adjacent Oceans during the Last Deglaciation: Geological Society of America, The Geology of North America, v. K3, p. 111-135.

Morrison, R.B., 1965, Principles of Quaternary soil stratigraphy, *in* Morrison, R.B., and Wright, H.E., Jr., eds., Quaternary soils, INQUA VII Congress, Proceedings, v. 9, p. 1-69.

\_\_\_\_\_, 1987, Long-term perspective: changing rates and types of Quaternary surficial processes: erosion-deposition-stability cycles, *in* Graf, W.L., ed., Geomorphic systems of North America: Geological Society of America, Decade of North American Geology Centennial Special Volume 2, p. 167-176.

Muhs, D.R., Bettis, E.A., III, Johnson, W.C., Rutledge, E.W., Maat, P.B., and Stafford, T.W., 1996, Pedologic evidence for the climate of the Last Interglacial Period in the Midcontinent of the United States: Geological Society of America Abstracts with Programs, v. 28, p. A-251.

North American Commission on Stratigraphic Nomenclature, 1983, North American Stratigraphic Code: American Association of Petroleum Geologists Bulletin, v. 67, p. 841-875.

Osborn, G., Clapperton, C., Davis, P.T., Reasoner, M., Rodbell, D.T., Seltzer, G.O., and Zielinski, G., 1995, Potential glacial evidence for the Younger Dryas event in the Cordillera of North and South America: Quaternary Science Reviews, v. 14, p. 823-832.

Oviatt, C.G., Karlstrom, E.T., and Ransom, M.D., 1988, Pleistocene loess, buried soils, and thermoluminescence dates in an exposure near Milford Lake, Geary County, Kansas (abst.): Geological Society of America, Abstracts with Programs, v. 20, p. 125-126.

Paterson, W.S.B., and Hammer, C.U., 1987, Ice core and other glaciological data, *in* Ruddiman, W.F., and Wright, H.E., Jr., eds., North America and adjacent oceans during the last deglaciation: Geological Society of America, Decade of North American Geology, The Geology of North American, v. K-3, p. 91-109.

Peteet, D., Rind, D., and Kukla, G., 1992, Wisconsinan ice-sheet initiation: Milankovitch forcing, paleoclimatic data and global climate modeling, *in* Clark, P.U., and Lea, P.D., eds., The Last Interglacial-Glacial Transition in North America: Geological Society of America Special Paper 270, p. 53-69.

Pierce, H.G., 1987, the gastropods, with notes on other invertebrates, *in* Johnson, E., ed., Lubbock Lake: Late Quaternary Studies on the Southern High Plains: Texas A&M University Press, College Station, p. 41-48.

Plog, S., 1976, Relative efficiencies of sampling techniques for archaeological surveys, *in* Flannery, K.V., ed., *the early Maesoamerican village*: Academic Press, p. 136-158.

Polach, H.A., and Costin, A.B., 1971, Validity of soil organic matter radiocarbon dating: buried soils in Snowy Mountains, southeastern Australia as example, *in* D.H. Yaalon, ed., *Paleopedology*: Isreal University Press, pp. 89-96.

Redman, C.L., 1974, Archaeological sampling strategies: Addison-Wesley Module in Anthropology 55, 35 p.

Reed, E.C., and Dreeszen, V.H., 1965, Revision of the classification of the Pleistocene deposits of Nebraska: Nebraska Geological Survey Bulletin 23, 65 p.

Richmond, G.M., and Fullerton, D.S., 1986a, Introduction to Quaternary glaciations in the United States of America, *in* V. Sibrava, D.Q. Bowen, and G.M. Richmond, eds., *Quaternary Glaciations in the Northern Hemisphere: Quaternary Science Reviews*, v. 5, 3-10 p.

\_\_\_\_\_, and Fullerton, D.S., 1986b, Summation of Quaternary glaciations in the United States of America, *in* Sibrava, V., Bowen, D.Q., and Richmond, G.M., eds., *Quaternary glaciations in the Northern Hemisphere: Quaternary Science Reviews*, v. 5, p. 183-196.

Ruddimann, W.F., 1987, Synthesis; the ocean/ice sheet record, *in* Ruddiman, W.L., and Wright, H.E. Jr., eds., *North America and adjacent oceans during the last deglaciation: Geological Society of America, The Geology of North America*, v. K-3, p. 463-478.

Ruhe, R.V., 1956, Geomorphic surfaces and the nature of soils, *Soil Science*, v. 82, p. 441-455.

\_\_\_\_\_, 1969, *Quaternary landscapes in Iowa*: Iowa State University Press, Ames, Iowa, 255 p.

\_\_\_\_\_, 1976, Stratigraphy of midcontinental loess, U.S.A., *in* Mahaney, W.C., ed., *Quaternary Stratigraphy of North America*: Dowden, Hutchinson, and Ross, Stroudsburg, PA, p. 197-211.

\_\_\_\_\_, 1983, Depositional environment of late Wisconsin loess in the Midcontinental United States, *in* Porter, S.C., ed., *Late-Quaternary environments of the United States, Volume 1. The Late Pleistocene*: Minneapolis, University of Minnesota Press, p. 130-137.

\_\_\_\_\_, Miller, G.A., and Vreeken, W.J., 1971, Paleosols, loess sedimentation, and soil stratigraphy, *in* Yaalon, D.H., ed., *Paleopedology--origin, nature, and dating of paleosols*: Hebrew University Press, Jerusalem, Israel, p. 41-60.

\_\_\_\_\_, and Olson, C.G., 1980, Clay-mineral indicators of glacial and nonglacial sources of Wisconsinan loesses in southern Indiana, U.S.A.: *Geoderma*, v. 24, p. 283-97.

Schaetzl, C.B., 1986, The Sangamon paleosol in Brown County, Kansas: Kansas Academy of Science, Transactions, v. 89, p. 152-161.

Shiffer, M.B., 1976, Behavioral archaeology: Academic Press, 222 p.

Schultz, C.D., Lueninghoener, G.C., and Frankforter, W.D., 1951, A graphic resume of the Pleistocene of Nebraska (with notes on the fossil mammalian remains): University of Nebraska State Museum Bulletin, v. 3(6), p. 1-41.

\_\_\_\_\_, and Martin, L.C., 1970, Quaternary mammalian sequence in the central Great Plains, *in* Wakefield Dort, Jr., and J.S. Jones, eds., Pleistocene and Recent Environments of the Central Great Plains: University of Kansas Press, p. 341-353.

\_\_\_\_\_, and Stout, T.M., 1945, Pleistocene loess deposits of Nebraska: American Journal of Science, v. 243, p. 231-244.

\_\_\_\_\_, and Stout, T.M., 1948, Pleistocene mammals and terraces in the Great Plains: Geological Society of America Bulletin, v. 59, p. 553-591.

\_\_\_\_\_, and Tanner, L.G., 1957, Medial Pleistocene fossil vertebrate localities in Nebraska: University of Nebraska State Museum Bulletin, v. 4, p. 59-81.

Simonson, R.W., 1941, Studies of buried soils formed from till in Iowa: Soil Science Society of America, Proceedings, v. 6, p. 373-381.

Souders, V.L., and Kuzila, M.S., 1990, A report on the geology and radiocarbon ages of four superimposed horizons at a site in the Republican River Valley, Franklin County, Nebraska (abst.): Proceedings, Nebraska Academy of Sciences, Lincoln, p. 65.

Thorpe, J., Johnson, W.M., and Reed, E.C., 1951, Some post-Pliocene buried soils of central United States: Journal of Soil Science, v. 2, p. 1-22.

Tien, P.L., 1968, Differentiation of Pleistocene deposits in northeastern Kansas by clay minerals: Clay and Clay Mineralogy, v. 16, p. 99-107.

Warren, R.E., and O'Brien, M.J., 1981, Regional sample stratification; one drainage class technique: Plains Anthropologist 26: 213-227.

Wedel, W. R., 1961, Prehistoric Man on the Great Plains: University of Oklahoma Press.

Welch, J.E., and Hale, J.M., 1987, Pleistocene loess in Kansas; Status, present problems, and future considerations, *in* W.C. Johnson, ed., Quaternary environments of Kansas: Kansas Geological Survey Guidebook Series 5, p. 67-84.

Wells, P.V., and Stewart, J.D., 1987, Cordilleran-boreal taiga and fauna on the central Great Plains of North America, 14,000-18,000 years ago: *The American Midland Naturalist*, v. 118, p. 94-106.

Wendland, W.M., 1982, Geomorphic responses to climatic forcing during the Holocene, *in* Thorn, C.E., ed., *Space and Time in Geomorphology*: London, Allen and Unwin Ltd., p. 355-371.

\_\_\_\_\_, 1978, Holocene Man in the Plains-The ecological setting and climatic background: *Plains Anthropologist*, v. 23, p. 273-287.

\_\_\_\_\_, and Bryson, R.A., 1974, Dating climatic episodes of the Holocene: *Quaternary Research*, v. 4, p. 9-24.

Willman, H.B. and Frye, J.C., 1970, Pleistocene stratigraphy of Illinois: Illinois State Geological Survey, Bulletin 94, 204 p.

Wood, W.R., and Johnson, D.L., 1978, A survey of disturbance processes in archaeological site formation, *in* Schiffer, M.B., ed., *Advances in archaeological method and theory*, v. 1: Academic Press, p. 315-381.