

PRELIMINARY SURFICIAL GEOLOGY OF THE MIAMI COUNTY
PORTION OF THE NEW LANCASTER QUADRANGLE, KANSAS


Geology by Rolfe D. Mandel and K. David Newell
2016

Computer compilation and cartography by John W. Dunham and Sarah F. Child



Open-File Report 2016-13

Funded in part by the
USGS National Cooperative
Geologic Mapping Program



Elevation contours are presented for general reference. Used in the U.S. Geological Survey's current US Topo 1:24,000-scale topographic map series, they were generated from hydrographically-improved 1/3 arc-second National Elevation Dataset (NED) data, and smoothed during processing for use at 1:24,000 scale. In some places the contours may be more generalized than the base data used for compilation of geologic outcrop patterns. Outcrop patterns on the map will typically reflect topographic variations more accurately than the annotated contour lines. Repeated fluctuation of an outcrop line across a contour line should be interpreted as an indication that the mapped rock unit is maintaining a relatively constant elevation along a generalized contour.

Geology was mapped in the field using a USGS 7.5-minute 1:24,000-scale topographic map. Roads and highways are shown on the base map as represented by data from the Kansas Department of Transportation (KDOT), U.S. Census Bureau, and other sources. U.S. Department of Agriculture - Farm Services Agency (USDA-FSA) National Agriculture Imagery Program (NAIP) imagery also was used to check road locations.

Shaded relief is based on 1-meter hydroflattened bare-earth DEMs from the State of Kansas LiDAR Database. The DEM images, in ERDAS IMAGINE format, were mosaicked into a single output DEM, downsampled to 5-meter resolution, and reprojected to decimal degrees. The output DEM was then converted to a hillshade, a multidirectional shaded-relief image using angles of illumination from 0°, 225°, 270°, and 315° azimuths, each 45° above the horizon, with a 4x vertical exaggeration.



This geologic map was funded in part by the USGS National Cooperative Geologic Mapping Program, award number G15AC00225 (FY2015).

This map was produced using the ArcGIS system developed by Esri (Environmental Systems Research Institute, Inc.).

This map is a preliminary product and has had less scientific and cartographic review than the Kansas Geological Survey's M-series geologic maps. KGS does not guarantee this map to be free from errors or inaccuracies and disclaims any responsibility or liability for interpretations made from the map or decisions based thereon.

SUGGESTED REFERENCE TO THE MAP

Mandel, R. D., and Newell, K. D., 2016, Preliminary surficial geology of the Miami County portion of the New Lancaster quadrangle, Kansas: Kansas Geological Survey, Open-File Report 2016-13, scale 1:24,000, unpublished.



GEOLOGIC UNITS
CENOZOIC
Quaternary System

Undifferentiated alluvium — Undifferentiated alluvium that typically is less than 4,000 years old comprises valley fills beneath the modern floodplain of the Marais des Cygnes River and its tributaries. This alluvium consists of clay, silt, sand, and gravel, with the coarsest sediments comprising the lower 3-6 ft (0.9-1.8 m) of the valley fills. The fine-grained floodplain facies are mostly dark gray, gray, dark grayish brown, and grayish brown silty clay, and silty clay loam. These deposits can reach thicknesses greater than 50 ft (15 m) in the Marais des Cygnes River valley.

Alluvial terraces — Alluvium that typically is 4,000 to 11,000 years old comprises valley fills beneath a low terrace (T-1) of the Marais des Cygnes River and its tributaries. This alluvium consists of clay, silt, sand, and gravel, with the coarsest sediments comprising the lower 3-6 ft (0.9-1.8 m) of the terrace fills. The fine-grained alluvium is mostly brown, dark yellowish brown, and yellowish brown silty clay loam. These deposits can reach thicknesses greater than 70 ft (21 m) in the Marais des Cygnes River valley.

Quaternary and Neogene alluvium — Clayey alluvium often interbedded with cherty gravel is common beneath high terraces and on hillslopes in Miami County. The numerical age of these alluvial deposits is unknown. However, based on their position in the landscape, the deposits on the hillslopes probably aggraded during the Neogene (Aber, 1998), and it is likely the high-terrace fills date back to the middle Pleistocene. The lower 3-6 ft (0.9-1.8 m) of the terrace fills and hilltop deposits consist of cherty gravel. The overlying fine-grained alluvium mostly consists of dark gray, gray, dark grayish brown, and pale brown silty clay. The distribution of fine-grained alluvium interbedded with cherty alluvial gravels is indicated by the Kenema soil series, as shown on the Soil Survey of Linn and Miami Counties, Kansas (Penner, 1981). Redoximorphic features, including strong brown and reddish brown mottles and iron and manganese oxide concretions, are common in the upper 3-10 ft (0.9-3 m) of the fine-grained alluvium. Alluvial deposits comprising the fills of high terraces are typically 20-30 ft (6-9 m) thick, but the alluvial deposits on hillslopes generally are less than 15 ft (4.5 m) thick. A veneer of loess that is less than 3 ft (0.9 m) thick often caps the high-terrace and hilltop alluvial deposits.


Colluvial apron and alluvial fan deposits — Deposits of clay, silt, sand, and gravel occur on footslopes and toeslopes in valley landscapes. These deposits mostly accumulated during the early and middle Holocene. The fan deposits are stratified and consist of the terminal Pleistocene, and they comprise both colluvial aprons and alluvial fans. The colluvial aprons consist of massive deposits of poorly sorted sediments that include many angular, bedrock-derived pebbles and cobbles. The alluvial fans formed where small, intermittent streams enter the Marais des Cygnes River valley and the valleys of its major tributaries. The fan deposits are stratified and consist of well-sorted alluvium dominated by brown, dark yellowish brown, and yellowish brown silty clay, silty clay loam, and silt loam. Thin lenses of gravel are common, and buried soils often occur at the top of upward-fining sequences.

PALEOZOIC
Carboniferous System -Pennsylvanian Subsystem
(Descriptions from Miller, 1966)

Kansas City Group

Wyandotte Limestone/Bonner Springs Shale — The uppermost part of the Kansas City Group is composed of the Wyandotte Limestone and the overlying Bonner Springs Shale. The Wyandotte Limestone is composed of five alternating shale and limestone members. From the base upward, these five units are the Frisbie limestone, Quindaro shale, Argentine limestone, Island Creek shale, and Farley limestone. The Wyandotte is a cuesta-forming formation, ranging in thickness from about 10 to 80 ft (3 to 24 m). The shale members are present only locally in Miami County. The absence of the shale members and the similarity in the lithology of the two upper limestone members makes identification of the units difficult. The Frisbie Limestone Member (2.5-3.6 ft; 0.8-1.1 m) is a light olive-gray to light brownish-gray, fine-grained, massive limestone. Locally, thin shales (<0.5 ft; 0.15 m) in thick can be present in the Frisbie. Crinoid stems and small productid brachiopods are the most common fossils and locally Osagia-like forms (coated grains) may be present. The Quindaro Shale Member (locally absent to 3.3 ft; 1 m) is a dusky-yellow to dark yellowish-orange, sandy shale. In places the lower part of this unit is composed of very dark gray shale. The Quindaro is very fossiliferous with Heterocoelia sp., Dielasma sp., and Phricodothyris sp. being the most common forms. Crinoid fragments and bryozoans are also abundant. The Argentine Limestone Member (~35 ft; 11 m) is probably the most persistent member of the Wyandotte. It is lithologically similar to the Farley limestone which is normally the uppermost limestone member of the Wyandotte. Locally the Farley limestone can be missing and the Argentine limestone is the uppermost member. The Argentine is a light olive-gray to grayish-orange, medium-grained, thin-bedded, locally cherty limestone, which weathers into thin fragments. The Argentine has a varied fauna with the brachiopods Composita sp., Echinaria sp., Antiquatonia sp., and Phricodothyris sp. being most common. Enteletes sp. is abundant west and south of Paola. The fusulid Trinitites sp. is common locally. The Island Creek Shale Member (~1.5 ft; 0.5 m) is not well exposed. It is a grayish-orange, clayey shale containing abundant gastropods. The Farley Limestone Member (locally absent to ~15 ft; 4.6 m), where present, is the uppermost member of the Wyandotte Limestone. It is a light olive-gray to pinkish-gray coarse-grained, wavy, thin- to thick-bedded limestone. Locally the lower part of this unit can be oolitic. The Farley limestone contains much the same fauna as the Argentine limestone. The Bonner Springs Shale (0.9-31.5 ft; 0.3-9.6 m) is the uppermost formation of the Kansas City Group. The lower part is composed of pale olive- to light-gray sandy shale which in places is a thin-bedded micaceous siltstone. The upper part is an olive-gray to yellowish-brown clayey shale in the upper part. Varying lithologies can be present near the middle of the unit - unfossiliferous grayish-red clayey shale or medium-gray clayey shale containing carbonaceous streaks, plant rootlets, and leaf impressions, or medium-grained, medium-bedded, calcareous sandstone is found in this interval. In areas where the Bonner Springs is primarily clayey shale, scattered limestone nodules are present, and a thin, argillaceous limestone bed (~1 ft; 0.3 m) is present 2.0 ft (0.6 m) below the top of the formation. The limestone bed is very fossiliferous and contains abundant pelecypods, algae, brachiopods, gastropods, and bryozoan fragments.

Table with 2 columns: EXPLANATION and symbols. Rows include: Boundaries and Locations (Township/range line, Section line, County line), Transportation (U.S. highway, Local road), Geologic Unit Boundaries (Observed contact), Hydrology and Topography (Perennial stream, Intermittent stream, Water body, Water body - manmade shoreline), Elevation contour (100-foot interval, Elevation contour, 20-foot interval, Depression contour, 20-foot interval).



Iola Limestone/Lane Shale — The Iola Limestone (10-12 ft; 3-3.7 m) is composed of two limestone members and an intervening shale member—the lower limestone is the Paola limestone, the middle shale is the Muncie Creek shale, and the upper limestone is the Raytown limestone. The Paola Limestone Member (absent to ~2.5 ft; 0.8 m) is composed of a massive bed of dark-gray to brownish-gray, fine-grained, dense limestone. The contact with the underlying shale is fairly smooth, but the upper surface of the unit is very irregular. Locally, ironstained "worm tubes" extend downward 1 to 3 inches (2.5-7.6 cm) from the upper surface of the Paola. Crinoid stems and small productid brachiopods are the most abundant fossils. Coated grains and bryozoans are common. The Muncie Creek Shale Member (0.5 ft; 0.15 m) is a bluish-gray to dusky-yellow sandy shale that locally has a dark-gray, carbonaceous, fissile facies that contains small (0.4-1.2 inches; 1-2 cm) phosphatic nodules. The Raytown Limestone Member (5-24 ft; 1.5-7 m) is a light olive-gray to light-gray medium- to coarse-grained limestone with numerous silty shale partings and abundant vugs lined with crystalline calcite. The Raytown is medium bedded in the northern part of Miami County but becomes progressively thin bedded toward the southern part. Abundant large productid brachiopods occur in the Raytown, with Echinaria sp. and Lingioproductus sp. being the most common. The Lane Shale (25 ft; 8 m, increasing in thickness westward to 80 ft; 24 m) is lithologically variable. In the western half of the county is an olive-gray to dusky-yellow silty to sandy shale and thin-bedded siltstone. In the eastern half it is an olive-gray to light-gray clayey shale. Thin carbonaceous streaks 1/16 to 1/4-inch (1.6 to 6.4 mm) thick are found locally in the Lane. In east-central Miami County, a thin light-gray, clayey shale containing laminae of reddish-brown calcareous silt occurs just below the overlying Wyandotte Limestone. The Lane is unfossiliferous except for sparse plant remains, and locally, small brachiopods and crinoids are present in the upper few feet of the formation.

Drum Limestone/Chanute Shale — The Drum Limestone (1.7-8 ft; 0.5-2.4 m) consists of a single massive bed of yellowish-gray to reddish-brown fine- to medium-grained cross-bedded limestone that is uniform in lithology and weathers into thin shales. The most characteristic feature of this unit is abundant, small, white crinoid segments scattered throughout the limestone. Other fossils are Neospirifer sp., Murgispirifer sp., and Composita sp., and locally the small sponge Heliospongia sp. The Chanute Shale (8-38 ft; 2.4-12 m) varies greatly in thickness and lithology over short lateral distances. It is yellowish-brown to greenish-yellow sandy to clayey shale, which locally contains sandstone in the lower and middle parts and commonly in the upper part. The Chanute Shale is a relatively unfossiliferous unit except for plant impressions found locally in the sandy layers. The Chanute Shale contains the Thayer coal (0-0.5 ft; 0.15 m), 11 to 15 feet (3.4-4.6 m) above the base of the formation. The Thayer occurs at differing stratigraphic positions in respect to the upper formational boundary, but is absent where the Chanute is thinner than 11 ft (3.4 m).

Dennis Limestone/Cherryvale Shale — The Dennis Limestone (~32 ft; 10 m) is resistant, forming prominent scarps. It is composed of three members, which, in ascending order, are the Canville limestone, Stark shale, and Winterset limestone. The Canville Limestone Member is absent in Miami County, but may be represented locally by a thin, impure, nodular limestone. The Stark Shale Member (0.9-6 ft; 0.3-1.8 m) is represented by a black, fissile to very thin bedded, carbonaceous shale in the lower part. The upper part is medium to greenish-gray sandy shale and is about twice the thickness of the underlying black shale. The Stark is fossiliferous in the upper, sandy part, with chonetid brachiopods being the most common fossil. The lower part is unfossiliferous except for rare inarticulate brachiopods. The Winterset Limestone Member (~29 ft; 9 m), the uppermost member of the Dennis Limestone, is light gray to olive gray, medium grained, medium bedded, dense in the lower part and more fine grained in the middle part. The lower and middle parts of the member characteristically contain shaly partings and abundant chert. The upper part is usually separated from the lower beds by a thin bed of calcareous gray shale (~1.5 ft; 0.5 m). The upper part of the Winterset is dense, fine-grained, massive limestone. It contains an abundance of darkgray chert. Locally, 1 to 3 ft (0.3-0.9 m) of oolitic limestone can be present near the top. The upper part of the Winterset is very fossiliferous and is characterized by numerous large productid brachiopods and locally by abundant Trinitites sp. is also very abundant in the upper part and is usually found in chert. The Cherryvale Shale (~60 ft; 18 m) comprises a shale-dominated unit between the top of the Dennis Limestone and the base of the Drum Limestone. It includes five members, in ascending order, the Fontana shale, the Block limestone, the Wea shale, the Westerville limestone, and the Quivira shale. The Fontana Shale Member (~15 ft; 4.6 m) is greenish-gray to olive-gray sandy shale. Locally a thin, nodular limestone occurs in the lower part. It is relatively unfossiliferous except for sparse chonetid brachiopods in the lower part. The Block Limestone Member (~12 ft; 3.7 m) is the lowermost limestone bed in the Cherryvale Shale, is a bluish-gray to olive-gray medium-grained, thin-bedded fossiliferous limestone with numerous thin, fossiliferous shale partings. Trinitites sp. is the most characteristic fossil, but Murgispirifer sp. and Syringospira sp. also are abundant. The Wea Shale Member (~1.8 ft; 5.5 m) is characteristically olive-gray sandy shale that contains sandstone beds locally near the base. A thin bed of maroon shale may occur near the top. Locally, pyritiferous nodules and limonite concretions occur in the Wea, but plant impressions and carbonaceous partings can be found at its top. In places a very thin coal occurs near the top of the member. The Westerville Limestone Member (absent to ~2 ft; 0.6 m) is poorly exposed, but where present, is composed of conglomeratic limestone containing abundant limonite nodules and quartz sand. It is an lithologically transition to yellowish-gray marly limestone. The Quivira Shale (~4 ft; 1.2 m) has a lower part (0.5-1 ft; 0.15-0.3 m) that is composed of black, carbonaceous, fissile shale or maroon clayey shale. The upper part consists of olive-gray sandy shale. The Quivira is unfossiliferous, except for sparse inarticulate brachiopods and conodonts in the black fissile shale.

Swope Limestone/Galesburg Shale — The Swope Limestone (34 ft; 10 m), the lower part of this map unit, is composed of two limestone members and an intervening shale member. In ascending order, these members are: Middle Creek limestone, Hushpuckney shale, and Bethany Falls limestone. The Middle Creek Limestone Member (1.8-2 ft) is a medium-gray to bluish-gray, fine-grained, dense limestone that is commonly expressed in outcrop as a single massive unit. A thin shale parting can be present approximately in the middle of this unit. The Hushpuckney Shale Member (4-7.5 ft; 1.2-2.3 m) is composed of black fissile shale overlain by grayish to yellow sandy shale. Locally, a thin bed of light olive-gray shale can overlie the black shale. The upper part of the Hushpuckney is fossiliferous, with chonetid and productid brachiopods. The Bethany Falls Limestone Member (13-28 ft; 4-8.5 m) is a light-gray to light brownish-gray, medium-grained, medium- to thick-bedded sandy limestone containing numerous thin shale partings. In most areas the Member can be divided into two distinct parts. The lower part (~12-16 ft; 3.7-4.9 m) contains fusulimids and the upper part (8-14 ft; 4.3 m) is cherty and locally cross-bedded. Pelletal limestone or oolites, or both, may be present at the very top of the Bethany Falls. Where the pelletal or oolitic limestone is present, the upper part locally contains vertical, tubular cavities (3 to 4 feet (0.9-1.2 m) in length and 1 to 2 inches (2.5-10.2 cm) in diameter) containing iron-stained crystalline calcite. The Bethany Falls is a fossiliferous unit in which Westella sp., Derbyella sp., Antiquatonia sp., and chonetid brachiopods are common. Trinitites sp. are found locally in the lower part of the unit. Algae are locally abundant. The Galesburg Shale (4-12 ft; 1.2-3.7 m), which overlies the Swope Limestone, consists of greenish-gray to dusky-yellow, sandy to calcareous, blocky shale. It is characteristically sparsely fossiliferous and poorly exposed.

Hertha Limestone/Ladure Shale — The Hertha Limestone (~8 ft; 2.4 m), the lower part of this map unit, is composed, in ascending order, of the Critzer Limestone Member, Mound City Shale Member, and Sniabar Limestone Member. The Hertha is commonly poorly exposed. The Critzer Limestone Member (~1 ft; 0.3 m) is composed of light olive-gray, medium grained, cherty limestone with abundant crinoid fragments and some bryozoans. The Mound City Shale Member (~0.4ft; 0.1m), if exposed, is present only as a thin, dusky yellow, unfossiliferous sandy shale. The Sniabar Limestone Member (~6 ft; 1.8 m) is a medium-to coarse-grained, medium- to thick-bedded limestone. It is oolitic at the top and contains chert in the lower part. Hematite pebbles and limestone conglomerate are locally present at the base of this unit. Generally over much of the county, the Sniabar Limestone Member is composed of two distinct limestone beds (both of which can be rich with gastropods), separated by a thin bed of shale. The Ladure Shale (~1.2-11.4 ft; 0.4-3.5 m), which overlies the Hertha Limestone, is light-gray to olive-gray and weathers to a yellowish-gray. It is usually sandy and in some localities calcareous, with a thin limestone bed near the middle. It is unfossiliferous, except for plant impressions in the sandy layers.

Pleasanton Group

Tackett Formation — The Tackett Formation of the Pleasanton Group is composed of three units—two shales that are separated by a middle limestone. Approximately 25 ft (7.6 m) of the upper shale, an olive-gray sandy shale locally weathered to a grayish orange, is exposed in Miami County. Lenses of fine-grained sandstone as thick as 25 ft (7.6 m), called "Knobtown sandstone" when penetrated in the subsurface, are locally present. The Tackett Formation in Miami County is primarily mapped wherever there are sparsely-exposed, non-resistant strata below the Hertha Limestone of the Kansas City Group.

REFERENCES

Aber, J. S., 1998, Chert Gravel and Neogene Drainage in East-Central Kansas: Current Research in Earth Science, Bulletin 240, part 3, Kansas Geological Survey, pp. 29-41.
Miller, D. E., 1966, Geology and ground-water resources of Miami County, Kansas: Kansas Geological Survey, Bulletin 181, 66 p. http://www. kgs.ku.edu/General/Geology/Miami/index.html.
Penner, H. L., 1981, Soil Survey of Linn and Miami Counties, Kansas: U.S. Department of Agriculture, Soil Conservation Service, U.S. Government Printing Office, Washington, D.C., 102 p.