



# PRELIMINARY SURFICIAL GEOLOGY OF THE LYONS SE QUADRANGLE, RICE COUNTY, KANSAS

by William C. Johnson, Nathan A. Schlagel, and John W. Dunham  
2017



Open-File Report 2017-22

Funded in part by the  
USGS National Cooperative  
Geologic Mapping Program



Elevation contours are presented for general reference. Used in the U.S. Geological Survey's current US Topo 1:24,000-scale topographic maps, they were generated cartographically from 1-meter resolution digital elevation models (DEMs) derived from lidar and smoothed during processing for use at 1:24,000 scale. In some places, the contours may be more generalized than the base data used for compilation of geologic outcrop patterns. Outcrop patterns on the map will typically reflect topographic variation more accurately than the associated contour lines. Repeated fluctuation of an outcrop line across a contour line should be interpreted as an indication that the mapped rock unit is maintaining a relatively constant elevation along a generally linear outcrop.

1-meter LiDAR hillshades and 1-meter 2010 and 2012 U.S. Department of Agriculture – Farm Services Agency (USDA-FSA) National Agriculture Imagery Program (NAIP) digital imagery were used as references in the digital mapping. USGS 7.5-minute 1:24,000-scale topographic maps, USDA Natural Resources Conservation Service (NRCS) soil surveys, and other geologic maps and bulletins were used to supplement the mapping. Roads and highways are shown on the base map as represented by data from the Kansas Department of Transportation (KDOT), U.S. Census Bureau, and other sources. U.S. Department of Agriculture – Farm Services Agency (USDA-FSA) National Agriculture Imagery Program (NAIP) imagery also was used to check road locations.

Shaded relief is based on 1-meter hydroflattened bare-earth DEMs from the State of Kansas LiDAR Database. The DEM images, in ERDAS IMAGINE format, were mosaicked into a single output DEM, downsampled to 2-meter resolution, and reprojected to decimal degrees. The output DEM was then converted to a hillshade, a multidirectional shaded-relief image using angles of illumination from 0°, 22.5°, 27.0°, and 31.5° azimuths, each 45° above the horizon, with a 4x vertical exaggeration.



This geologic map was funded in part by the USGS National Cooperative Geologic Mapping Program, award number G16AC00195 (FY2016).

This map was produced using the ArcGIS system developed by Esri (Environmental Systems Research Institute, Inc.).

This map is a preliminary product and has had less scientific and cartographic review than the Kansas Geological Survey's M-series geologic maps. KGS does not guarantee this map to be free from errors or inaccuracies and disclaims any responsibility or liability for interpretations made from the map or decisions based thereon.

## SUGGESTED REFERENCE TO THE MAP

Johnson, W. C., Schlagel, N. A., and Dunham, J. W., 2017, Preliminary surficial geology of the Lyons SE quadrangle, Rice County, Kansas: Kansas Geological Survey, Open-File Report 2017-22, scale 1:24,000, unpublished.



QUADRANGLE LOCATION




The following descriptions consist of a compilation of several sources, including field notes and measured sections, data from shallow (<10 m) cores collected during the course of mapping, Kansas Department of Transportation geologic reports, U.S. Department of Agriculture Natural Resources Conservation Service databases, and lithologic and other data in Bayne and Ward (1974).

## GEOLOGIC UNITS

### CENOZOIC

#### Quaternary System

##### Holocene



**Floodplain alluvium** — Floodplain deposits are associated with the Arkansas River and its tributaries and are Holocene in age. These deposits consist of silt, sand, and gravel. In the Arkansas River sand, to gravel-size sediment consists to a large extent of material derived from the Rocky Mountains, whereas the coarse sediment in its tributaries (e.g., Little Arkansas River, Cow Creek, and Little Cow Creek) is derived primarily from local shale and sandstone, with the exception of Rattlesnake Creek, which is dominated by sand-size material derived from reworked Holocene sediments. Core data indicate that the thickness of the floodplain alluvium ranges from 0 to 45 ft (0–14 m) and averages about 25 ft (8 m).

**Aeolian dune and sheet sand** — Aeolian sand deposits occur in scattered bodies on the floodplain (Qds deposit). These dune sediments consist of moderately well-rounded quartz grains with lesser amounts of silt and clay. The primary source of the dune sediments is the Arkansas River channel. Luminescence dating of selected aeolian dunes mantling the Arkansas River alluvium indicates they were last active about 2,200 to 220 years ago, ages which is consistent with reports from Reno County by Haffen et al. (2012). Arbogast (1998) reported an age range of 22,000 to 11,000 years ago for the sandy alluvium underlying the dunes and sand sheets in the Great Bend Sand Prairie, including southwestern Rice County. The dunes are stable with grass or crop cover except for a few isolated areas where livestock and other disturbances have encouraged activation. Dune sand thickness is highly variable but ranges up to 50 ft (15 m).

##### Pleistocene

**Loess** — Undrained in the northern and eastern parts of the county are mantled with loess, which consists of wind-transported silt with minor amounts of clay and very fine sand. The loess is buff in color and calcareous. The age of the loess is primarily late Pleistocene and is recognized as the Peoria Formation throughout the midcontinent; it ranges up to 13 ft (4 m) in thickness. The older, underlying Loveland Formation is a loess unit ranging up to 8 ft (2.5 m) in thickness. Additionally, minor loess deposition occurred in the early and middle Holocene but could not be differentiated from terminal Peoria Formation deposits in Rice County. Loess has a total thickness ranging up to 21 ft (6.5 m).

##### MESOZOIC

#### Cretean System

**Kiowa Formation** — The Kiowa Formation crops out in the eastern third of Rice County, mainly north of a large sand dune area associated with Cow Creek in the southeast corner. A smaller outcrop area exists on the north side of the Arkansas River valley in southwestern Rice County. Originally called the Kiowa shale, the formation consists of a gray to black fissile shale. Thin fossiliferous (marine mollusks) limestone beds and fossil plant debris occur locally. Selenite crystals commonly appear on weathered surfaces, particularly slopes. Lenses of sandstone, often with calcite cementation, occur in and commonly dominate some of the outcrops. Though not expressed in Rice County, a bench-forming sandstone in the upper Kiowa has ripple marks and associated sedimentary structures in addition to calcite concretions. The basal part of the formation contains a zone of siltstone underlain by mudstone and claystone with red mottles. Notably, lenses and concretions of cone-in-cone structures are common in the shale bedding. Maximum exposed thickness of the Kiowa Formation in Rice County is about 110 ft (34 m).

##### PALEOZOIC

#### Permian System

**Harper Sandstone** — The Harper Sandstone is a red argillaceous siltstone and very fine silty sandstone cropping out only in east-central Rice County. The unit is divided into two members: the Chikasla sandstone and the overlying Kingman sandstone, which are separated by a conspicuous 0.9 m thick bed of sandy, white siltstone. The Harper Sandstone is about 220 ft (67 m) thick and attains an exposed thickness of about 65 ft (20 m) in Rice County.

**Stone Corral Formation** — The Stone Corral Formation consists of dolomitic, anhydrite, gypsum, and halite, though only the dolomite persists in outcrops. Dissolution of the other components results in calcite- or gypsum-filled vugs. In places, the dolomite is oysteric. The formation is mainly light gray with occasional red to orange zones or bands. Ripple marks are common. A few exposures of the formation are dominated by a red shale bounded by thin dolomite beds. This formation is an established marker bed in the Permian red-bed sequence of Kansas. In Rice County, it only crops in a narrow band along the upper bluff line of the Little Arkansas River valley in the east-central part of the county. The Stone Corral Formation is up to 100 ft (31 m) thick and occurs up to 12 ft (4 m) thick in outcrops.

**Ninnescah Shale** — The Ninnescah Shale is mainly a red argillitic dolomite, calcareous mudrock, with some thin green shale beds; however, numerous (mostly gray) beds do occur and occasionally contain conchostraceans (clam shrimp). Thin, cross-cutting secondary saline spar gypsum veins may be present. Beds containing rosette-shaped calcareous concretions weather to prominent stratigraphic features. Outcrops occur in the extreme east-central part of Rice County immediately below the Stone Corral Formation along the lower bluffs of the Little Arkansas River valley. The Ninnescah Shale is 300–450 ft (91–137 m) thick, and outcrops are up to 16 ft (5 m) thick.

##### CITED REFERENCES

Arbogast, A. F., 1998, Late Quaternary paleoenvironments and landscape evolution on the Great Bend Sand Prairie: Kansas Geological Survey, Bulletin 242, 74 p.

Bayne, C. K., and Ward, J. R., 1974, Geology and hydrology of Rice County, central Kansas: Kansas Geological Survey, Bulletin 206, pt. 3, 17 p.

Haffen, A. F., Johnson, W. C., Hanson, P. R., Woodburn, T. L., Ludvigson, G. A., and Young, A. R., 2012, Activation history of the Hutchinson dunes in east-central Kansas, USA during the past 2200 years: Aeolian Research 5, p. 9–20.

##### ADDITIONAL SOURCES


Bayne, C. K., Franks, P. C., and Ives, W., Jr., 1971, Geology and ground-water resources of Ellsworth County, central Kansas: Kansas Geological Survey, Bulletin 201, 84 p.

Fent, O. S., 1950, Geology and ground-water resources of Rice County, central Kansas: Kansas Geological Survey, Bulletin 85, 142 p.

Plummer, N. V., and Romary, J. F., 1947, Kansas clay, Dakota Formation: Kansas Geological Survey, Bulletin 67, 241 p.

##### EXPLANATION

Boundaries and Locations	Hydrology	Topography
Township/range line	Perennial stream	Elevation contour (50-foot interval)
Section line	Intermittent stream	Elevation contour (10-foot interval)
	Water body	Depression contour (50-foot interval)
Local road	Water body + manmade shoreline	Depression contour (10-foot interval)
Geologic Unit Boundaries		
Observed contact		



##### RICE COUNTY QUADRANGLES

- |    |    |    |    |    |
|----|----|----|----|----|
| 1  | 2  | 3  | 4  | 5  |
| 6  | 7  | 8  | 9  | 10 |
| 11 | 12 | 13 | 14 | 15 |
| 16 | 17 | 18 | 19 | 20 |